

Notes on the distribution range of *Sedum rubens* L. (Crassulaceae) in continental Portugal

Vasco Silva¹, Gideon F. Smith² & Estrela Figueiredo²

¹Centre for Applied Ecology “Prof. Baeta Neves” (CEABN, InBIO), School of Agriculture, University of Lisbon, Tapada da Ajuda, 1349-017 Lisbon (Portugal)

²Department of Botany, Nelson Mandela University, P.O. Box 77000, Gqeberha [Port Elizabeth], 6031 (South Africa)

Correspondencia

V. Silva

e-mail: silvadavasco@gmail.com

Recibido: 12 octubre 2021

Aceptado: 20 octubre 2021

Publicado on-line: 25 octubre 2021

Editado por: Marta Recio Criado

Abstract

The natural geographical distribution range of *Sedum rubens* in continental Portugal is reassessed. Based on the *Flora iberica* provinces recognised for the country, the occurrence of the species in Ribatejo is confirmed but it is thus far excluded from Beira Alta.

Key words: Beira Alta, chorology, *Flora iberica*, new records, Ribatejo.

Resumen

Notas sobre la distribución geográfica de Sedum rubens L. (Crassulaceae) en Portugal continental

Se reevalúa la distribución geográfica de *Sedum rubens* L. en Portugal continental. Según las provincias de *Flora iberica*, se confirma la presencia de la especie en Ribatejo, pero se excluye, hasta el momento, de Beira Alta.

Palabras clave: Beira Alta, corología, *Flora iberica*, nuevas citas, Ribatejo.

In continental Portugal the family Crassulaceae is represented by 37 species, nothospecies, and infraspecific taxa, of which 23 are indigenous and 14 variously naturalised (Smith & Figueiredo, 2013: 77; Smith & Stephenson, 2015). Of the crassuloid genera represented in Portugal, *Sedum* s.l. is the most diverse, with 18 annual and perennial taxa having been recorded for the country. Although several clades in Crassulaceae subfam. Sempervivoideae have been accorded recognition at the rank of genus, such as *Petrosedum* Grulich, Messerschmid *et al.* (2020) recently proposed that an expanded, ‘super’ *Sedum* genus that contains about 750 species might be the most sensible solution to the taxonomy of the group. This view largely coincides with the treatment of *Sedum* made for the *Flora iberica* Project (Castroviejo & Velayos, 1995; Smith, 2015).

Sedum rubens L. is an annual species which has an extensive natural geographical distribution range in the Iberian Peninsula and Balearic Islands, as well as further north and east in continental and islandic Europe (Stephenson & Stephenson, 1994: 4–5; Castroviejo & Velayos, 1997: 130; ’t Hart & Eggli, 2003: 100–101). As *S. rubens* is virtually indistinguishable from *S. mucizonia* (Ortega) Raym.-Hamet, also an annual species, when plants are not

yet in flower (Stephenson, 1993: 439, 2007: 267; Smith *et al.*, 2016), we here clarify the occurrence of *S. rubens* in continental Portugal (Figure 1).




Figure 1. Delimitation of provinces in Portugal based on *Flora iberica* (<http://www.floraiberica.es>). The provinces where *Sedum rubens* occurs are in darker shade.

Figura 1. Delimitación de las provincias portuguesas según la *Flora iberica* (<http://www.floraiberica.es>). Las provincias donde se encuentra *Sedum rubens* están en un tono más oscuro.

Brotero (1804: 213) was the first to record this species in Portugal indicating that it occurred “near Lisbon and elsewhere in Estremadura, rarely around Coimbra and in the North of Portugal”, a distribution area that reflects its occurrence in the provinces of Beira Litoral and Estremadura established in *Flora iberica*. Still in the 19th century, *S. rubens* was found in the Algarve by the French naturalist E. Bourgeau (GBIF, 2021) and in Trás-os-Montes by Mariz (1889). Afterwards, Coutinho (1913: 280, 1939: 335) gave its distribution as restricted to the centre and south of the country, while Sampaio (1947: 373) indicated the occurrence of the species south of the Mondego River. Franco (1971: 253) extended the distribution range of *S. rubens* northwards to the Douro River basin, based on collections made by L.G. Barbosa and F.J. Garcia in Trás-os-Montes and Alto Douro (Mendonça & Vasconcellos, 1962; GBIF, 2021), and also recorded the species from the central and southeastern parts of the country (respectively C and SE, Figure 1 in Franco, 1971; see Stephenson 1991: 16 for a map). The southeastern record was probably based on Fernandes & Fernandes (1948) who made collections in Alto Alentejo. Fernandes (1952) presented a record for the north of Portugal, specifically for the region of Porto in the province of Douro Litoral, and later confirmed the occurrence of *S. rubens* in the north by recording a new location in Trás-os-Montes (Fernandes, 1960). Several collections were given by Malato Beliz *et al.* in Alentejo, including Baixo Alentejo (GBIF, 2021), while for Beira Baixa, apart from the reference in Fernandes (1952), there are recent occurrence references in Cano *et al.* (2007). Lastly, the occurrence of *S. rubens* in the province of Beira Alta was recorded by Castroviejo & Velayos (1997) but without specimen citations. This is likely based on specimens collected by Barbosa and Garcia at Almendra (near túnel de Almendra, Vila Nova de Foz Côa, 29TPF6844, 01-V-1946, LISI 11530/1999) (Mendonça & Vasconcellos, 1962; GBIF, 2021). However, according to the provinces of *Flora iberica*, this municipality belongs to the Trás-os-Montes province, not to Beira Alta (see Anthos, 2021).

Although several places where *S. rubens* has been observed and collected in central Portugal fall in the province of Estremadura, no specimens were cited from the adjacent province of Ribatejo (GBIF, 2021). *Sedum rubens* was recorded for Ribatejo by Smith & Figueiredo (2013: 86) and the occurrence is confirmed here with an herbarium sheet (LISI 32811) and images (Figures 2 and 3) which were taken by one of us (VS) during fieldwork at Serras de Aire e Candeeiros Natural Park, in 2020. Several individuals were detected dispersed on walls and rocks in dry places, which confirms the ecology of the species as noted by Coutinho (1939), a character species of the

class *Tuberarietea guttatae* Rivas Goday & Rivas-Martínez 1963 nom. mut. (Costa *et al.*, 2010).



Figure 2. *Sedum rubens* growing near Valverde (Ribatejo) in central Portugal. Photo: V. Silva.

Figura 2. *Sedum rubens* cerca de Valverde (Ribatejo) en el centro de Portugal. Foto: V. Silva

Sedum rubens L. in Sp. Pl.: 432 (1753)

PORTUGAL. Ribatejo: Rio Maior, Alcobertas, Casais Monizes, 39°26'30.1"N 8°53'18.3"W, 325 m, V. Silva & A.P. Paes, 19-V-2020 (LISI 32811).

In addition, observations have been made in the municipalities of Rio Maior, Azambuja, and Coruche of that province, and recorded on the Flora-On portal (Porto *et al.*, 2021).

The distribution of *S. rubens* is now clarified, extending to all Portuguese provinces, with the exception of Minho and Beira Alta, following the delimitation of *Flora iberica*.




Figure 3. *Sedum rubens* growing near Casais Monizes (Ribatejo) in central Portugal. Photo: V. Silva.

Figura 3. *Sedum rubens* cerca de Casais Monizes (Ribatejo) en el centro de Portugal. Foto: V. Silva.

Acknowledgements

We thank the reviewer for valuable comments and suggestions, which helped improve the manuscript.

References

- Anthos (2021). *Sistema de información sobre las plantas de España*. Real Jardín Botánico, CSIC-Fundación Biodiversidad. Available at: www.anthos.es (Accessed: 10 August 2021).
- Brotero, F.A. (1804). *Flora Lusitanica, seu plantarum, quae in Lusitania vel sponte crescunt, vel frequentius coluntur, ex florum praesertim sexibus systematice distributarum, synopsis*. 2 vols., 18 pl. + 607 pp. and 558 pp. Lisbon: Typographia Regia.
<https://catalog.hathitrust.org/Record/100427823>
- Cano, E., Rodríguez-Torres, A., Pinto Gomes, C., García-Fuentes, A., Torres, J.A., Salazar, C., Ruiz-Valenzuela, L., Cano-Ortiz, A. & Montilla, R.J. (2007). Analysis of the *Juniperus oxycedrus* L. communities in the centre and south of the Iberian peninsula (Spain and Portugal). *Acta Botanica Gallica*, 154(1), 79–99.
<https://doi.org/10.1080/12538078.2007.10516046>
- Castroviejo, S. & Velayos, M. (1995). Notas y comentarios sobre el género *Sedum* L. (Crassulaceae) y su tratamiento para Flora iberica. *Anales del Jardín Botánico de Madrid*, 53(2), 271–279.
- Castroviejo, S. & Velayos, M. (1997). *Sedum* L. In: Castroviejo, S. et al. (Eds.), *Flora iberica* 5, 121–153. Madrid: Real Jardín Botánico and CSIC.
- Costa J.C, Espírito-Santo D. & Arsénio P. (2010). Guia geobotânico da excursão ao Parque Natural das Serras de Aire e Candeeiros. *Quercetea*, 10, 5–106.
- Coutinho, A.X.P. (1913). *A Flora de Portugal (Plantas Vasculares): Disposta em Chaves Dichotomicas*. Lisbon: Aillaud, Alves & Cia.
- Coutinho, A.X.P. (1939). *Flora de Portugal*, 2.ª Ed. Lisbon: Irmãos Bertrand, Lda.
- Fernandes, A. & Fernandes, R.B. (1948). Herborizações nos domínios da Fundação da Casa de Bragança. II – Vila Viçosa. *Boletim da Sociedade Broteriana*, Sér. 2, 22, 17–96.
- Fernandes, R.B. (1952). Notas sobre a Flora de Portugal. III. *Anuário da Sociedade Broteriana*, 18, 9–29.
- Fernandes, R.B. (1960). Notas sobre a Flora de Portugal. VIII. *Boletim da Sociedade Broteriana*, Sér. 2, 34, 99–155.
- Franco, J.A. (1971). *Nova Flora de Portugal (Continente e Açores)*, vol. I *Lycopodiaceae—Umbelliferae*. Lisbon: Sociedade Astória Lda.
- GBIF (2021) GBIF.org. GBIF Occurrence Download. Available at: <https://doi.org/10.15468/dl.dspcn> (Accessed: 10 August 2021).
- Mariz, J. de (1889). Uma excursão botânica em Traz os Montes. *Boletim da Sociedade Broteriana*, 7, 3–34.
- Mendonça, F.A. & Vasconcellos, J.C. (1962). *Estudo Fitogeográfico da Região Duriense IV*. Anais do Instituto do Vinho do Porto [1960]. Porto: Edição do Instituto do Vinho do Porto.
- Messerschmid, T.F.E., Klein, J.T., Kadereit, G. & Kadereit, J.W. (2020). Linnaeus's folly—phylogeny, evolution and classification of *Sedum* (Crassulaceae) and Crassulaceae subfamily Sempervivoideae. *Taxon*, 69(5), 892–926.
<https://doi.org/10.1002/tax.12316>
- Porto, M., Pereira, A.J., Carapeto, A., Clemente, A., Araújo, P.V., Covelo, F., Almeida, J.D., Amado, A. & al. (2021). *Sedum rubens* L. - mapa de distribuição. Flora-On: Flora de Portugal Interactiva, Sociedade Portuguesa de Botânica. Available at:
<http://www.flora-on.pt/#wSedum+rubens> (Accessed: 10 August 2021).
- Sampaio, G. (1947). *Flora Portuguesa*. 4th facsimile edn (1990) of the 2nd revised edn by A. Pires de Lima. Lisbon: Instituto Nacional de Investigação Científica.
- Smith, G.F. (2015). *Sedum xestrelae* (Gideon F.Sm. & R.Stephenson) Gideon F.Sm. (Crassulaceae): a new combination for a natural hybrid in *Sedum* series *Rupestria* A.Berger (=*Petrosedum* Grulich) from Portugal. *Haseltonia*, 21, 95–96.
<https://doi.org/10.2985/026.021.0114>
- Smith, G.F. & Figueiredo, E. (2013). The family Crassulaceae in continental Portugal. *Bradleya*, 31, 76–88.
<https://doi.org/10.25223/brad.n31.2013.a10>
- Smith, G.F., Figueiredo, E. & Silva, V. (2016). Notes on the geographical distribution range of *Sedum mucizonia* (Ortega) Raym.-Hamet (Crassulaceae), a miniature, annual succulent, in continental Portugal. *Bradleya*, 34, 133–141.
<https://doi.org/10.25223/brad.n34.2016.a3>
- Smith, G.F. & Stephenson, R. (2015). *Petrosedum xestrelae* Gideon F.Sm. & R.Stephenson (Crassulaceae) [*Petrosedum forsterianum* (Sm.) Grulich × *Petrosedum sediforme* (Jacq.) Grulich]: a new nothospecies from central Portugal. *Bradleya*, 33, 2–8.
<https://doi.org/10.25223/brad.n33.2015.a2>
- Stephenson, R. (1991). Sedum hunting in southern Portugal. *Sedum Society Newsletter*, 18, 16–18.
- Stephenson, R. (1993). Alpine stonecrops from Portugal. *Bulletin of the Alpine Garden Society*, 61, 434–439.

Stephenson, R. (2007). Portuguese succulents. *Cactus & Succulent Journal (U.S.)*, 79(6), 265–268.
[https://doi.org/10.2985/0007-9367\(2007\)79\[265:PS\]2.0.CO;2](https://doi.org/10.2985/0007-9367(2007)79[265:PS]2.0.CO;2)

Stephenson, R. & Stephenson, J. (1994). Andalucian break, *Sedum Society Newsletter*, 30, 4–10.
't Hart, H. & Eggli, U. (2003). *Sedums of Europe—stonecrops and wallpeppers*. Lisse, Abingdon, Exton (PA), Tokyo: A.A. Balkema Publishers.