

A new species of *Clinopodium* (Lamiaceae) from E Cuba

Authors: Santos, Isidro E. Méndez, and Téllez, Julio C. Rifá

Source: *Willdenowia*, 47(2) : 173-177

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.47.47209>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ISIDRO E. MÉNDEZ SANTOS^{1*} & JULIO C. RIFÁ TÉLLEZ¹

A new species of *Clinopodium* (Lamiaceae) from E Cuba

Version of record first published online on 1 September 2017 ahead of inclusion in August 2017 issue.

Abstract: *Clinopodium rankiniae*, a new species from the SE region of Cuba, is described and illustrated. It appears to be closely related to the other endemic species of Cuba and Hispaniola (*C. alpestre*, *C. banaoense*, *C. bucheri* and *C. ekmanianum*), which grow isolated in mountainous ecosystems. It differs primarily by the disposition of the leaves (mostly on short, lateral brachyblasts), the colour of the corolla (white, with a very light purple tinge) and the hirsute to tomentose calyx.

Key words: Caribbean, *Clinopodium*, Cuba, endemism, *Lamiaceae*, *Lamiales*, new species, taxonomy, West Indies

Article history: Received 5 April 2017; peer-review completed 14 June 2017; received in revised form 4 July 2017; accepted for publication 8 July 2017.

Citation: Méndez Santos I. E. & Rifá Téllez J. C. 2017: A new species of *Clinopodium* (Lamiaceae) from E Cuba [Novitiae florae cubensis No. 52]. – Willdenowia 47: 173–177. doi: <https://doi.org/10.3372/wi.47.47209>

The genus *Clinopodium* L. (Lamiaceae), sensu Wagstaff & al. (1995), Cantino & Wagstaff (1998) and Harley & Granda Paucar (2000), comprises about 100 species, most in the New World and only some in Eurasia and Africa (Harley & al. 2004). In Cuba, six species of *Clinopodium* were known (Pool 2008; Greuter & al. 2016), which have been considered indistinctly as *Micromeria* Benth. (Grisebach 1866; Sauvalle 1873; Gómez de la Maza 1897; Urban 1919; Britton & Wilson 1922; Borhidi 1981) or *Satureja* L. (Urban 1924a, 1924b; Alain 1956, 1957; Méndez Santos & al. 2005).

While reviewing the genus *Clinopodium* for the *Flora de la República de Cuba*, we found a gathering in B, HAJB and JE collected in 1989 in Santa María del Loreto (Sierra de Los Ciegos), Santiago de Cuba province, with

characters that differ from all the species hitherto described. Studies of this natural population and comparison with similar taxa from the Greater Antilles, described by Urban (1919), Britton & Wilson (1922), Alain (1968) and Méndez Santos & al. (2005), lead us to propose a new species.

Clinopodium rankiniae I. E. Méndez, **sp. nov.** – Fig. 1, 2. Holotype: Cuba, Provincia Santiago de Cuba, altiplanicie de Santa María del Loreto, finca Los Monieles, al este de la subestación eléctrica con paneles fotovoltaicos, vegetación secundaria sobre suelos derivados de areniscas y conglomerados, al borde de los farallones, 1 Oct 2016, I. E. Méndez, J. C. Rifá & C. Regalado 12016 (HIPC; isotypes: B, HAC, HAJB, JE, PAL-Gr, ULV).

¹ Centro de Estudios de Gestión Ambiental, Universidad “Ignacio Agramonte Loynaz”, CP-74670, Camagüey, Cuba; *e-mail: isidro.mendez@reduc.edu.cu, imendezs1958@gmail.com (author for correspondence); julio.rifa@reduc.edu.cu



Fig. 1. *Clinopodium rankiniae* – holotype (I. E. Méndez & al. 12016, HIPC).




Fig. 2. *Clinopodium rankiniae* – A: lateral brachyblast; B: disposition of leaf and flower; C: flower; D: calyx; E: adaxial surface of leaf blade; F: abaxial surface of leaf blade. – All from live plants from the type locality.

Description — *Shrubs* erect, to 2 m tall, usually much branched. *Twigs* quadrangular, densely covered with a pale tomentose indumentum of tubercle-shaped, bulbous-based, stellate and/or branched hairs (also found on petioles, leaf blades, pedicels, bracteoles and calyxes). *Leaves* disposed on short, lateral brachyblasts; *petiole* c. 1 mm long; *leaf blade* obovate to oblanceolate, 3–10 × 2–5 mm, coriaceous, adaxial surface dark green, veins scarcely sunken, abaxial surface somewhat lighter, veins prominent, both surfaces hirsute-tomentose, more densely so abaxially, base acute to decurrent toward petiole, margin entire, revolute, apex acute or sometimes obtuse. *Flowers* solitary, inserted in leaf axils on short, lateral brachyblasts; *pedicel* c. 1 mm long; *bracteoles* lanceolate to filiform, c. 2 mm long. *Calyx* narrowly campanulate or cylindric, 2-labiate; tube straight, 2–2.5 mm long, 10–12-nerved, hirsute to tomentose outside; upper lip oblong, 3-dentate, teeth acute, central one lightly longer; lower lip 2-dentate, teeth acute to lanceolate, 1-nerved. *Corolla* white, with a very light purple tinge, funnel-shaped, 2-labiate; tube straight, c. 4.5 mm long; upper lip emarginate; lower lip 3-lobate, lobes rounded to spatulate, c. 1.5 mm long. *Stamens* 4, c. 1.5 mm long above insertion point in corolla, glabrous, anterior pair slightly exserted from corolla tube. *Ovary* elongate, glabrous; *style* exserted. *Nutlets* elongate, c. 1 mm long, puberulous only in distal third.

Distribution and ecology — *Clinopodium rankiniae* is known only from the Santa María del Loreto plateau, municipality of La Maya, province of Santiago de Cuba. It grows on a substratum originating from gritty rocks formed by granitic sediments. At this locality, the average annual precipitation is 1200–1400 mm. Associated species include: *Bellonia spinosa* Sw., *Calycogonium rhamnoides* Naudin, *Casearia hirsuta* Sw., *Citharexylum tristachyum* Turcz., *Guapira obtusata* (Jacq.) Little, *Ipomoea subrevoluta* Choisy, *Malpighia suberosa* Small and *Plumeria filifolia* Griseb., among others.

Eponymy — The epithet is in honour of Rosa Rankin Rodríguez, Cuban botanist, co-editor of the *Flora de la República de Cuba* and member of the team that first gathered the species in 1989.

Affinities — *Clinopodium rankiniae* is similar to the other four endemic species of Cuba and Hispaniola (Haiti: *C. ekmanianum* (Epling & Alain) Harley; Dominican Republic: *C. alpestre* (Urb.) Harley; Cuba: *C. banaoense* (P. Herrera & al.) Melnikov and *C. bucheri* (P. Wilson) Harley). All of them are typical of mountainous ecosystems, characterized by a peculiar indumentum on the branches, leaves and inflorescences, with tubercle-shaped, bulbous-based, stellate and/or branched hairs. Within this group, *C. rankiniae* is more similar to *C. alpestre* and *C. ekmanianum* by its leaves being ± compacted on short, lateral brachyblasts and the colour of the corolla being basically white. *Clinopodium ekmanianum* and *C. rankiniae* differ from *C. alpestre* by the obovate to oblanceolate (instead of triangular) leaf blade with a revolute (instead of ± flat) margin. *Clinopodium rankiniae* is finally distinguished by the hirsute to tomentose calyx and a very light purple tinge of the corolla. Besides, *C. rankiniae* occurs distant and completely isolated from the species of Hispaniola, to which it is most morphologically similar, and occurs under markedly different geological and edaphic conditions (Cuba: substratum originating from gritty rocks formed by granitic sediments; Hispaniola: soils derived from calcareous rocks).

Melnikov (2014) classified all these species as belonging to *Clinopodium* subg. *Xenopoma* (Willd.) Melnikov

in Novosti Sist. Vysš. Rast. 45: 148. 2014 [= *Xenopoma* Willd. in Mag. Neuesten Entdeck. Gesammten Naturk. Ges. Naturf. Freunde Berlin 5: 399. 1811 = *Micromeria* sect. *Xenopoma* (Willd.) Benth. in Candolle, Prodr. 12: 222. 1848 = *Satureja* sect. *Xenopoma* (Willd.) Briq. in Engler & Prantl, Nat. Pflanzenfam. 4(3a): 300. 1895].

Conservation status — *Clinopodium rankiniae* is at present considered a very rare species. So far, it is only documented in one locality with a total of fewer than 50 mature individuals. The extent of occurrence is smaller than 100 km² and the area of occupancy is smaller than 10 km² with proven reduction since 1989. The species can therefore be provisionally assessed as Critically Endangered CR B1ab(ii,iii,v)+2ab(ii,iii,v); C2a(i); D according to the IUCN Red List categories and criteria (IUCN 2012).

Additional specimens seen — CUBA: PROVINCIA SANTIAGO DE CUBA: altiplanicie de Santa María del Loreto, finca Los Monieles, al este de la subestación eléctrica con paneles fotovoltaicos, vegetación secundaria sobre suelos derivados de areniscas y conglomerados, al borde de los farallones, 10 May 2016, C. Regalado 12002 (HIPC); Municipio Santiago de Cuba [in error; correctly: Mun. Songo – La Maya], altiplanicie de Santa María del Loreto [Sierra de Los Ciegos], Alto de la Torre, vegetación secundaria sobre caliza [in error; correctly: sobre suelos derivados de areniscas y conglomerados], 3 May 1989, Gutiérrez & al. 67956 (B 100363784, HAJB, JE).

Acknowledgements

The authors thank César Regalado Díaz, who explored the Santa María del Loreto region unceasingly until locating the new species and who provided invaluable help in the realization of the field studies, and the Moniele family for their support in studying the new species inside their property. We are also grateful to the Botanic Garden and Botanical Museum Berlin for liberal access to the research facilities of that institution and various kinds of help through the collaborative Flora de Cuba programme; to the Verein der Freunde des Botanischen Gartens und Botanischen Museums Berlin-Dahlem e.V. for financial support; and to the directors and curators of many herbaria (B, BM, F, G, GH, HAC, HAJB, JE, MO, NY, S and US) for lending material for the purpose of our study. Dr Rosa Rankin Rodríguez, Prof. Werner Greuter and three anonymous reviewers are acknowledged for their advice on, and critical revision of, earlier versions of this paper.

References

Alain [bro.] 1956: Novedades en la flora cubana (VII). – Contr. Ocas. Mus. Hist. Nat. Colegio “De La Salle” **15**: 6–23.

- Alain [bro.] 1957: Flora de Cuba, 4. Dicotiledoneas: *Melastomataceae* a *Plantaginaceae*. – Contr. Ocas. Mus. Hist. Nat. Colegio “De La Salle” **16**.
- Alain [Liogier A. H.] 1968: Novitates antillanae III. – Brittonia **20**: 148–161.
- Borhidi A. 1981 [“1980”]: New names and new species in the flora of Cuba, II. – Acta Bot. Acad. Sci. Hung. **26**: 255–275.
- Britton N. & Wilson P. 1922: Notes on plants collected by Mr. Bucher on Pico Turquino, Cuba. – J. New York Bot. Gard. **23**: 91–94.
- Cantino P. D. & Wagstaff S. J. 1998: A reexamination of North American *Satureja* s.l. (*Lamiaceae*) in light of molecular evidence. – Brittonia **50**: 63–70.
- Gómez de la Maza M. 1897: Flora Habanera. Fanerógamas. – Habana: Sellos de Goma, “La Moderna Poesía”.
- Greuter W., Rankin Rodríguez R. & Méndez Santos I. E. 2016: *Lamiaceae*. – Pp. 185–193 in: Greuter W. & Rankin Rodríguez R. 2016: Espermatófitos de Cuba: inventario preliminar. Parte II: Inventario. The *Spermatophyta* of Cuba: a preliminary checklist. Part II: Checklist. – Berlin: Botanischer Garten & Botanisches Museum Berlin-Dahlem; La Habana: Jardín Botánico Nacional, Universidad de La Habana. – Published at: <http://dx.doi.org/10.3372/cubalist.2016.2>
- Grisebach A. 1866: Catalogus plantarum cubensium exhibens collectionem Wrightianam aliasque minores ex insula Cuba missas. – Lipsiae: apud Guilielmum Engelmann.
- Harley R. M., Atkins S., Budantsev A. L., Cantino P. D., Conn B. J., Grayer R., Harley M. M., Kok R. de, Krestovskaja T., Morales R., Paton A. J., Ryding O. & Upson T. 2004: *Labiatae*. – Pp. 167–275 in: Kadereit J. W. (ed.), The families and genera of vascular plants **7**. – Berlin, Heidelberg, New York: Springer.
- Harley R. & Granda Paucar A. 2000: List of species of tropical American *Clinopodium* (*Labiatae*), with new combinations. – Kew Bull. **55**: 917–927.
- IUCN 2012: IUCN Red List categories and criteria: version 3.1, ed. 2. – Gland & Cambridge: IUCN. – Published at <http://www.iucnredlist.org/technical-documents/red-list-documents>
- Melnikov D. 2014: New combinations in the genus *Clinopodium* (*Lamiaceae*). – Novosti Sist. Vysš. Rast. **45**: 148–149.
- Méndez Santos I. E., Bécquer Granados E. R. & Herrera Oliver P. P. 2005: A new species of *Satureja* (*Lamiaceae*) from central Cuba. – Willdenowia **35**: 167–171.
- Pool A. 2008: New combination in *Clinopodium* (*Lamiaceae*) from Mesoamerica and Cuba. – Novon **18**: 508–510.
- Sauvalle F. A. 1873: Flora cubana. Enumeratio nova plantarum cubensium vel revisio catalogi Grisebachiani, exhibens descriptiones generum specierumque novarum Caroli Wright (Cantabrigiae) et Francisci Sau-

- valle, synonymiis nominibusque vulgaribus cubensis adjectis. – Havanae: Imp. “La Antilla,” de Cacho-Negrete.
- Urban I. 1919: Sertum antillanum IX. – Repert. Spec. Nov. Regni Veg. **16**: 132–151.
- Urban I. 1924a: Plantae cubenses novae vel rariores a clo. Er. L. Ekman lectae. II. – Pp. 177–272 in: Urban I. (ed.), Symbolae antillanae **9**. – Lipsiae: Fratres Born-
traeger; Parisiis: Paul Klincksieck; Londini: Williams & Norgate.
- Urban I. 1924b: Sertum antillanum. XXI. – Repert. Spec. Nov. Regni Veg. **20**: 337–352.
- Wagstaff S. J., Olmstead R. G. & Cantino P. D. 1995: Parsimony analysis of cpDNA restriction site variation in subfamily *Nepetoideae* (*Labiatae*). – Amer. J. Bot. **82**: 886–892.

Willdenowia

Open-access online edition www.bioone.org/loi/will  BioOne

Online ISSN 1868-6397 · Print ISSN 0511-9618 · Impact factor 0.680

Published by the Botanic Garden and Botanical Museum Berlin, Freie Universität Berlin

© 2017 The Authors · This open-access article is distributed under the CC BY 4.0 licence