

A new species of
Ouratea Aubl.
(*Ochnaceæ*)
from South America

Gisele SILVESTRE SALVADOR,
Élide PEREIRA DOS SANTOS
& Armando Carlos CERVI

FONTQUERIA **55**(39): 293-296 [seorsim 1-4]
MADRID, 22-IV-2006

FONTQUERIA is a series of botanical publications without administrative affiliation. It publishes original works in Botany, particularly those that are of interest to the editors. Its publications are in any language, the only limitation being the ability of the editorial team.

Accredited with the *International Association for Plant Taxonomy* for the purpose of registration or new non-fungal plant names.

PRODUCTION

Database consultant: Guillermo GONZÁLEZ GARCÍA
Typesetting: Ambrosio VALTAJEROS POBAR, Ulpiano SOUTO MANDELOS
Screen operators: Samuel FARENA SUBENULLS, Emilio NESTARES SANTAINÉS
Preprinting: Sonja MALDÍ RESTREPO, Demetrio ONCALA VILLARRASO

DISTRIBUTION

Postal distribution: contact the editor
Mail for electronic distribution: Fontqueria@yahoo.com

EDITOR

Francisco Javier FERNÁNDEZ CASAS. Madrid (MA)

JOINT EDITORS

Ramón MORALES VALVERDE. Madrid. German texts
Antonio Manuel REGUEIRO y GONZÁLEZ-BARROS. Madrid. English texts
Francisco Javier SÁNCHEZ GARCÍA. Cáceres. Latin texts
Ulpiano Emilio SOUTO MANDELOS. Pontevedra. Chartographic work

EDITING CONSULTANTS for this fascicle

Matías MAYOR LÓPEZ (FCO)
María Antonia RIVAS PONCE (UAM, Madrid)

ISSN: 0212-0623

Depósito legal: M-29282-1982

A new species of *Ouratea* Aubl. (*Ochnaceæ*) from South America

Gisele SILVESTRE SALVADOR,

Universidade Federal do Paraná. Departamento de Botânica. Programa de Pós-Graduação em Botânica. Caixa Postal 19031, CEP 81531-980, Curitiba, Paraná, Brasil
(gisele.salvador@ibest.com.br)

Élide PEREIRA DOS SANTOS

Universidade Federal do Paraná e Bolsista Pesquisadora do CNPq (elide@ufpr.br)

& Armando Carlos CERVI

Universidade Federal do Paraná e Bolsista Pesquisador do CNPq (accervi@ufpr.br)

SALVADOR, G. SILVESTRE, E. PEREIRA DOS SANTOS & A. C. CERVI (22-iv-2006). A new species of *Ouratea* Aubl. (*Ochnaceæ*) from South America. *Fontqueria* **55**(39): 293-296 [seorsim 1-4].

Keywords. Systematics, new species; *Ouratea* (*Ochnaceæ*); Bolivia (Santa Cruz de la Sierra), Brazil (Mato Grosso do Sul, Paraná, Tocantins), South America.

Abstract. A new species of *Ouratea* (*Ochnaceæ*) from Bolivia and Brazil is described: *O. Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi.

Zusammenfassung. Es wird hier eine neue Art der Gattung *Ouratea* (*Ochnaceæ*) aus Bolivia und Brasil: *O. Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi beschrieben.

Resumé. Une nouvelle espèce est décrite de la Bolivie et du Brésil dans le genre *Ouratea* (*Ochnaceæ*): *O. Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi.

Resumo. Descreve-se uma nova espécie boliviana e brasileira no gênero *Ouratea* (*Ochnaceæ*): *O. Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi.

Resumen. Se describe una especie nueva en el género *Ouratea* (*Ochnaceæ*), nativa de Bolivia y de Brasil: *O. Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi.

Ouratea Aubl. is a neotropical genus comprising about 120 species, cf. A. KANIS (1968), C. SASTRE (1988). The new species described here is clearly distinct from related species due to its smooth anthers and elongated gynophore.

***Ouratea Claudei* G. S. Salvador, E. P. dos Santos & A. C. Cervi, spec. nov.,**
Fontqueria **55**(39): 293-296 (22-iv-2006)

TYPUS: Brasil. Paraná, «(munic. Vila Alta) Parque Nacional de Ilha Grande», C. Kozera 1847, 25-I-2003 (MBM, holo-; MBM, iso-)

ILLUSTR.: lamina nostra i (figuræ a-j), pagella 295.

DISTR. GEOGR.: **America australis:** Bolivia (Santa Cruz de la Sierra); Brasil (Mato Grosso do Sul, Paraná, Tocantins). Tabula nostra 1, pagella 296.

ABBREVIATA DIAGNOSIS (vel character essentialis)

Affinis *Ourateæ salicifoliae* (A. Saint-Hilaire & Tulasne) Engler sed antheris lævibus, gynophoro elongato atque nervis non reticulatis differt. Affinis *Ourateæ castaneifoliae*

Engler sed antheris lævibus foliisque chartaceis differt.

MAIN DIAGNOSTIC FEATURES (or diagnostic characters)

Ouratea Claudei is similar to *O. castaneifolia* Engler and *O. salicifolia* (A. Saint-Hilaire & Tulasne) Engler but differs from *O. castaneifolia* by its smooth anthers, gynophore twice as long as the ovary, and leaf blades papery; it differs from *O. salicifolia* by its smooth anthers, gynophore twice as long as the ovary, globose carpophore, and lateral veins ascending with subparallel intersecondaries.

DESCRIPTION

Tree or shrub 2-5 m tall; branches glabrous. LEAF blade papery, ovate or elliptic, 9-13 × 4-5.5 cm, acute or acuminate; base obtuse; margin serrate; midrib prominent on the abaxial surface; lateral veins ascending with subparallel intersecondaries. INFLORESCENCE terminal, paniculate, 8-12 cm long. Bracts triangular, 1-2.5 mm long, deciduous. Pedicels 6-13 mm long; flower bud 6-9 × 2-4 mm, conical; sepals five, ovate or elliptic, 7-9.5 × 2-3.5 mm; petals five, spatulate, 7-12 × 4.5-5.5 mm; stamens 10, anthers 5-7 mm long, smooth; filament ca. 0.2 mm long; carpels five, ovary 0.7-1 mm long, gynophore 1.5-2.5 mm long, twice as long as the ovary, style 5-8 mm long. FRUIT with carpophore globose, 4-7 mm ø; carpels ellipsoid, 5-7.5 × 3.5-6 mm; seed ellipsoid, 5-7 × 3-4.5 mm (fig. 1, page 295).

ETYMOLOGY. The specific epithet was chosen in honour of Claude Sastre, French botanist, for his important studies on the family *Ochnaceae*.

DISTRIBUTION AND ECOLOGY. *Ouratea Claudei* is found in the gallery forests of Bolivia (department of Santa Cruz de la Sierra) and Brazil (states of Mato Grosso do Sul, Paraná and Tocantins); cf. map 1, page 296.

BOLIVIA, (08) SANTA CRUZ DE LA SIERRA:

- 20LNJ48** -14.60°, -062.58°; «(prov. Ñuflo Chávez) Perseverancia: vecindad del río Negro, tributario del río Baures, a 75 km sur del límite del depto. de Beni y 150 km W del río Paragua, 14°38'S, 062°37'W», *B. Mostacedo* 71, 16-V-1991, fl. (F).

BRASIL, (11) MATO GROSSO DO SUL:

- 21KXT23** -20.48°, -055.80°; Aquidauana (ppl), 20°28'16"S, 055°47'14"W, «(munic. Aquidauana) fazenda Fazendinha: área de brejo do rio Negro. Pantanal», *S. M. Salis* 773 & al., 08-IX-1993, fl. (CPAP).
- 21KXT23** Aquidauana (ppl), 20°28'16"S, 055°47'14"W, «(Largo do Mamão) fazenda Fazendinha: sub-região de Aquidauana, região do rio Negro. Pantanal», *A. Pott* 5784 & al., 05-II-1991, fl. (CPAP).

BRASIL, (18) PARANÁ:

- 22KBU29** -23.55°, -053.69°; Parque Nacional de Ilha Grande (rpds), 23°30'30"S, 053°43'40"W, «(munic. Vila Alta) Parque Nacional de Ilha Grande», *C. Kozera* 1690 07-XI-2002 (MBM).
- 22KBU29** Parque Nacional de Ilha Grande (rpds), 23°30'30"S, 053°43'40"W, «(munic. Vila Alta) Parque Nacional de Ilha Grande», *C. Kozera* 1826, 24-I-2003, fr. (MBM).
- 22KBU29** Parque Nacional de Ilha Grande (rpds), 23°30'30"S, 053°43'40"W, «(munic. Vila Alta) Parque Nacional de Ilha Grande», *C. Kozera* 1847, 25-I-2003, fl., fr. (MBM, holo-; MBM, iso-; typus *O. Claudei*).
- 22KBU29** «Pôrto da Balsa: 4 km para o S, no lado W do rio Paraná», *R. S. Ziller* 1185,





FIGURE 1. *Ouratea Claudei*. a) Branch with inflorescence. b) Detail of the leaf venation (adaxial side). c) Flower bud. d) Flower. e) Sepals. f) Petal. g) Anther. h) Ovary, gynophore, base of the style and filaments. i) Fruit. j) Seed, lateral view. a-c, e, f, h) *C. Kozera 1847* (MBM). d) *S. F. Lolis & E. dos Santos, s. n.* (MBM). g) *C. Kozera 1690* (MBM). i, j) *R. S. Ziller 1185* (MBM).



14-XI-1995, fr. (EFC; MBM).

22KBU29 «entre Pôrto Figueira e Pôrto Cerâmica: no rio Paraná», *J. Carneiro* 127, 06-XII-1995, fl. (MBM).

BRASIL, (31) TOCANTINS:

22LFP50 -10.81°, -049.58°; Lagoa da Confusão (ppl), 10°47'37"S, 049°37'25"W, «(munic. Lagoa da Confusão) Ilha do Formoso: margem esquerda do rio Formoso», *S. F. Lolis & E. dos Santos*, s/n, 21-VIII-1996, fl. (UEC; RB; UPCB).

CITED LITERATURE

- KANIS, A. (1968). A revision of the *Ochnaceae* of the Indo-Pacific Area. *Blumea* **16**: 1-82.
 SASTRE, C. (1988). Studies on the Flora of the Guiana 34. Synopsis generic *Ouratea* Aubl. (*Ochnaceae*). *Bull. Mus. Natl. Hist. Nat. Paris, sér. 4, sect. B, Adansonia* **1**: 47-67.