

Identification, cloning and lactonase activity of recombinant protein of *N*-acyl homoserine lactonase (AiiA) from *Bacillus thuringiensis* 147-115-16 strain

Identificación, clonación y actividad lactonasa de la proteína recombinante de *N*-ácil homoserina lactonasa (AiiA) de *Bacillus thuringiensis* cepa 147-115-16

Álvaro M. Florez^{*†}, Adriana González^{*}, Carmen J. Pedroza^{**}, Elizabeth Correa^{**},
Nohora J. Rueda^{*}, Sergio Orduz^{**}

Abstract

The quorum-quenching *N*-acyl homoserine lactonases are a family of bacterial metalloenzymes that participate in degradation of *N*-acyl homoserine lactones (AHLs), disrupting the quorum sensing system of gram negative bacterial species. From a collection of *Bacillus thuringiensis* strains isolated in Colombia from plants and exhibiting toxic activity against lepidopteran insects, 310 bacterial isolates were tested to determine lactonase activity by using biosensor systems in presence of synthetic *N*-hexanoyl-L-homoserine lactone (C6-HSL) and *N*-octanoyl-L-homoserine lactone (C8-HSL). From them, 251 strains showed degrading activity to both C6-HSL and C8-HSL, 57% exhibited degrading activity to C6-HSL and 43% to C8-HSL. One *B. thuringiensis* strain, denoted as 147-115-16, that exhibit high degrading activity to C6-HSL and C8-HSL, was able to attenuate soft rot symptoms in infected potato slices with *Pectobacterium carotovorum*. This strain contains an homologous of the *aiiA* gene that was cloned, sequenced and expressed in *Escherichia coli* DE3. The recombinant protein AiiA₁₄₇₋₁₁₅₁₆ display activity to C6-HSL, C8-HSL, *N*-(β -ketocaproyl) (3-O-C6-HSL) and *N*-3-oxo-dodecanoyl (3-O-C12-HSL). The recombinant strain in the presence of *P. carotovorum* cultures was able to attenuate the infection, suggesting that it interferes either on the accumulation or response to the AHLs signals. According to this data and based on previous report from recombinant AiiA₁₄₇₋₁₁₅₁₆, this enzyme exhibit activity to wide range of catalytic substrates suggesting its industrial application in the disease control programs by plants transformation.

Key words: lactones, Quorum sensing, Quorum quenching, Lactonases, *Pectobacterium carotovorum*

Resumen

Las *N*-ácil homoserina lactonasas son una familia de metaloenzimas bacterianas que participan en la degradación de *N*-ácil homoserina lactonas (AHLs) interrumpiendo el sistema de detección de quórum sensing de bacterias Gram negativas. A partir de una colección de cepas de *Bacillus thuringiensis* aisladas del filoplano de plantas colombianas que presentaron actividad tóxica contra insectos lepidópteros, 310 fueron probadas para determinar actividad lactonasa mediante el uso de sistemas de biosensores en presencia de *N*-hexanoilo-L-homoserina lactona (C6-HSL) y la *N*-octanoilo-L-homoserina lactona (C8-HSL) sintéticas. De estas cepas, el 251 mostraron actividad para ambas lactonas y de estas, el 57% mostró actividad a C6-HSL y el 43% a C8-HSL. Una cepa de *B. thuringiensis* denominada 147-115-16 que mostró alta actividad para C6-HSL y C8-HSL, fue capaz de atenuar los síntomas de la pudrición blanda en rodajas de papa infectadas con *Pectobacterium carotovorum*. Esta cepa contiene un gen homólogo a *aiiA*, el cual este fue clonado, secuenciado y expresado en *Escherichia coli* DE3. La proteína recombinante AiiA₁₄₇₋₁₁₅₁₆ exhibe actividad para C6-HSL y C8-HSL, así como para *N*-(β -cetocaproil) (3-O-C6-HSL) y *N*-3-oxo-dodecanoil (3-O-C12-HSL). La cepa recombinante en presencia de *P. carotovorum* fue capaz de atenuar la infección, sugiriendo que interfiere con la acumulación o respuesta de las señales AHLs. Según estos datos y

* Laboratorio de Biología Molecular y Biotecnología, Instituto MASIRA, Facultad de Salud, Universidad de Santander, UDES, Bucaramanga, Santander, Colombia.

** Escuela de Biociencias, Facultad de Ciencias, Universidad Nacional de Colombia sede Medellín, Medellín, Colombia.
Correspondence author amflores@udes.edu.co. Phone +57 6516500 Ext. 1375. Fax: +57 6516492

basados en el reporte previo sobre la actividad hidrolítica de la proteína recombinante AiiA₁₄₇₋₁₁₅₁₆, esta enzima posee un actividad contra un amplio número de sustratos lo cual sugiere su aplicación en la industria en el control de enfermedades, mediante la transformación de plantas.

Palabras clave: Lactones, Quorum sensing, Quorum quenching, Lactonases, *Pectobacterium carotovorum*.

Recibido: octubre 25 de 2013

Aprobado: mayo de 2014

Introduction

In the majority of gram negative bacteria quorum sensing is mainly mediated by signaling cascades, involving *N*-acyl-homoserine lactones (AHLs) (Von Bodman *et al.*, 2003). AHLs contain a common homoserine lactone ring, differing in their carbon length (4 to 18 carbons), degree of saturation, and the presence of C3 acyl chain modifications. AHLs are multifunctional molecules that when they accumulate regulate relevant functions during bacterial growth such as production of extracellular enzymes, bioluminescence, biofilm formation, and coordinate cellular activities associated with host pathogen interactions (Atkinson and Williams, 2009; Williams and Camara, 2009). Many bacterial species, such as *Pectobacterium carotovorum*, are highly pathogenic to plants by triggering the expression of virulence factors via quorum sensing (QS) (Williams, 2007). *P. carotovorum* is an etiologic agent of soft rot disease in a wide variety of cultivated plants (Charkowski, 2009). This bacterium regulates the production of different exoenzymes by secreting AHLs, including *N*-(3-oxohexanoyl)-L-homoserine lactone (3-O-C6-HSL) and *N*-(3-oxooctanoyl)-L-homoserine lactone (3-O-C8-HSL) (Jones *et al.*, 1993; Pirhonen *et al.*, 1993; Cha *et al.*, 1998; Smadja *et al.*, 2004; Chatterjee *et al.*, 2005; Liu *et al.*, 2008). The role of QS in *P. carotovorum* pathogenicity have been reviewed (Barnard *et al.*, 2007).

In contrast, there are microorganisms acting as antagonists of QS by producing signal interference mechanisms known as quorum quenching (QQ), (Kalia, 2013) mediated by AHL-degrading enzymes. There are two families of QQ enzymes, lactonases and acylases. The lactonase family cleaves the lactone ring of AHLs releasing *N*-acyl homoserines, which are not recognized as a QS signal, some examples include AiiA from *Bacillus* sp., AttM from *Agrobacterium tumefaciens* strain A60, AiiB from *A. tumefaciens* strain C58 (Carlier *et al.*, 2003; Dong *et al.*, 2000) and AhID from *Arthrobacter* sp. (Park *et al.*, 2003). Acylases are known to be capable of cleaving the acyl bonds and releasing a homoserine lactone ring (Lin *et al.*, 2003; Sio *et al.*, 2006), some representative enzymes are AiiD from *Ralstonia* XJ12B, and PvdQ from *Pseudomonas aeruginosa* PAO1. After the cleavage of the AHLs molecules, the released products become more soluble in water, making them an available nutrient source to the surrounding microorganisms, like *Variovorax paradoxus*

(VAI-C), which is capable of using AHLs as a sole nutrient source (Leadbetter and Greenberg, 2000).

Production of QQ *N*-acyl homoserine-lactonases was first reported in a strain of the genus *Bacillus* sp. (Dong *et al.*, 2000; Dong *et al.*, 2004). It was also reported in *Bacillus thuringiensis*, *B. cereus* and *B. mycoides* (Dong *et al.*, 2002; Lee *et al.*, 2002). The first AHL-lactonase that was completely characterized was AiiA_{240B1} and it was found that belongs to the metallo- β -lactamase family based on the presence of a HXHXDH motif, and an identical Zn²⁺ binding motif (Dong *et al.*, 2000). This enzyme has the ability to inactivate AHLs with different acyl chain lengths and substitutions at the C3 position, including *N*-butyryl-L-homoserine lactone (C4-HSL), 3-O-C6-HSL, 3-O-C8-HSL, and 3-O-C12-HSL (Dong *et al.*, 2001).

The second AHL-lactonase (AiiA_{BTK}) reported was isolated from *B. thuringiensis* subsp. *kurstaki*, and work on a wider range of AHLs substrates, with maximum activity towards C10-HSL. This enzyme contains two Zn²⁺ ions into the active site and possesses the same conserved residues present in all AHL-lactonases whose activity depends in metal coordination (Kim *et al.*, 2005). For both characterized enzymes, AiiA_{240B1} and AiiA_{BTK}, important differences in the affinity for metal ions under chelating agents have been described (Wang *et al.*, 2004; Kim *et al.*, 2005; Thomas *et al.*, 2005).

Based on that there is widespread existence of AHL-lactonases in *B. thuringiensis* (Dong *et al.*, 2002; Lee *et al.*, 2002), and in Colombia more than 600 *B. thuringiensis* isolates have been characterized looking for insecticidal activity (Uribe *et al.*, 2003; Jara *et al.*, 2006; Armengol *et al.*, 2007), we aimed at isolating AHL degrading from a *B. thuringiensis* collection obtained from Colombian phyloplane. In this study we screened a *B. thuringiensis* collection and described the cloning, sequencing and the activity of recombinant protein of a *N*-acyl homoserine lactonase from *B. thuringiensis* ₁₄₇₋₁₁₅₁₆.

Materials and methods

Bacterial strains and growth conditions

Two different AHL biosensors strains were used to detect AHL-lactonase activity, *Chromobacterium violaceum* CVO26 (mini Tn5 mutant of 31532) (McClellan *et al.*, 1997; Winson *et al.*, 1998) and *A. tumefaciens* NTL4 *tral-lacZ* fusion (pCF218) (pCF372) (Fuqua *et al.*,

1994). *C. violaceum* CVO26 was grown in LB medium (1% peptone, 0.5% yeast extract, 0.5% NaCl) solidified with 1.2% agar, and supplemented with kanamycin (20 µg/ml) (Invitrogen®). *A. tumefaciens* NTL4 was grown in AT minimal salts medium as was described in Tempé *et al.* (1977), at 30°C, supplemented with tetracycline 4.5 µg/ml (Sigma-Aldrich®), spectinomycin 50 µg/ml (Sigma-Aldrich®) and 40 µg/ml of X-gal (Promega®). *Escherichia coli* ArticExpress (DE3) cells were used for cloning and expression and were cultured at 37°C in LB medium supplemented with 100 µg/ml of ampicillin (Sigma-Aldrich®). *B. thuringiensis* strains were cultured in LB medium at 30°C in agar plates and broth.

Plant material with symptoms of soft rot disease were selected in order to isolate bacterial plant pathogens as described (Pérombelon and Kelman, 1980). Samples were taken from affected zones from lettuce (*Lactuca sativa* var. Batavia), calla lily (*Zantedeschia aethiopica* var. Spreng), potato (*Solanum tuberosum* var. Diacol capiro) and lulo (*S. quitoense* var. Castilla), and processed according to phytopathological methods (Agrios, 2005). Different bacterial colonies were obtained in nutrient agar from serial dilutions of macerated tissues. The isolated colonies were grown in D-3 selective medium for *P. caratovorum* (Kado *et al.*, 1970). Complementary tests for species identification were carried out as described (Schaad *et al.*, 2001).

***B. thuringiensis* AHL-lactonase activity**

AHL-lactonase activity was evaluated in 310 *B. thuringiensis* isolates as described by Lee *et al.* (2002), with some modifications. A culture of each *B. thuringiensis* strain was grown overnight until reaching an optical density of 1.1 at 600 nm. Then, 20 µM of synthetic C6-HSL and C8-HSL (Sigma-Aldrich®) were added independently into each culture and the reaction mixture was incubated at 30°C with shaking during 30, 60 and 120 min. Bacterial cells were centrifuged at 3000 x g for 10 min and the supernatants corresponding to each incubation time and to each *B. thuringiensis* strain were heated for 3 min at 95°C and placed into petri dishes with 10 wells with LB medium or AT agar containing an agar-suspension culture of the biosensor strain. These plates were incubated for 24 h at 30°C. The AHL-lactonase activity was evaluated at different times by measuring the diameters of the colored areas surrounding the wells in CVO26 (purple) and NTL4 (blue). Based on the sizes of the colored areas, a scale was designed; diameters between 0.0 and 0.39 cm² were assigned with (+++) indicating *B. thuringiensis* strains displaying high AHL-lactonase activity; between 0.4 and 0.74 cm² were assigned with (++) indicating a moderate activity; and between 0.75 and 0.89 cm² were assigned with (+) indicating a low activity. Finally, diameters larger than 0.9 cm² were considered without AHL-lactonase activity. Only the *B. thuringiensis* strains that exhibit high AHL lactonase activity (+++) to each

substrate to either C6-HSL or C8-HSL at 30, 60 and 120 min or 60 and 120 min were reported.

Screening of AHL production by *P. caratovorum*

The plant pathogenic bacterium *P. caratovorum* was streaked in parallel to the biosensor strains in LB agar for CVO26 and in the same way for NTL4 but in AT medium supplemented with 50 µg X-gal (Ravn *et al.*, 2001; Chu *et al.*, 2011), and plates were incubated for 24 h at 30°C. To evaluate the presence of AHLs, CVO26 was streaked on LB agar supplemented with C6-HSL, because its sensor system is inhibited by long acyl chain molecules. For this last assay, all plates with CVO26 were incubated for 24 h at 30°C for production of purple pigment, and then plates were re-incubated at 30°C for 24 h in order to evaluate inhibition of pigment formation.

In vitro control of soft rot symptoms

To evaluate the effect of AHL-lactonase activity of *B. thuringiensis* on *P. caratovorum* infection, potatoes (*S. tuberosum* var Diacol Capiro) were washed with tap water and treated as described Dong *et al.* (2004). Five mm thick potato slices were dipped in a *B. thuringiensis* 12 hr culture containing 2x10⁷ CFU/ml for 20 s and then were air dried in a laminar flow cabinet for 20 min. Then, three streaks were made on the surface of the slices and inoculated with 2.5 µl of cell culture media containing 1x10⁵ UFC/ml of an *P. caratovorum* suspension isolated from potato. Each slice was placed in a petri dish in a humid chamber at 30°C. Disease symptoms were evaluated after 24 and 48 h of treatment. Each treatment was performed in quadruplicate. Based on the severity of the soft rot symptoms produced in the potato slices, a scale was developed to qualify the effect of *B. thuringiensis* on the disease caused by *P. caratovorum*. Absence of soft rot was assigned with (-), presence of small soft rot area and little wet was considered with (+), presence of moderate soft rot area, dark and wet was considered with (++) and, presence of high soft rot area, very dark and very wet was considered with (+++).

Cloning and sequencing of the *aiiA* gene

The *B. thuringiensis* strain displaying greater AHL-lactonase activity with the biosensor strains *A. tumefaciens* NTL4 and *C. violaceum* CVO26 and the best control of *P. caratovorum* pathogenicity was selected and its DNA was extracted (Ceron *et al.*, 1995). Using PCR the *aiiA* gene was amplified with the following primers: 5' ATCGGATCCATGACAGTAAAGAAGCTTTATTTCG3' and 5' GTCGAATTCTCAACAAGATACTCCTAATGATGT3' designed according to the *aiiA* sequences previously reported (Dong *et al.*, 2002). Amplification conditions were: denaturation at 94°C for 3 min followed by 28

cycles at 94°C for 30 s, 50°C for 30 s, and 72°C for 1 min in a MJ Research Thermocycler (Bio-Rad®). The amplified fragments were purified with a QIAquick PCR purification system (Qiagen®) and cloned into pGEM-T vector (Promega®). Clones were selected in LB media supplemented with IPTG (40 µg/ml) (Invitrogen®), X-gal (40 µg/ml) (Invitrogen®) and ampicillin (100 µg/ml) (Sigma-Aldrich®). The gene was sequenced in both strands and the results were analyzed using Blastn, Blastp and ClustalW algorithm in Bioedit (Hall, 1999). The similarity and identity analyses were performed using MatGat (Campanella *et al.*, 2003), and evolutionary relationships between AHL-lactonases were done using MEGA5 (Tamura *et al.*, 2011).

The subcloning into the expression vector pCold IV (Takara-BIO-INC®), gene expression and protein purification are described by Pedroza *et al.* (2014).

Homology and phylogenetic dendrogram of AHL-degrading enzyme

The similarity and identity analyses were performed using MatGat (Campanella *et al.*, 2003), and evolutionary relationships between AHL-lactonases were done using MEGA5 (Tamura *et al.*, 2011). The evolutionary history was inferred using the Neighbor-Joining Method (Saitou and Nei, 1987). The optimal tree with the sum of branch length was = 0.34721705. The tree was drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Maximum Composite Likelihood Method (Tamura *et al.*, 2004) and are in the units of the number of base substitutions per site. The analysis involved 22 nucleotide sequences. Codon positions included were 1st+2nd+3rd+Noncoding. All positions containing gaps and missing data were eliminated. There were a total of 570 positions in the final dataset.

AHL-lactonase activity of recombinant *E. coli* DE3

The recombinant *E. coli* DE3 containing the expression cassette pColdIV-AiiA was grown at 37°C in LB broth with ampicillin (50 µg/ml) at 200 rpm. After 5 hours of growth (O.D. 600 nm = 0.83), the IPTG (3.2 mM) was added and the incubation was continued overnight to reach an O.D. of 2.6 at 600 nm (Zhang *et al.*, 2007). The cell culture was divided in several fractions in order to determine the AHL-lactonase activity. The first and second fractions were added each to LB medium in presence of 100 µM of synthetic C8-HSL and C6-HSL, the third and fourth fractions were added to LB medium in the presence of 100 µM of synthetic 3-O-C6-HSL and 3-O-C12-HSL, respectively. The fifth fraction was co-inoculated with 20 µl of a *P. caratovorum* culture (OD₆₀₀ = 0.8) in 5 ml of LB media. All the fractions were incubated at 37°C for 18 h at 200 rpm. The experiments included *B. thuringiensis*₁₄₇₋₁₁₅₁₆, as positive control and *E. coli*

DE3 strain and *P. caratovorum* alone as negative control. Once the incubation time was finished each culture was centrifuged at 5800 x g during 10 min at 21°C and the supernatants were collected. Six lines were made with a sterile blade on each petri dish containing either LB media for CV026 strain or AB media with X-gal for NTL4 strain. Each line was 1 cm wide and 3 mm apart. Five µl of each supernatant was served in one end of each groove, and the samples were allowed to dry, and then each respective reporter strain was streaked. The petri dishes were incubated at 30°C for 24 hours. To determine the pigment produced by the reporter strains, the length was measured and the reproducibility was determined by 20 replicates.

Statistical Analysis

Data was analyzed by using the ANOVA test. The confident values were set at 95%.

Results

***B. thuringiensis* AHL-inactivating activity**

Inactivation of AHLs by *B. thuringiensis* strains is considered effective when in a given evaluation period, the size of the colored area decreases with the time. When no color is observed at the first time of evaluation (30 min), the strain is listed as a candidate for *in vitro* pathogenicity inhibition assays. From the 310 *B. thuringiensis* strains isolated from phylloplane plants tested, 251 showed AHL-lactonase activity with C6-HSL and C8-HSL; C6-HSL was degraded by 57% of the *B. thuringiensis* strains, while C8-HSL was inactivated by 43% of the *B. thuringiensis* strains. *B. thuringiensis* strains that exhibit strong AHL lactonase activity against C6-HSL and C8-HSL are shown in table 1. The screening to these strains using CV026 in are shown in figure 1. The detectable concentration of the reporter strains was between 100-120 µM of AHLs.

Effect of *B. thuringiensis* cultures on soft rot symptoms caused by *P. caratovorum*

To test the attenuation of soft rot symptoms, *B. thuringiensis* strains displaying the highest AHL-lactonase activity according to the results with the biosensor strains were selected. The plant pathogen *P. caratovorum* isolated from potato was used in these assays and its pathogenicity was evaluated on potato slices showing severe tissue maceration. When potato slices were pretreated with *B. thuringiensis* suspension (2x10⁷ UFC/ml) before inoculation with *P. caratovorum*, the soft rot symptoms were considerably less severe compared with the soft rot produced on slices without previous treatment. According to the evaluation scale of soft rot symptoms, the strain *B. thuringiensis*₁₄₇₋₁₁₅₋₁₆ displayed the highest

efficiency inhibiting *P. carotovorum* pathogenicity (figure 2).

Cloning and sequence analysis of the *AiiA* gene

*B. thuringiensis*₁₄₇₋₁₁₅₁₆ strain displayed the best results in the biosensor assays, as well as in the *in vitro* pathogenicity assays; therefore, it was selected for molecular characterization. The length of expected amplified PCR product containing the gene coding *AiiA* was around 820 bp. Several clones that showed this length, exhibit a 754 bp ORF with sequence homology to *aiiA* genes reported in GenBank. The DNA sequence of *B. thuringiensis*₁₄₇₋₁₁₅₁₆ gene was reported in GenBank with accession number EF379241. Comparison of the *B. thuringiensis*₁₄₇₋₁₁₅₁₆ *AiiA* gene with other *AiiA* genes revealed maximum identity in the nucleic acid sequence between 93% to 99%. The dendrogram show that *aiiA* gene of *B. thuringiensis*₁₄₇₋₁₁₅₁₆ is most similar to *B. thuringiensis* subsp. *aizawai* and *indiana* (figure 3). At protein level three conserved domains in the protein *aiiA* of the *B. thuringiensis*₁₄₇₋₁₁₅₁₆ were detected by BLAST CD search of Conserved Domain Database

(rpsblast; <http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>; domain COG0491GloB, Zn dependent hydrolases, including glyoxylases (residues 78-250); domain pfam00753, the lactamase-β domain, metallo-beta-lactamase superfamily (residues 33-235); and domain COG1237, metal dependent hydrolases domain of the β lactamase superfamily (residues 22-161). Using MEME and MAST programs (Bailey and Gribskov, 1998), in which protein sequences are analyzed for similarities among hydrolases, three motifs were also found in *B. thuringiensis*₁₄₇₋₁₁₅₁₆ *AiiA*: motif 1 (residues 11-60); motif 2 (residues 76-125), and motif 3 (residues 128-177). These motifs of the *AiiA* are also present in the three Zn-dependent hydrolases.

Recombinant *E. coli* DE3- pCold-*AiiA* shown lactonase activity

The lactonase activity of the recombinant *E. coli* DE3 was tested with synthetic lactones, C8-HSL and C6-HSL in the presence of the reporter strains CV026 and, with 3-O-C6-HSL and 3-O-C12-HSL in the case of NTL4. The results showed lack of purple pigmentation of strain




Figure 1. *C. violaceum* CVO26 with C6-HSL (HHL) as substrate. Well (+), reaction buffer with C6-HSL (20 μM); well (-), reaction buffer only.

Table 1. AHL-lactonase activity of eight *B. thuringiensis* strains determined by biosensor strains. **A:** *C. violaceum* CVO26 with C6-HSL as substrate. Well (+), reaction buffer with C6-HSL (20 μM); well (-), reaction buffer only.

Well number	Strain	Phylloplane origin	Evaluation time		
			30 min	60 min	120 min
1	147-11516	<i>Phaseolus vulgaris</i>	+++	+++	+++
2	146-17202	<i>Piper glanduligeron</i>	-	-	-
3	146-15505	<i>Piper calceolarium</i>	-	-	-
4	147-0206	<i>Zea mays</i>	+	+++	+++
5	146-16312	<i>Piper crassinervium</i>	+	+++	+++
6	146-16702	<i>Piper danielgonzalezi</i>	-	+++	+++
7	146-15533	<i>Piper calceolarium</i>	+++	+++	+++
8	147-12003	<i>Zea mays</i>	+++	+++	+++




Figure 2. Attenuation of *P. carotovorum* pathogenicity by AHL-lactonase activity of *B. thuringiensis*. (E) Potato slice inoculated with *P. carotovorum* alone, (E+Bt) potato slice pre-treated with *B. thuringiensis* and inoculated with *P. carotovorum*, (Bt) potato slice pre-treated with *B. thuringiensis* suspension only. **A.** No control of *P. carotovorum* symptoms by strain *B. thuringiensis*₁₄₆₋₁₇₂₀₂, **B.** Moderate control of *P. carotovorum* symptoms by strain *B. thuringiensis*₁₄₆₋₁₆₇₀₂, **C.** Total control of *P. carotovorum* symptoms by strain *B. thuringiensis*₁₄₇₋₁₁₅₁₆.

CV026 (figure 4) and in the case of NTL4, exhibit lack of blue pigmentation. In the presence of the pathogen *P. carotovorum*, the QQ signals were disrupted (figure 4A, lane 2 and 4B lane 1).

In all the cases, disruption of QS by the recombinant *E. coli* DE3, is shown by the lack of pigmentation in the presence of biosensor strains in comparison with the control, with the pathogen alone, which exhibits pigment production in response to both, long and short chain lactones. ANOVA analysis showed statistically significant differences between lactones indicating the preference to short chain than those that exhibit long-chain oxo substitutions. There was no statistical difference within the same group of lactones (figure 5). These results demonstrated the preference of recombinant AiiA₁₄₇₋₁₁₅₁₆ for C8-HSL and C6-HSL.

Discussion

N-acyl homoserine lactones are signaling molecules that are inactivated by the enzyme AHL-lactonase by hydrolyzing the lactone ring, and in consequence, inhibiting the virulence of gram negative pathogens such as *P. carotovorum* (Dong *et al.*, 2004). In this study, the strain *B. thuringiensis*₁₄₇₋₁₁₅₁₆ isolated from the phylloplane of Colombian plants, was found to display attenuation of *P. carotovorum* symptoms in potato bioassays.

It has been established that the potential of some *B. thuringiensis* strains to attenuate infections depends on

breaking down AHL molecules that contain different acyl side chain lengths (Lee *et al.*, 2002). In this study, two synthetic AHLs with different acyl chain lengths, C6-HSL and C8-HSL, were inactivated by *B. thuringiensis*₁₄₇₋₁₁₅₁₆ detected by the biosensor strains *A. tumefaciens* NTL4 and *C. violaceum* CV026. The biosensor strain NTL4 showed blue color as response in the presence of AHLs produced by *P. carotovorum*. In the *in vitro* assays using pretreated potatoes slices with *B. thuringiensis*₁₄₇₋₁₁₅₁₆ cultures, attenuation of the soft rot symptoms caused by *P. carotovorum* were observed, which could suggest an interference of the *B. thuringiensis*₁₄₇₋₁₁₅₁₆ AiiA towards AHL signaling molecules produced by *P. carotovorum*. This signaling molecules including those that have been reported to *P. carotovorum* such as 3-oxo substitutions in the acyl chain (Chatterjee *et al.*, 2005), have been reported as a main regulators of extracellular enzymes involved in potato infection and responsible for symptoms development (Jones *et al.*, 1993; Pirhonen *et al.*, 1993). The *B. thuringiensis*₁₄₇₋₁₁₅₁₆ strain contains also the *aiaA* gene that encodes a AiiA protein with a molecular weight of approximately 28,2 kDa, which is similar to the predicted molecular mass of 28,036 Da of the protein reported by Dong *et al.* (2000). The amino acid sequence alignments showed identities of 90.8% to 99.6% to other AHL-lactonases described (Lee *et al.*, 2002). The evolutionary distances used to infer a phylogenetic tree confirm the wide distribution of *aiaA* homologous genes in *B. thuringiensis* subspecies and individuals and with identities above 93%. At protein level it was also observed the region




Figure 3. Phylogenetic dendrogram of AHL-degrading enzymes constructed using the neighbor-joining method. The bar indicates 1% estimated sequence divergence and the black circle highlight *B. thuringiensis*₁₄₇₋₁₁₅₁₆




Figure 4. A. Attenuation activity of AiiA using the reporter strain CVO26. Lane 1. Pathogen *P. carotovorum*; Lane 2: Recombinant strain (DE3) co-inoculated with *P. carotovorum*; Lanes 3 and 4: wild type strain *B. thuringiensis*₁₄₇₋₁₁₅₁₆ co-inoculated with *P. carotovorum*; Controls: Lane 5: C6-HSL in the presence of CV026; Lane 6: C8-HSL in the presence of CV026 reporter strain. **B.** Attenuation activity of AiiA using the reporter strain NTL4 in the presence of AHL. Lane 1. Recombinant strain DE3 co-inoculated with the pathogen *P. carotovorum*; Lane 2: Recombinant strain (DE3); Lane 3: Recombinant strain DE3 in presence of 3-O-C12-HSL; Lane 4: Recombinant strain DE3 in the presence of 3-O-C6-HSL. Lane 5: Pathogen *P. carotovorum* grown alone; Lane 6: negative control using the reporter strain alone without culture or extract added to the loading area.




Figure 5. Graphic obtained of the ANOVA (95%) analysis performed from the data obtained on the activity essays. The contrast between ($a-b=-0.666667$) and ($c-d=-123.939$) there are not shown significant difference. However, between ($a-c=-10.397$) ($a-d=-116.364$) ($b-c=-97.303$) and ($b-d=-109.697$) there are shown a significant difference. All data analysis shown \pm Limits=2,278940. The lowest pigmentation means highest activity. To C6-HSL and C8-HSL was obtained the greatest activity. To 3-O-C6-HSL and 3-O-C12-HSL was observed larger pigmentation in the biosensor NTL4, the activity among these AHLs was not significant.

$^{104}\text{HLHFDH}^{109} \approx \text{H}^{169}$, that correspond to the motif $^{104}\text{HXHDXDH}^{109} \approx \text{H}^{169}$ which is a completely conserved region among AiiA proteins. This motif contains a dinuclear Zn^{2+} center in which the Zn1 is coordinated by H^{104} , H^{106} and H^{169} while the Zn2 is coordinated by H^{109} , H^{235} , D^{108} and D^{191} (Kim *et al.*, 2005). According to this data, the AiiA from *B. thuringiensis* $_{147-11516}$ seems to be metallo hydrolytic enzyme. The differences observed between AiiA from *B. thuringiensis* $_{147-11516}$ (Pedroza *et al.*, 2014) and AHL-lactonase $_{\text{BTK-AiiA}}$ (Kim *et al.*, 2005) in 23 amino acids in which Ile-73 and Met-138 are part of hydrophobic channel and Tyr-165 and Ser-170 differ from the conserved region $^{165}\text{HTPGHTPGH}^{173}$ involved in Zn coordination, could explain the capability of AiiA $_{147-11516}$ to retain the activity in presence of high concentrations of EDTA and increase three and four times its activity to C4-HSL and C8-HSL respectively (Pedroza *et al.*, 2014).

The lactonase activity of the recombinant strain DE3 harboring the construct pCOLDIV:*aaiA* was evaluated in the presence of synthetic lactones with different acyl side chain lengths, C6-HSL, C8-HSL, 3-O-C6-HSL and 3-O-C12-HSL in the presence of the reporter strains *C. violaceum* CVO26 and *A. tumefaciens* NTL4. Based on the observation that the lactones are diffusible on solid media, and that the lactonase activity is inversely proportional to the color intensity, a light blue pigment of the biosensor was observed to 3-O-C6-HSL and

3-O-C12-HSL when they were tested with the biosensor strain NTL4. These data indicates that lactonase inactivation was not complete and that there was remaining lactones in the media (data not shown). It was also observed the ability of recombinant *E. coli* DE3 strain to disrupt *P. caratovorum* communication when they were co-inoculated in the presence of the reporter strain. This attenuation suggests that the lactonase produced by the recombinant strain exhibit activity to modified lactones and interferes to signaling molecules used by *P. caratovorum*.

The high homology and the presence of one conserved motif $^{104}\text{HXHDXDH}^{109} \approx \text{H}^{169}$ indicate that the AHL-lactonase $_{147-11516}$ belongs to the AHL-lactonase family. The ability of the recombinant strain in *P. caratovorum* suggest an important role of AHL-lactonase to interfere the AHLs signaling or AHLs response. Recently, we report the enzymatic hydrolysis from recombinant AiiA $_{147-11516}$ lactonase in order to determine its application (Pedroza *et al.*, 2014). The activity to broad catalytic spectrum of lactones including as *N*-butyryl-L-Homoserine lactone (C4-HSL), *N*-heptanoyl-L-Homoserine lactone (C7-HSL) and the enzymatic hydrolysis to C6-HSL in alkaline pH, activity in high temperature (60°C) and decreasing of enzymatic activity by two metal ions, suggest that this enzyme is a promissory candidate for the disease control programs by plants transformation. Other scenarios such as the addition of enrichment cultures of AHL degrading bacteria *Bacillus sp.* that increase the survival of aquaculture species (Tinh *et al.*, 2008), the control over QS based bacterial biofilms in biorefineries (Hong *et al.*, 2012) and the incorporation of QS inhibitors in medical and dental implants and catheters (Choudhary and Schmit-Dannert, 2010), provide further applications.

Aknowledgments

The authors thank Dr. Clay Fuqua (Indiana University, Bloomington, Indiana) for providing *C.violaceum* CVO26 and *A. tumefaciens* NTL4. We thank Dr. Oscar Alzate, for his critical review of the manuscript. This work was supported by grants from The Colombian Department of Science, Technology and Innovation (2213-12-13773 and 1299-12-17827) to AMF.

Bibliography

- Agrios, G. C., 2005. Plant Pathology, Fifth ed. Academic Press, (Ed.), New York. Pp 952.
- Armengol G., Escobar M.C., Maldonado M.E., Orduz S. 2007. Diversity of Colombian strains of *Bacillus thuringiensis* with insecticidal activity against dipteran and lepidopteran insects. *Journal of applied microbiology*. 102(1): 77-88.
- Atkinson S., Williams P. 2009. Quorum sensing and social networking in the microbial world. *Journal of The Royal Society Interface*. 6(40): 959-678.
- Bailey T.L., Gribskov M. 1998. Methods and statistics for combining motif match scores. *Journal of Computational Biology*. 5(2): 211-221.

- Barnard A.M., Bowden S.D., Burr T.S., Coulthurst J., Monson R.E., Salmond G.P. 2007. Quorum sensing, virulence and secondary metabolite production in plant soft-rotting bacteria. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 362(1483): 1165-1183.
- Campanella J.J., Bitincka L., Smalley J. 2003. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences. *BMC Bioinformatics*. 4(1): 29.
- Carlier A., Uroz S., Smadja B., Fray R., Latour X., Dessaux Y., Faure D. 2003. The Ti plasmid of *Agrobacterium tumefaciens* harbors an attM-paralogous gene, *aiiB*, also encoding N-Acyl homoserine lactonase activity. *Applied and Environmental Microbiology*. 69(8): 4989-4993.
- Cerón J., Ortiz A., Quintero R., Guereca L., Bravo A. 1995. Specific PCR primers directed to identify *cryI* and *cryIII* genes within a *Bacillus thuringiensis* strain collection. *Applied and Environmental Microbiology*. 61(11): 3826-3831.
- Cha C., Gao P., Chen Y.C., Shaw P.D., Farrand S.K. 1998. Production of acyl-homoserine lactone quorum-sensing signals by gram negative plant-associated bacteria. *Molecular Plant-Microbe Interactions*. 11(11): 1119-1129.
- Charkowski A.O. 2009. Decaying signals: will understanding bacterial-plant communications lead to control of soft rot?. *Current opinion in Biotechnology*. 20(2): 178-184.
- Chatterjee A., Cui Y., Hasegawa H., Leigh N., Dixit V., Chatterjee A.K. 2005. Comparative analysis of two classes of quorum-sensing signaling systems that control production of extracellular proteins and secondary metabolites in *Erwinia carotovora* subspecies. *Journal of Bacteriology*. 187(23): 8026-8038.
- Choudhary S., Schmidt-Dannert C. 2010. Applications of quorum sensing in biotechnology. *Applied Microbiology and Biotechnology*. 86(5): 1267-1279.
- Chu W., Vatter D.A., Maitin V., Barnes M.B., McLean R.J.C. 2011. Bioassays of Quorum Sensing Compounds using *Agrobacterium tumefaciens* and *Chromobacterium violaceum*. In: Quorum Sensing: Methods and Protocols, Methods in Molecular Biology. Rumbaugh, K.P. (Ed). Springer Science-Business Media.
- Dong Y.H., Xu J.L., Li X.Z., Zhang L.H. 2000. *AiiA*, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proceedings of the National Academy of Sciences*. 97(7): 3526-3531.
- Dong Y.H., Wang L.H., Xu J.L., Zhang H.B., Zhang X.F., Zhang L.H. 2001. Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature*. 411(6839): 813-817.
- Dong Y.H., Gusti A.R., Zhang Q.J., Xu L. Zhang L.H. 2002. Identification of quorum-quenching N-acyl homoserine lactonases from *Bacillus* species. *Applied and Environmental Microbiology*. 68(4): 1754-1759.
- Dong Y.H., Zhang X.F., Xu J.L., Zhang L.H. 2004. Insecticidal *Bacillus thuringiensis* silences *Erwinia carotovora* virulence by a new form of microbial antagonism, signal interference. *Applied and Environmental Microbiology*. 70(2): 954-960.
- Fuqua W.C., Winans S.C., Greenberg E.P. 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *Journal of Bacteriology*. 176(2): 269-275.
- Hall T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. In: *Nucleic Acids Symposium Series*. 41: 95-98.
- Hong S.H., Hegde M., Kim J., Wang J., Jayaram A., Wood T.K. 2012. Synthetic quorum-sensing circuit to control consortial biofilm formation and dispersal in a microfluidic device. *Nature Communications*. 3: 613. doi:10.1038/ncomms1616
- Jara S., Maduell P., Orduz S. 2006. Diversity of *Bacillus thuringiensis* strains in the maize and bean phylloplane and their respective soils in Colombia. *Journal of applied microbiology*. 101(1): 117-124.
- Jones S., Yu B., Bainton N.J., Birdsall M.B., Bycroft W., Chhabra S.R., Cox A.J.R., Golby P., Reeves P.J., Stephens S., Winson M.K., Salmond G.P.C., Stewart G.S.A.B., Williams P. 1993. The lux autoinducer regulates the production of exoenzyme virulence determinants in *Erwinia carotovora* and *Pseudomonas aeruginosa*. *The EMBO Journal*. 12(6): 2477-2482.
- Kado C.I., Heskett M.G. 1970. Selective media for isolation of *Agrobacterium*, *Corynebacterium*, *Erwinia*, *Pseudomonas*, and *Xanthomonas*. *Phytopathology*. 60: 969-976.
- Kalia C.V. 2013. Quorum sensing inhibitors: An overview. *Biotechnology Advances*. 31(2): 224-245.
- Kim M.H., Kang H.O., Kang B.S., Kim K.J., Choi W.C., Oh T.K., Lee C.H., Lee J.K. 2005. Crystallization and preliminary crystallographic analysis of *Bacillus thuringiensis* AHL-lactonase. *Biochimica et Biophysica Acta (BBA)-Proteins and Proteomics*. 1750(1): 5-8.
- Leadbetter J.R., Greenberg E.P. 2000. Metabolism of acyl-homoserine lactone quorum-sensing signals by *Variovorax paradoxus*. *Journal of Bacteriology*. 18: 6921-6926.
- Lee S.J., Park S.Y., Lee J.J., Yum D.Y., Koo B.T., Lee J.K. 2002. Genes encoding the N-acyl homoserine lactone-degrading enzyme are widespread in many subspecies of *Bacillus thuringiensis*. *Applied and Environmental Microbiology*. 68(8): 3919-3924.
- Lin Y.H., Xu J.L., Hu J., Wang L.H., Ong S.L., Leadbetter J.R., Zhang L.H. 2003. Acyl-homoserine lactone acylase from *Ralstonia* strain XJ12B represents a novel and potent class of quorum-quenching enzymes. *Molecular Microbiology*. 47(3): 849-860.
- Liu H., Coulthurst S.J., Pritchard L., Hedley P.E., Ravensdale M., Humphris S., Burr T., Takle G., Brurberg M.B., Birch P.R., Salmond G.P., Toth I.K. 2008. Quorum sensing coordinates brute force and stealth modes of infection in the plant pathogen *Pectobacterium atrosepticum*. *PLoS Pathogens*. 4(6): e1000093.
- McClellan K.H., Winson M.K., Fish L., Taylor A., Chhabra S.R., Camara M., Daykin M., Lamb J.H., Swift S., Bycroft B.W., Stewart G.S., Williams P. 1997. Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. *Microbiology*. 143(12): 3703-3711.
- Park S.Y., Lee S.J., Oh T.K., Oh J.W., Koo B., Yum D.Y., Lee J.K. 2003. AhLD, an N-acylhomoserine lactonase in *Arthrobacter* sp., and predicted homologues in other bacteria. *Microbiology*. 149(6): 1541-1550.
- Pérombelon M.C., Kelman A. 1980. Ecology of the Soft Rot *Erwinias*. *Annual Review of Phytopathology*. 18(1): 361-387.
- Pedroza C.J., Flórez A.M., Ruiz, O.S., Orduz S. 2014. Enzymatic hydrolysis of molecules associated with bacterial quorum sensing using an acyl homoserine lactonase from a novel *Bacillus thuringiensis* strain. *Antonie van Leeuwenhoek*. 105:253-264
- Pirhonen M., Flego D., Heikinheimo R., Palva E.T. 1993. A small diffusible signal molecule is responsible for the global control of virulence and exoenzyme production in the plant pathogen *Erwinia carotovora*. *The EMBO Journal*. 12(6): 2467-2476.
- Ravn L., Christensen A.B., Molin S., Givskov M., Gram L. 2001. Methods for detecting acylated homoserine lactones produced by gram negative bacteria and their application in studies of AHL-production kinetics. *Journal of Microbiological Methods*. 44(3): 239-251.
- Saitou N., Nei M. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*. 4(4): 406-425.
- Schaad N.W., Jones J.B., Chun W. 2001. Laboratory Guide of Identification of Plant Pathogenic Bacteria. APS Press (Ed). St. Paul, Minnesota, USA.
- Sio C.F., Otten L.G., Cool R.H., Diggle S.P., Braun P.G., Bos R., Daykin M., Camara M., Williams P., Quax W.J. 2006. Quorum quenching by an N-acyl-homoserine lactone acylase from *Pseudomonas aeruginosa* PAO1. *Infection and immunity*. 74(3): 1673-82.
- Smadja B., Latour X., Faure D., Chevalier S., Dessaux Y., Orange N. 2004. Involvement of N-acylhomoserine lactones throughout plant infection by *Erwinia carotovora* subsp. *atroseptica* (*Pectobacterium atrosepticum*). *Molecular Plant-Microbe Interactions*. 17(11): 1269-1278.

- Tamura K., Nei M., Kumar S. 2004. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences of the United States of America*. 101(30): 11030-11035.
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M., Kumar S. 2011. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution*. 28(10): 2731-2739.
- Tempé J., Petit A., Holsters M., Van Montagu M., Schell J. 1977. Thermosensitive step associated with transfer of the Ti plasmid during conjugation: Possible relation to transformation in crown gall. *Proceedings of the National Academy of Sciences of the United States of America*. 74(7): 2848-2849
- Thomas P.W., Stone E.M., Costello A.L., Tierney D.L., Fast W. 2005. The quorum-quenching lactonase from *Bacillus thuringiensis* is a metalloprotein. *Biochemistry*. 44(20): 7559-7569.
- Tinh N.T.N., Yen V.H.N., Dierckens K., Sorgeloos P., Bossier P. 2008. An acyl homoserine lactone-degrading microbial community improves the survival of first-feeding turbot larvae (*Scophthalmus maximus* L.). *Aquaculture*. 285(1): 56-62.
- Uribe D., Martínez W., Cerón J. 2003. Distribution and diversity of *cry* genes in native strains of *Bacillus thuringiensis* obtained from different ecosystems from Colombia. *Journal of Invertebrate Pathology*. 82(2): 119-127.
- Von Bodman S.B., Bauer W.D., Coplin D.L. 2003. Quorum sensing in plant pathogenic bacteria. *Annual Review of Phytopathology*. 41(1): 455-482.
- Wang L.H., Weng L.X., Dong Y.H., Zhang L.H. 2004. Specificity and enzyme kinetics of the quorum-quenching N-Acyl homoserine lactone lactonase (AHL-lactonase). *Journal of Biological Chemistry*. 279(14): 13645-13651.
- Williams P. 2007. Quorum sensing, communication and cross-kingdom signalling in the bacterial world. *Microbiology*. 153(12): 3923-39238.
- Williams P., Camara M. 2009. Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Current Opinion in Microbiology*. 12(2): 182-191.
- Winson M.K., Swift S., Fish L.J., Throup P., Jorgensen F., Chhabra S.R., Bycroft B.W., Williams P., Stewart G. 1998. Construction and analysis of luxCDABE-based plasmid sensors for investigating N-acyl homoserine lactone-mediated quorum sensing. *FEMS Microbiology Letters*. 163(2): 185-192.
- Zhang H.B., Wang L.H., Zhang L.H. 2007. Detection and Analysis of Quorum-Quenching Enzymes against Acyl Homoserine Lactone Quorum-Sensing Signals. *Current Protocols in Microbiology*. 5:1C.3.1-1C.3.15. DOI: 10.1002/9780471729259.mc01c03s05