

Epidural analgesia in laparoscopic radical prostatectomy

A. Tejedor Navarro, M. García Martínez, M. Vendrell Jordà, E. Gómez Lanza, S. García Aranda and J. Masdeu Castellví

Hospital Sant Joan Despí Moisès Broggi. Barcelona. Spain

Tejedor Navarro A, García Martínez M, Vendrell Jordà M, Gómez Lanza E, García Aranda S and Masdeu Castellví J. Epidural analgesia in laparoscopic radical prostatectomy. Rev Soc Esp Dolor 2018;25(1):13-20.

ABSTRACT

Introduction: Epidural analgesia is an effective technique for postoperative pain relief. Our aim in this retrospective study was to assess the postoperative pain control and complications relating to epidural technique in laparoscopic radical prostatectomy (LRP).

Material and methods: A retrospective analysis of 193 patients who underwent LRP, in which epidural analgesia was the postoperative pain approach, was performed. The procedure is generally performed under combined anesthesia. Data collected was postoperative pain data; appearance of neurological disorders (Bromage scale was used for motor weakness assessment); data related to the epidural technique and possible difficulties when performing it; data on epidural infusion; catheter-related complications; hospital stay, postoperative complications, and outcome.

Results: Firstly, average VAS at rest was 1.2 ± 1.6 ; and upon movement, average VAS was 1.9 ± 1.8 during the hospital stay. Secondly, complications related to epidural technique appeared in 37 % of patients. There were 3 cases of hematic puncture; 3 accidental catheter disconnections; 1 dural puncture, and 1 subdural block. As for neurological secondary effects of local anesthetics in the epidural technique, 56 patients (30.1 %) presented with motor block of one or both lower extremities after surgery, and 5 (2.7 %) with paresthesia. Statistical analysis showed that motor weakness was not related to age, weight, type of local anesthetic used, infusion rate, level of epi-

dural puncture nor length of catheter within the epidural space ($p > 0.05$).

Conclusion: Postoperative epidural analgesia offers excellent analgesic quality but it can be associated with several complications secondary to the use of local anesthetics, which could disagree with the terms of Fast-track surgery. New techniques like the TAP block could offer the same analgesic quality, without the epidural's technique potential complications.

Key words: Laparoscopic prostatectomy, epidural analgesia, postoperative pain, neurologic complications.

RESUMEN

Objetivos: La analgesia epidural es una técnica eficaz para el control del dolor postoperatorio. Nuestro objetivo en este estudio retrospectivo fue evaluar el control del dolor postoperatorio mediante la escala visual analógica del dolor (EVA) y las complicaciones relacionadas con la técnica epidural, en la prostatectomía radical laparoscópica (PRL).

Material y métodos: Llevamos a cabo un análisis retrospectivo de 193 pacientes sometidos a PRL, en los que se realizó la técnica epidural analgésica para el control del dolor postoperatorio. El procedimiento se hizo bajo una anestesia combinada. Registramos los datos relacionados con el dolor postoperatorio; la aparición de sintomatología neurológica (la escala de Bromage se utilizó para evaluar la debilidad motora); datos relacionados con la técnica epidural y posibles dificultades al realizarla; datos sobre la infusión epidural, como tipo de anestésico local utilizado; complicaciones relacionadas con el catéter y complicaciones postoperatorias asociadas al mismo, estancia hospitalaria y resultado.

Resultados: En primer lugar, durante la estancia hospitalaria de los pacientes, el EVA promedio en reposo fue $1,2 \pm 1,6$; y durante el movimiento, el EVA promedio fue de $1,9 \pm 1,8$. En

segundo lugar, las complicaciones relacionadas con la técnica epidural aparecieron en el 37 % de los pacientes. Hubo 3 casos de punción hemática; 3 desconexiones accidentales del catéter, 1 punción dural y 1 bloqueo subdural. En cuanto a las complicaciones neurológicas debidas a los efectos secundarios de los anestésicos locales en la técnica epidural, 56 (30,1 %) pacientes presentaron bloqueo motor de una o ambas extremidades inferiores después de la cirugía y 5 (2,7 %) refirieron parestesias. No hubo ninguna complicación neurológica que persistiese tras el alta hospitalaria. El análisis estadístico mostró que la debilidad motora no estaba relacionada con la edad, el peso, el tipo de anestésico local utilizado, la velocidad de infusión, el nivel de punción epidural ni la longitud del catéter en el espacio epidural ($p > 0,05$).

Conclusión: La analgesia epidural postoperatoria ofrece una excelente calidad analgésica, pero puede estar asociada a varias complicaciones secundarias al uso de anestésicos locales, lo que podría estar en contraposición con las tendencias actuales de cirugía fast-track. Las nuevas técnicas emergentes podrían ofrecer la misma calidad analgésica evitando las potenciales complicaciones de la técnica epidural.

Palabras clave: Prostatectomía laparoscópica, analgesia epidural, dolor postoperatorio, complicaciones neurológicas.

INTRODUCTION

Laparoscopic radical prostatectomy (LRP) is considered the standard treatment for clinically localized prostate cancer.

LRP pain is lesser than that of open surgery, but it is still considered as moderate (1). Literature results on the optimal analgesic technique are conflicting (2-5), with actual guidelines encouraging a multimodal approach (6) to reduce the amount of opioids administered, thus minimizing side effects (2,7). Several studies have been published, where different analgesic methods are considered: patient-controlled analgesia (PCA) of morphine chloride (2), epidural technique (3), use of drugs for neuropathic pain control (4), or surgical wound infiltration. However, there is no clear consensus on the best analgesic technique for this type of surgery.

The benefits of optimal postoperative analgesia are clear, and include a reduction in the postoperative stress response, which can lead to organ dysfunction and prolong recovery time (8). Appropriate analgesia translates in lesser postoperative morbidity and improved surgical outcomes, accelerating rehabilitation without increasing the number of readmissions or complications (9).

Hence, the usual practice in our hospital is the epidural technique with local anesthetics, which has been considered the most adequate analgesic technique for this type of surgery, as it improves pain control and patient comfort, decreasing secondary effects of other analgesics, and the number of complications (7,10). However, this technique

has certain contraindications and is associated to complications such as numbness or motor weakness (11,12), requiring close monitoring to ensure its safety and efficacy, not delaying the rehabilitation period. Moreover current practices advocate in favor of a less invasive analgesic therapy, which seems to be sufficient for this kind of surgical approach.

As part of a larger, prospective study comparing the analgesic efficacy of epidural technique vs. TAP technique for PRL pain, we performed this retrospective study of all epidural-controlled LRP performed in our hospital between 2010 and 2015. Our aim was to describe the analgesic quality of this technique, as well as its complications in the postoperative period.

MATERIAL AND METHODS

A retrospective analysis of all patients who underwent LRP for prostate cancer between 2010 and 2015, in which epidural analgesia was the postoperative approach, was performed. This analysis was carried out in the design of a study comparing the epidural technique with another analgesic technique. We performed the analysis of the last 5 years to see the results obtained and to be able to evaluate them with those that will be obtained in the study that is currently in progress. Exclusion criteria included those patients where regional anesthesia was contraindicated (due to patient's rejection or lack of collaboration, bleeding disorders, including heparin use, puncture site infection, allergy to local anesthetics, neurological disorders), and those patients where surgery was reconverted from LRP to open surgery.

Management of patients undergoing LRP

The procedure is generally performed under combined anesthesia, as follows. Under standard monitoring (electrocardiogram, pulse oximetry and noninvasive blood pressure), a peripheral venous access is obtained and intravenous midazolam (1-2 mg) is administered for anxiolysis.

In those patients without contraindications (13), epidural technique with loss of resistance (serum or air) approach is performed at either thoracic or lumbar level, depending on each anesthesiologist's usual practice. A test dose of 3 mL epidural 2 % lidocaine is given.

The patient is transorally intubated, after anesthetic induction is performed with fentanyl (1.5 mcg/kg), propofol (1.5-2 mg/kg) and rocuronium (0.6 mg/kg). Maintenance is performed with sevoflurane (CAM 0.6-1). For analgesic control, an initial epidural bolus of between 6 and 8 mL levobupivacaine 0.25 % is administered, with further hourly bolus of between 5 and 8 mL, depending on pain assessment by usual parameters.

After surgery, the patient is extubated and transferred to the postoperative acute care unit (PACU), where he remains for the first 12 hours after surgery, for pain and bleeding control. An infusion of local anesthetic (LA) (levobupivacaine 0.125 % or bupivacaine 0.125 % with or without fentanyl 2 mcg/mL) is begun and maintained at either 5 or 8 mL/h (elastomeric pump reservoir). If motor blockade or hypotension due to vasoplegia is detected, the infusion is suspended until reversal of effects. On the other hand, in case of uncontrolled pain, a bolus of 5 mL 1 % lidocaine is administered, with reposition of epidural catheter if there is no improvement.

It has been demonstrated that postoperative pain control is effective both as PCEA (Patient-controlled epidural analgesia) (14,15) as in CEI (continuous epidural infusion) (16,17) mode. As PCEA pumps are not available in our center, CEI was used.

Intravenous postoperative analgesia consists in paracetamol 1 g/8 h ev and dexketoprofen 50 mg/8 h ev. In situations of renal impairment, gastric problems or allergies, metamizol 2 g/8 h ev was administered.

During the post-operative period, the anesthesiologist and the nurse from the acute pain unit (APU) visit the patients daily, evaluating and registering catheter function, pain, and possible complications. Catheter removal is performed when VAS were consistently <3 in patients with low doses of epidural analgesia, without blood coagulation alterations nor low molecular weight heparin (LMWH) administration within the previous 6 hours.

Data Collection

Data was obtained via retrospective review of clinical histories, including the anesthesia sheet, nurse's charts and urology charts, and the APU's registries. From these, data on the epidural technique, drug administration, general evolution and possible complications was obtained.

According to the previous description, the following data was recorded: anthropometric parameters (age, weight, height, ASA); data related to the epidural technique (distance to skin of epidural space, length of catheter inserted) and possible difficulties when performing it (vascular puncture, spinal puncture, impossibility of performance, subdural block); data on epidural infusion (type of LA administered, infusion rate); catheter-related complications (accidental disconnection); postoperative pain data (visual analog scale (VAS) at rest and in movement at PACU, and daily); appearance of neurological side effects (motor block according to the Modified Bromage scale: *I: no block: free movement of legs and feet; II: Partial block: just able to flex knees with free movement of feet; III: Almost complete block: unable to flex knees, but with free movement of feet; IV: complete block: unable to move legs or feet*); postoperative complications and total length of admission.

Data analysis

The collected data was analyzed using SPSS version 22.0 program. The results are expressed as mean \pm SD or percentages and range. To analyze and compare the variables, non-parametric tests (Kruskal-Wallis, Mann-Whitney) were applied. Results were accepted as statistically significant when $p < 0.05$.

RESULTS

One hundred ninety-three medical records were reviewed. Six of them were excluded due to partial absence of data and one of them was excluded due to epidural catheter malfunction, requiring a switch to intravenous morphine for pain treatment.

Average surgery time was 161 ± 71 minutes (110-500 minutes, range). Anthropometric results and epidural technique results are available in Tables I and II. Epidural infusion, rate and type of LA varied depending on the anesthesiologist's initial prescription and subsequent adjustments depending on the pain and side effects. LA used was levobupivacaine 0.125 % in 156 cases (85 % total); with fentanyl in 87 cases (46.8 % total) and without in 71 cases (38.2 % total). In the 28 (15 %) remaining cases, LA was bupivacaine, with fentanyl in 16 (8.6 %) and without in 12 (5.9 %) cases. In brief, in 55.4 % of the cases, LA + fentanyl was used, and LA without fentanyl in the remainder cases. No statistically significant differences in motor blockade between groups were seen ($p > 0.05$). Average infusion rate was $5.12 \text{ mL/h} \pm 1.1 \text{ mL/h}$.

Complications related to epidural technique appeared in 37 % of patients. Of all the technique-related compli-

TABLE I
ANTHROPOMETRIC RESULTS OBTAINED FROM STUDIED SUBJECTS, DATA FROM ALL PATIENTS WHO UNDERWENT LAPAROSCOPIC RADICAL PROSTATECTOMY FOR PROSTATE CANCER BETWEEN 2010 AND 2015 WITH EPIDURAL ANALGESIA USED FOR POSTOPERATIVE PAIN TREATMENT

Age ($X \pm SD$) years	66 \pm 6	
Weight ($X \pm SD$) kg	78.8 \pm 10.5	
Height ($X \pm SD$) cm	169 \pm 70	
ASA score (%)	I	4.3 %
	II	85.5 %
	III	10.2 %

X: mean. SD: standard deviation. ASA: American Society of Anesthesiology.

cations there were 3 cases (1.5 %) of hematic puncture, and 1 (0.5 %) adverted dural puncture, in which technique was successfully repeated without further complications; 3 (1.5 %) accidental catheter disconnections after the first 24 h, where conventional analgesia was continued, as pain was well-controlled; and 1 (0.5 %) subdural block, with catheter removal and intravenous morphine used for pain relief (data relative to this patient was included up until that point). The subdural blockade and the dural puncture cases were strictly followed-up by the APU nurse, and no complications were observed.

TABLE II
EPIDURAL TECHNIQUE RESULTS OBTAINED FROM STUDIED SUBJECTS, DATA FROM ALL PATIENTS WHO UNDERWENT LRP FOR PROSTATE CANCER BETWEEN 2010 AND 2015 WITH EPIDURAL ANALGESIA USED FOR POSTOPERATIVE PAIN TREATMENT

<i>Epidural puncture level (%)</i>	≥ T12	25.8 %
	L1-L2	47.3 %
	L2-L3	18.3 %
	L3-L4	8.6 %
<i>Epidural space (X±SD, [range]) cm</i>	5.3 ± 0.9 [3-8]	
<i>Length of catheter left in epidural space (X ± SD, [range]) cm</i>	4.4 ± 0.8 [3-7]	
<i>Epidural complications</i>	69 (37 %) yes / 117 (63 %) no	

X: mean. SD: standard deviation.

As for neurological side effects, 56 patients (30.1 %) presented with motor block of one or both lower extremities after surgery, and 5 (2.7 %) with paresthesia. When talking specifically of the 56 patients with motor blockade, in 42 cases (75 %) it was a unilateral Bromage I degree block; and in 14 patients (25 %) it was a bilateral block (8 Bromage I and 6 Bromage II) (Table III).

The highest frequency of motor weakness was observed during the first 6 to 12 hours after surgery. The degree of motor block in relation to the level of epidural puncture is shown in Table III. Motor weakness was greater in the epidural lumbar punctures than in thoracic level punctures, although results were not statistically significant. In the 42 cases of unilateral motor blockade (see previous results), the catheter was removed 0.5-1 cm, if possible, and the patient was placed on his side, with the blocked side on top. In all cases blockade was reversed. In the cases of bilateral block, the 8 Bromage grade I blocks were treated by decreasing the LA infusion rate; in the remaining 6 bilateral Bromage II patients, infusion was suspended until improvement, which was complete in all cases. In most cases catheter mobilization was performed at the PACU. In those cases of bilateral block, drug infusion was suspended until reversal of motor blockade was observed, at which point infusion was re-started at lower rates. These patients were controlled strictly by the APU nurse.

Postoperative pain

Patients were followed during the time epidural catheter was in place. Average VAS at rest (r) and upon movement (m) in PACU and on first, second and third day was r 0.7 ± 1.2, m 2.1 ± 1.7; r 0.8 ± 1.3, m 1.9 ± 0.9; r 0.61 ± 1.2, m 2.01 ± 1.7; and r 0.8 ± 1.1, m 1.78 ± 2; respectively

TABLE III
INCIDENCE AND DEGREE OF MOTOR BLOCKADE DEPENDING ON EPIDURAL LEVEL IN STUDIED SUBJECTS

<i>Epidural puncture level</i>	<i>N</i>	<i>Incidence of motor weakness</i>	<i>Bromage scale</i>			
			<i>I</i>	<i>II</i>	<i>III</i>	<i>IV</i>
T7-T8	3	0	-	-	-	-
T8-T9	2	0	-	-	-	-
T9-T10	4	1 (1.8 %)	1	-	-	-
T10-T11	5	2 (3.6 %)	2	-	-	-
T11-T12	11	1 (1.8 %)	1	-	-	-
T12-L1	23	5 (8.9 %)	5	-	-	-
L1-L2	88	31 (55.5 %)	28	3	-	-
L2-L3	34	11 (19.6 %)	9	2	-	-
L3-L4	16	5 (8.9 %)	4	1	-	-

T: thoracic. L: lumbar.

(Figure 1). Overall, average VAS at rest was 1.2 ± 1.6 ; and upon movement, average VAS was 1.9 ± 1.8 .

Epidural catheter remained in place for over 72h in only 5 % of the cases. 26 (14 %) patients required epidural analgesia for 24 h, 106 (57 %) for 48 h, and 44 (23.5 %) for 72 h. Of the 10 patients (5 %) were the catheter was maintained for over 96 h; in 8 cases it was due to surgical revision; and in 2 patients the infusion was stopped but catheter couldn't be removed due to abnormal clotting parameters due to excessive bleeding.

Postoperative care

Ten cases required hospital admission above 10 days, due to surgical complications. These were 4 cases of urinary leakage, 2 urinary tract infection cases, 1 surgical wound infection, 1 case of pulmonary embolism, 1 bladder dome section that coursed with anuria, and 1 paralytic ileus. There were 5 cases of postoperative bleeding treated conservatively. Length of hospital stay was 5.98 ± 2.92 (minimum 2 days and maximum 30 days).

Statistical analysis showed that motor weakness was not related to age, weight, type of LA used, addition of fentanyl, infusion rate, level of epidural puncture nor length of catheter within the epidural space ($p > 0.05$).

DISCUSSION

In LRP, the pain is defined as moderate (1). With the goal of increasing the welfare of postoperative patients undergoing this type of intervention, studies comparing open versus laparoscopic technique have been conducted, with inconclusive results. Results by Webster et al. (18) concluded that necessity of analgesics diminished if it the procedure


is laparoscopic; whereas D'Alonzo et al. (19) did not find statistically significant differences between the two techniques in terms of opioid consumption.

Systematic reviews reveal a significant lack of evidence in the pain management protocol of patients undergoing LRP (5). Most studies evaluate pain under unimodal analgesic management (20), not multimodal management, as is the current trend. Moreover, while it is accepted that LPR decreases postoperative pain, studies are few. It is therefore necessary to determine the optimal analgesic technique for this type of surgery (2-4). Studies have shown that pain in this type of surgery can be controlled with opioids, but because of their side effects (2,7), nowadays other analgesic techniques are advocated.

Since the opening of our center in 2010, control of postoperative pain in the LRP has been performed using lumbar epidural analgesia. The epidural technique allows for optimal pain control, improving patient comfort (8), reducing side effects of other analgesics (2,7), and the number of complications such as postoperative infarction, bleeding, pulmonary comorbidities, respiratory depression, kidney failure, and deep vein thrombosis (21-25).

When evaluating our results (Figure 1), the average VAS at rest (1.2 ± 1.6) and on movement (1.9 ± 1.8) remained below 3 during follow-up in all patients. Although studies have been published that correlate younger age with a higher pain score (26), our VAS records were excellent, in spite of an average age of 66 years. However, the complications associated with this technique must be kept in mind: catheter infection, post-dural puncture headache, stroke, intracranial hematoma, cerebral thrombosis, meningitis, and neurological disorders (11,27,28). No major complications were recorded, but thirty-seven per cent of analyzed patients presented some type of minor complication.

As summarized in Table 3, 56 patients (30.1 %) had motor weakness, especially during the first 12 hours after surgery, which disappeared with standard maneuvers such as change in patient's position and / or reduction or suspension of LA infusion. These results are similar to those published by Broekedema (16) et al. and Ahmed (27) et al. Broekedema (16) studied surgeries performed with either lumbar or thoracic epidural. A single motor blockade was detected in a patient with a lumbar epidural. Numbness could be detected in patients with either lumbar or low thoracic epidurals, with symptoms disappearing when infusion rate decreased. In the study by Ahmed et al. (27), 36.5 % of patients with local anesthetic epidural infusion presented motor weakness. The most likely cause of this initial motor blockade is attributed to the persistence of effect of LA administered during the surgery (12). Regarding the incidence of motor block according to puncture level, we observed that this is higher at lumbar (74.2 %) versus thoracic (25.8 %) level ($p > 0.05$); being these results similar to those described in other studies (11,27). This is thought to be due to anatomical reasons.



VAS: visual analog scale.

Fig. 1. Daily VAS in patients with epidural catheter for postoperative pain management of laparoscopic radical prostatectomy, evaluated upon repose and movement.

In addition to motor block, 5 cases of paresthesias, 3 hematic punctures, 3 accidental catheter disconnections (with consequent sterility loss) and 1 dural puncture were recorded. In 2 of the 5 cases of postoperative bleeding, the catheter was not removed until the 5th day due to coagulation disorders, and in the case of bladder dome section, it wasn't removed until the 8th day. Although risk of infection due to prolonged catheter placement is described as low (28), it nonetheless increases when catheter isn't timely removed. Vascular punctures, unnoticed in 9% of the cases (29), can lead to disastrous consequences (hemodynamic disturbances, heart failure). Other complications include urinary retention (28) or direct spinal cord injuries. For this reason, several studies advocate for other analgesic techniques (30,31).

Even if epidural technique covers pain in LRP excellently, the current trend is toward drain-less, less invasive surgery. This tendency raises the question of whether such analgesic technique is really necessary for lower abdominal surgeries, taking into account not only possible complications but also increased costs associated with epidural follow-up (32). Moreover, despite the advantages of the epidural technique, there is a lack of evidence to support its superiority over intravenous analgesia. While there are studies that advocate the benefits of epidural analgesia technique (33), others find no superiority in comparison to systemic analgesia (20).

In addition to this, we must remember that the current trend is the implementation of Fast-track protocols, in order to obtain patient's early discharge and a return to normal life as quickly possible (1,8,9).

Fast-track protocols are new multimodal approaches that deviate from standard treatments that include an increased fluid administration, use of nasogastric tubes and post-operative drains, prolonged postoperative oral intake restriction and immobilization; in order to favor early discharge and decrease hospital length-of-stay (34). The fact that LRP is not a bowel procedure favors the implementation of Fast-track therapy. As well as the use of minimally invasive techniques associated to a decrease in inflammatory response and immunological dysfunctions, opioid-free anesthetic and / or analgesic techniques are advocated, minimizing postoperative complications and facilitating recovery (8,9,35). Early-recovery-after surgery (ERAS) protocols have been included in various surgical paths (36-38), with scarce bibliography in urologic surgeries (39).

After reviewing the published literature, we decided to carry out a study comparing the epidural technique with another analgesic technique. In this context, we decided to evaluate the data of the epidural technique of the previous 5 years.

Epidural technique fulfills the requirements for analgesia in Fast-track surgery, providing optimal pain control with minimal opioid administration, as seen in our study and in the literature. However, it has its drawbacks, as it

may hinder early mobilization when paresthesia, motor blockade or other complications appear; and hospital costs related to monitoring of these complications increase. Neural block, not only epidural technique but any local technique that blocks nerve impulses, inhibits the endocrine and sympathetic response to surgery, more effectively in lower than in upper abdominal surgery (40). Thus, epidural benefits can be obtained with other neural blockade analgesic techniques, avoiding possible complications.

Our average length-of-stay is 5.98 ± 2.92 days. Magheli et al. found that with the implementation of the Fast-track protocol in the LRP, hospital stay was reduced from 7 days to 3.7 days, and complication rate decreased (1). However, recent studies demonstrate that epidural analgesia increases length-of-stay (41,42). We cannot assure that the average hospital stay could have been lower if another analgesic technique had been performed, but after analyzing the complications obtained would be an important factor to take into account. To clarify these data, comparative studies between the epidural technique and new anesthetic techniques would be necessary, which is what we are currently undertaking. For example, several studies supporting TAP block as an analgesic technique in colorectal surgery have been published (30,31). McDonnell et al. (43) evaluated the TAP blockade in colorectal surgery and obtained very good pain control. In a meta-analysis, which evaluated the TAP blockade as analgesic technique in laparoscopic surgery, it concluded that TAP technique is an effective analgesic technique in patients undergoing laparoscopic Fast-track surgery, providing analgesic quality without associated complications (44). However, specific literature regarding TAP block in LRP is scarce.

Our study has some limitations. Being an observational retrospective study, anesthesia management of patients did not follow strict protocols and this produces an important bias. Furthermore, data was collected retrospectively, with some data missing, compromising data analysis and reducing power of statistical conclusions. Motor blockade was greater in those patients with lumbar catheters than those with thoracic catheters, as has been previously described (12,25,27), but we can only speak of tendencies, as there was no technique randomization, and results may be subject to bias. The same is applicable when considering the use of fentanyl.

Ideally, the epidural technique should be compared to the new emerging analgesic techniques, as retrospective studies hinder the association of causality between variables. We are currently carrying out a comparative, prospective study, to reduce these bias.

CONCLUSION

To conclude, the epidural technique offers an excellent analgesic quality, but complications and/or side effects

associated to the use of LAs disagree with the terms of Fast-track surgery, with risk of increased length-of-stay. Other analgesic techniques may offer the same pain management without the complications of the epidural technique.

CONFLICT OF INTEREST

The authors declare they have no conflicts of interest.

REFERENCES

- Magheli A, Knoll N, Lein M, Hinz S, Kempkensteffen C, Gralla O. Impact of Fast-track postoperative care on intestinal function, pain, and length of hospital stay after laparoscopic radical prostatectomy. *J endourol* 2011;25(7):1143-7. DOI: 10.1089/end.2011.0020.
- Senagore A, Delaney C, Mekhail N, Dugan A, Fazio VW. Randomized clinical trial comparing epidural anaesthesia and patient-controlled analgesia after laparoscopic segmental colectomy. *Br J Surg* 2003;90(10):1195-9.
- Costello TG, Webb P. Anaesthesia for robot-assisted anatomic prostatectomy. Experience at a single institution. *Anaesth Intensive Care* 2006;34(6):787-92.
- Trabulsi E, Patel J, Viscusi E, Gomella LG, Lallas CD. Preemptive multimodal pain regimen reduces opioid analgesia for patients undergoing robotic-assisted laparoscopic radical prostatectomy. *Urology* 2010;76(5):1122-4. DOI: 10.1016/j.urology.2010.03.052.
- Joshi GP, Jaschinski T, Bonnet F, et al. Optimal pain management for radical prostatectomy surgery: what is the evidence? *BMC Anesthesiology* 2015;15:159. DOI: 10.1186/s12871-015-0137-2.
- American Society of Anesthesiologists Task Force on Acute Pain Management. Practice guidelines for acute pain management in the perioperative setting: an updated report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology* 2012;116(2):248-73. DOI: 10.1097/ALN.0b013e31823c1030.
- Gottschalk A, Smith DS, Jobs DR, Kennedy SK, Lally SE, Noble VE, et al. Preemptive epidural analgesia and recovery from radical prostatectomy: a randomized controlled trial. *JAMA* 1998;279(14):1076-82.
- Power I, Barratt S. Analgesic agents for the postoperative period. *Nonopioids*. *Surg Clin North Am* 1999;79(2):275-97.
- Lassen K, Soop M, Nygren J, Cox PB, Hendry PO, Spies C, et al; Enhanced Recovery After Surgery (ERAS) Group. Consensus review of optimal perioperative care in colorectal surgery: Enhanced Recovery After Surgery (ERAS) Group recommendations. *Arch Surg* 2009;144(10):961-9. DOI: 10.1001/archsurg.2009.170.
- Ben-David B, Swanson J, Nelson J, et al. Multimodal analgesia for radical prostatectomy provides better analgesia and shortens hospital stay. *J Clin Anesth* 2007;19(4):264-8.
- Königsrainer I, Bredanger S, Drewel-Frohnmeyer R, et al. Audit of motor weakness and premature catheter dislodgement after epidural analgesia in major abdominal surgery. *Anesthesia* 2009;64(1):27-31. DOI: 10.1111/j.1365-2044.2008.05655.x.
- Merson N. A comparison of motor block between ropivacaine and bupivacaine for continuous labor epidural analgesia. *AANA J* 2001;69(1):54-8.
- Brull R, Macfarlane A, Chan V. Spinal, epidural, and caudal anesthesia. In: Roland D. Miller, ed. *Miller's Anesthesia 8th Edition*. Philadelphia: Churchill Livingstone; 2015.
- Standl T, Burmeister M, Ohnesorge H, Wilhelm S, Striepke M, Gottschalk A, et al. Patient-controlled epidural analgesia reduces analgesic requirements compared to epidural infusion after major abdominal surgery. *Can J Anesth* 2003;50(3):248-64.
- Liu SS, Allen HW, Olsson GL. Patient-controlled epidural analgesia with bupivacaine and fentanyl on hospital wards. Prospective experience with 1.030 surgical patients. *Anesthesiology* 1998;88(3):688-95.
- Broekedema AA, Gielen MJ, Hennis PJ. Postoperative analgesia with continuous epidural sufentanil and bupivacaine: a prospective study in 614 patients. *Anesth Analg* 1996;82(4):754-9.
- George KA, Chisakuta AM, Gamble JA, Browne GA. Thoracic epidural infusion for postoperative pain relief following abdominal aortic surgery: bupivacaine, fentanyl or a mixture of both? *Anaesthesia* 1992;47(5):388-94.
- Webster TM, Herrell SD, Chang SS, Cookson MS, Baumgartner RG, Anderson LW, et al. Robotic assisted laparoscopic radical prostatectomy versus retropubic radical prostatectomy: a prospective assessment of postoperative pain. *J Urol* 2005;174(3):912-4.
- D'Alonzo RC, Gan TJ, Moul JW, Albala DM, Polascik TJ, Robertson CN, et al. A retrospective comparison of anesthetic management of robot-assisted laparoscopic radical prostatectomy versus radical retropubic prostatectomy. *J Clin Anesth* 2009;21(5):322-8. DOI: 10.1016/j.jclinane.2008.09.005.
- Höhwu L, Akre O, Bergenwald L, Törnblom M, Gustafsson O. Oral oxycodone hydrochloride versus epidural anaesthesia for pain control after radical retropubic prostatectomy. *Scand J Urol Nephrol* 2006;40(3):192-7.
- Poikolainen E, Hendolin H. Effects of lumbar epidural analgesia and general anaesthesia on flow velocity in the femoral vein and postoperative deep vein thrombosis. *Acta Chir Scand* 1983;149(4):361-4.
- Hendolin H, Mattila MA, Poikolainen E. The effect of lumbar epidural analgesia on the development of deep vein thrombosis of the legs after open prostatectomy. *Acta Chir Scand* 1981;147(6):425-9.
- Ballantyne JC, Carr DB, de Ferranti S, Suarez T, Lau J, Chalmers TC, et al. The comparative effects of postoperative analgesic therapies on pulmonary outcome: cumulative meta-analyses of randomized, controlled trials. *Anesth Analg* 1998;86(3):598-612.
- Beattie WS, Badner NH, Choi P. Epidural analgesia reduces postoperative myocardial infarction: a meta-analysis. *Anesth Analg* 2001;93(4):853-8.
- Rodgers A, Walker N, Schug S, McKee A, Kehlet H, van Zundert A, et al. Reduction of postoperative mortality and morbidity with epidural or spinal anaesthesia: results from overview of randomised trials. *BMJ* 2000;321(7275):1-12.
- Ene KW, Nordberg G, Sjöström B, Bergh I. Prediction of postoperative pain after radical prostatectomy. *BMC Nurs* 2008;7:14. DOI: 10.1186/1472-6955-7-14.
- Ahmed A, Baig T. Incidence of lower limb motor weakness in patients receiving postoperative epidural analgesia and factors associated with it: an observational study.

- Saudi J *Anaesth* 2016;10(2):149-53. DOI: 10.4103/1658-354X.168806.
28. Tornero Tornero JC, Gómez Gómez M, Fabregat Cid G, et al. Aliaga Font L, Roqués Escolar V, Escamilla Cañete B, et al. *Rev Esp Anesthesiol Reanim* 2008;55(9):552-62.
 29. Souza MP, Magalhães E, Cascudo Ede F, Jogaib MA, Silva MC. Accidental catheterization of epidural venous plexus: tomographic analysis. *Rev Bras Anesthesiol* 2016;66(2):208-11. DOI: 10.1016/j.bjan.2013.03.022.
 30. Niraj G, Kelkar A, Jeyapalan I. Comparison of analgesic efficacy of subcostal transversus abdominis plane blocks with epidural analgesia following upper abdominal surgery. *Anaesthesia* 2011;66(6):465-71. DOI: 10.1111/j.1365-2044.2011.06700.x.
 31. Sharma P, Chand T, Saxena A, Bansal R, Mittal A, Shrivastava U. Evaluation of postoperative analgesic efficacy of transversus abdominis plane block after abdominal surgery: a comparative study. *J Nat Sci Biol Med* 2013;4(1):177-80. DOI: 10.4103/0976-9668.107286.
 32. Cameron CM, Scott DA, McDonald WM, Davies MJ. A review of neuraxial epidural mortality: experience of more than 8000 cases at a single teaching hospital. *Anesthesiology* 2007;106(5):997-1002.
 33. Brodner G, Mertes N, Van Aken H, Pogatzki E, Buerkle H, Marcus MA, et al. Epidural analgesia with local anesthetics after abdominal surgery: early motor recovery with 0.2 % ropivacaine than 0.175 % bupivacaine. *Anesth Analg* 1999;88(1):128-33. DOI: 10.1213/00000539-199901000-00024.
 34. Kehlet H, Wilmore DW. Evidence-based surgical care and the evolution of fast-track surgery. *Ann Surg* 2008;248(2):189-98. DOI: 10.1097/SLA.0b013e31817f2c1a.
 35. Schwenk W, Row W, Haase O. Fast-track kolon chirurgie. *Chir urg* 2004;75:508-14.
 36. Wind J, Polle SW, Fung Kon Jin PH, Dejong CH, von Meyenfeldt MF, Ubbink DT, et al; Laparoscopy and/or Fast Track Multimodal Management Versus Standard Care (LAFA) Study Group; Enhanced Recovery after Surgery (ERAS) Group. Systematic review of enhanced recovery programmes in colonic surgery. *Br J Surg*. 2006;93(7):800-9.
 37. Podore PC, Throop EB. Infrarenal aortic surgery with a 3-day hospital stay: A report on success with a clinical pathway. *J Vasc Surg* 1999;29(5):787-92.
 38. Tovar EA, Roethe RA, Weissig MD, Lloyd RE, Patel GR. One-day admission for lung lobectomy: an incidental result of a clinical pathway. *Ann Thorac Surg* 1998;65(3):803-6.
 39. Arumainayagam N, McGrath J, Jefferson KP, Gillat DA. Introduction of an enhanced recovery protocol for radical cystectomy. *BJU Int* 2008;101(6):698-701. DOI: 10.1111/j.1464-410X.2007.07319.x.
 40. Kehlet H. Modification of responses to surgery by neural blockade clinical implications. In: Cousins HJ, Bridenbough PO, eds. *Neural blockade in clinical anesthesia and management of pain*. Philadelphia: JB Lippincot; 1998. p. 129-75.
 41. Borzellino G, Francis NK, Chapuis O, Krastinova E, Dyevre V, Genna M, et al. Role of Epidural Analgesia within an ERAS Program after Laparoscopic Colorectal Surgery: A Review and Meta-Analysis of Randomised Controlled Studies. *Surg Res Pract* 2016;2016:7543684. DOI: 10.1155/2016/7543684.
 42. Levy BF, Scott MJ, Fawcett W, Fry C, Rockall TA. Randomized clinical trial of epidural, spinal or patient-controlled analgesia for patients undergoing laparoscopic colorectal surgery. *Br J Surg* 2011;98(8):1068-78. DOI: 10.1002/bjs.7545.
 43. McDonnell J, O'Donnell B, Curley G, Hefferman A, Power C, Laffey JG. The analgesic efficacy of transverses abdominis plane block after abdominal surgery: a prospective randomized controlled trial. *Anesth Analg* 2007;104(1):193-7.
 44. De Oliveira GS, Castro-Alves L, Nader A, Kendall MC, McCarthy RJ. Transversus abdominis plane block to ameliorate postoperative pain outcomes after laparoscopic surgery: a meta-analysis of randomized controlled trials. *Anesth Analg* 2014;118(2):454-63. DOI: 10.1213/ANE.0000000000000066.