

Risk for decreased cardiac tissue perfusion and activity intolerance: Association study*

Riesgo de disminución de la perfusión del tejido cardíaco e intolerancia a la actividad: estudio asociativo

Risco de diminuição da perfusão do tecido cardíaco e intolerância à atividade: estudo de associação

*The data for this study were obtained from the dissertation "Clinical validation of the nursing diagnosis of decreased cardiac tissue perfusion in patients with coronary artery obstruction," presented by Julia Leme Gonçalves at Universidade de Campinas, Brazil.

How cite: Pendiente

1 Suellen Cristina Dias Emidio

Universidade Federal de Tocantins (Palmas, TO, Brazil).
ORCID: <https://orcid.org/0000-0003-2790-0271>
E-mail: suellen.emidio@outlook.com

Contribution: Data analysis and interpretation, writing of the article, critical review of the article.

2 Laís Pereira Giovanini

Universidade de Campinas (Campinas, SP, Brazil).
ORCID: <https://orcid.org/0000-0001-9455-3607>
E-mail: lagiovanini@gmail.com

Contribution: Research conception and design, data collection, data analysis and interpretation.

3 Paula Rocco Gomes Lima

Universidade de Campinas (Campinas, SP, Brazil).
ORCID: <https://orcid.org/0000-0001-9601-0999>
E-mail: paularglima@gmail.com

Contribution: Research conception and design, data analysis and interpretation, writing of the article, critical review of the article.

4 Julia Leme Gonçalves

Universidade de Campinas (Campinas, SP, Brazil).
ORCID: <https://orcid.org/0000-0002-1570-3898>
E-mail: julia_lemeg@yahoo.com.br

Contribution: Research conception and design, data collection, data analysis and interpretation.

5 Ana Railka de Souza Oliveira

Universidade de Campinas (Campinas, SP, Brazil).
ORCID: <https://orcid.org/0000-0002-7075-7987>
E-mail: ana.railka@gmail.com

Contribution: Research conception and design, data analysis and interpretation, writing of the article, critical review of the article.

DOI: <http://doi.org/10.15446/avenferm.v39n3.94571>

Recibido: 00/00/2020 Aceptado: 00/00/2021

ISSN (impreso): 0121-4500
ISSN (en línea): 2346-0261



Abstract

Objective: To identify the nursing diagnoses of risk for decreased cardiac tissue perfusion and risk for activity intolerance and establish the association between the components of both diagnoses and the signs and symptoms of acute coronary syndromes.

Materials and methods: Observational and associative study with 75 patients diagnosed with acute coronary syndrome in a Brazilian public hospital. Inferential statistics and multiple Poisson regression models were applied.

Results: 84% of the patients showed risk for activity intolerance and 80% reported risk for decreased cardiac tissue perfusion. Risk factors were present in more than 50% of the sample. The diagnoses showed a high frequency in patients with acute coronary syndrome. Pharmacological agents, presence of comorbidities, and family and personal history were associated with the diagnoses. Absence of evidence of decreased cardiac tissue perfusion as dyspnea, radiation to shoulder and jaw, and pain time less than ten hours acted as protective factors.

Conclusions: There is a high cardiovascular vulnerability of patients with acute coronary syndrome to the proposed nursing diagnoses. Therefore, we recommend further studies to determine the predictive power of the assessed risk diagnoses for those focusing on this health problem.

Descriptors: Acute Coronary Syndrome; Signs and Symptoms; Risk Factors; Nursing Diagnosis; Nursing Process (source: DeCS, BIREME).

Resumen

Objetivo: identificar los diagnósticos de enfermería sobre el riesgo de disminución de la perfusión tisular cardíaca y el riesgo de intolerancia a la actividad y establecer la asociación existente entre los componentes de ambos diagnósticos y los signos y síntomas de los síndromes coronarios agudos.

Materiales y métodos: estudio observacional y asociativo realizado con 75 pacientes diagnosticados con síndrome coronario agudo en un hospital público de Brasil. Se aplicaron estadísticas inferenciales y modelos de regresión de Poisson múltiples.

Resultados: 84 % de los pacientes presentó riesgo de intolerancia a la actividad y 80 % riesgo de disminución de la perfusión del tejido cardíaco. Los factores de riesgo estuvieron presentes en más del 50 % de la muestra. Los diagnósticos mostraron una alta frecuencia en pacientes con síndrome coronario agudo. Factores como el uso de agentes farmacológicos, la presencia de comorbilidades y antecedentes familiares y personales fueron asociados con ambos diagnósticos. La ausencia de evidencia sobre disminución de la perfusión del tejido cardíaco como disnea, radiación en el hombro y la mandíbula o la presencia de dolor por un período inferior a diez horas actuaron como factores protectores.

Conclusiones: existe una alta vulnerabilidad cardiovascular de los pacientes con síndrome coronario agudo a los diagnósticos de enfermería propuestos, por lo que se sugiere realizar más estudios con el fin de determinar el poder predictivo de los diagnósticos de riesgo evaluados para aquellos encargados de estudiar este problema.

Descriptoros: Síndrome Coronario Agudo; Signos y Síntomas; Factores de Riesgo; Diagnóstico de Enfermería; Proceso de Enfermería (fuente: DeCS, BIREME).

Resumo

Objetivo: identificar os diagnósticos de enfermagem risco de perfusão tecidual cardíaca diminuída e risco de intolerância à atividade e estabelecer a associação entre os componentes desses diagnósticos e os sinais e sintomas das síndromes coronarianas agudas.

Materiais e métodos: estudo observacional e associativo realizado com 75 pacientes com diagnóstico de síndrome coronariana aguda em um hospital público brasileiro. Estatísticas inferenciais e modelos de regressão de Poisson múltiplos foram aplicados.

Resultados: 84 % apresentaram risco para intolerância à atividade e 80 % com risco para diminuição da perfusão do tecido cardíaco. Os fatores de risco estiveram presentes em mais de 50 % da amostra. Os diagnósticos apresentaram alta frequência em pacientes com síndrome coronariana aguda. Agentes farmacológicos, presença de comorbidades, história familiar e pessoal estiveram associados aos diagnósticos. Ausência de evidências de diminuição da perfusão do tecido cardíaco como dispneia, radiação para ombro e mandíbula e tempo de dor menor que dez horas atuaram como fatores de proteção.

Conclusões: há alta vulnerabilidade cardiovascular dos pacientes com síndrome coronariana aguda aos diagnósticos de enfermagem propostos, assim, recomendamos novos estudos para determinar o poder preditivo dos diagnósticos de risco avaliados para aqueles com enfoque no problema.

Descritores: Síndrome Coronariana Aguda; Sinais e Sintomas; Fatores de Risco; Diagnóstico de Enfermagem; Processo de Enfermagem (fonte: DeCS, BIREME).

Introduction

Continuous comprehensive care with patients requires a common language between nursing healthcare. In this context, the use of NANDA International Inc. (NANDA-I) taxonomy stands out internationally. The Nursing Diagnoses (NDS) of NANDA-I are based on the diagnosis of human responses through clinical reasoning in which needs are identified and treatment and prevention strategies proposed. These answers can be based on potential or real problems (1).

As for the care of patients with susceptibility to decreased cardiac tissue perfusion (DCTP), we identify the ND Risk for Decreased Cardiac Tissue Perfusion (00200) in the taxonomy. To assist in the development of NANDA-I Taxonomy II, two recent surveys have proposed and validated (2, 3) the ND of DCTP, presenting acute coronary syndromes (ACS) as the most common associated condition. These works also discuss the relationship of DCTP with other diagnoses concerning Domain 4 – Activity/Rest, Class 4 – Cardiovascular/Pulmonary responses, such as decreased cardiac output and activity intolerance.

Rethinking care with a focus on NDS is still a major challenge for nurses, where we highlight two points. First, because the focus on the disease is the prevailing approach. Second, because we find it difficult to think both of care as a continuum and on the interrelationships among NDS, especially when we think of risky and real NDS.

Several studies focusing on nursing diagnosis have been developed (4-6), especially when we think about real NDS. Thus, the main objectives of this research were to identify the nursing diagnoses of risk for decreased cardiac tissue perfusion (RDCTP) [code: 00200] and risk for activity intolerance (RAI) [code: 00094] and to establish the relationship between the components of these diagnoses and the signs and symptoms of decreased cardiac tissue perfusion. As a specific objective, we seek to characterize the sociodemographic and clinical profile of the studied population. NDS were chosen for this research since they are frequent in the clinical practice of nurses and because they remain little studied subjects that could contribute to better subsidize health care planning.

Materials and methods

Design

This is an observational, associative, quantitative, and cross-sectional study derived from a clinical construction validation study based on accuracy test models (3). The research scenario comprised the emergency, hospitalization, intensive care, and hemodynamic units in Brazil.

Participants and sample

The inclusion criteria for the study established the selection of patients aged 18 or older with a medical diagnosis of ACS and who were in the acute phase of the disease. Patients who did not present the clinical condition (signs and symptoms) at the time of admission or who had undergone reperfusion therapy before admission to the service were excluded from the study. Eligible patients were informed before discharge and enrolled in the study.

As the present study is derived from primary research to verify the accuracy of the nursing diagnosis regarding decreased cardiac tissue perfusion in patients with ACS (3), sample size calculation considered the clinical construct validation methodology for the latent class analysis (LCA) (7).

The LCA is used to calculate accuracy measures (sensitivity and specificity) of clinical indicators when there is no perfect reference standard. For this reason, the sample size may vary from 5 to 30 individuals for each defining characteristic (7). Therefore, considering that the ND of DCTP has 15 defining characteristics, a total of 5 patients were considered per each defining characteristic, totaling 75 patients (3).

Data collection

Data collection with patients and medical records occurred from August 2018 to February 2019. This process was carried out by a nurse and a nursing senior student.

In order to standardize data collection, conceptual and operational definitions for each component of the NDS under study were developed as a protocol. This construction was submitted to the appreciation of three judges through a focus group. These were nurses, teachers, and researchers in the area of Standardized Language System or cardiology and

with experience in caring for patients with ACS. The protocol was adjusted according to the consensus of the discussion group and was used by the research team to reduce the evaluation and instrument bias. In addition, to decrease the evaluation bias related to measurement accuracy, the research team carried out a theoretical-practical training of eight hours to apply the instrument built for the collection of data and to carry out the physical examination in a standardized way. After a pre-test with five patients, who were not included in the final version, it was possible to verify the adequacy of the data recording instrument to start data collection.

Interviews included questions about patients' baseline sociodemographic characteristics and social background: sex, age, marital status, education, employment status, family income, and number of inhabitants per household. Data regarding the presence of risk factors for ACS (arterial hypertension, diabetes mellitus, smoking, dyslipidemia, and obesity) and others risk factors were also collected, following the protocol previously defined. In this case, for example, an open-ended question about the factors that could lead to ACS was formulated, so the examiner could check which risk factor(s) were known to patients and thus minimize bias in the instrument. Data regarding the associated conditions and whether patients fit into risk populations for the NDS under study were investigated in medical records.

To complement data collection, we sought medical records information on clinical indicators of DCTP on admission, as recommended by Santos *et al.* (2) and validated by Gonçalves *et al.* (3), namely: angina, pain score higher than seven, electrocardiographic changes, arrhythmia, altered cardiac enzymes, dyspnea, irradiation of pain to shoulder and jaw, pain duration for less than ten hours, nausea, altered blood pressure, dizziness, vomiting, pulmonary rales, altered heart rate, and third heart sound.

The diagnostic inference was carried out independently by a nursing student, a nurse specialized in cardiology, who is also enrolled in a master's degree, and a nurse doctor and researcher in the study area, all of them with broad knowledge of nursing taxonomies. This helped to limit the spectrum bias, referring the researcher to determine which patient has a diagnosis through their perception of patient impairment. Each evaluator received the clinical information of all patients in an Excel software spreadsheet. Afterwards, the agreement reached by the three evaluators was used as the gold standard for the presence of NDS, as in the study by Oliveira *et al.* (8).

Data analysis

All statistical analyses were performed in SAS, version 9.4. Descriptive analysis was presented using means, standard deviation (SD), median, and range for continuous variables. Categorical variables were expressed as numbers and percentages. The Shapiro-Wilk test was used to verify the normality of the data. Pearson's Chi-square test or Fisher's exact test was used for associative measures of categorical data.

Multiple Poisson regression models were constructed with robust variance, considering the two NDS studied as dependent variables. In these models, prevalence ratios and their confidence intervals and p-values were presented (9). A significance level of 5% was adopted for all tests.

Ethical considerations

This study was performed following the Helsinki Declaration. Ethical approvals were obtained from the Institutional Review Boards at University of Campinas in 2018, under number 2.641.527. Each participant signed informed consent forms before enrolling in this research.

Results

Patients' sociodemographic characteristics are shown in Table 1.

Table 1. Patients' baseline demographic characteristics

Variables	No. (%)
Gender	
Male	54 (72)
Female	21 (28)
Marital status	
Married	49 (65.3)
Unmarried	26 (34.7)
Education	
Illiterate	3 (4.2)
Incomplete primary school	39 (54.2)
Complete primary school	12 (16.7)
Incomplete secondary school	3 (4.2)
Complete secondary school	9 (12.5)
Incomplete higher education	5 (6.9)
Employment status	
Unemployed	41 (54.7)
Employed	34 (45.3)

Source: authors, based on research data.

The study included 75 patients with ACS, with an average age of 60.2 years (standard deviation - SD = 10.4), in age ranges from 36 to 86 years, and who in most cases were married (65.3%). There was a prevalence of men among the sample (72%). About 54.2% of patients did not complete elementary school, the average monthly family income was 2.7 minimum wages (SD = 1.5), and the number of inhabitants/household was in mean 1.9 (SD = 1.7), with a range from 1 to 9. Besides, 30.7% of the sample was aged 65 years or over, and 84% had a history of cardiovascular disease.

As for ACS risk factors, we highlight that 72% of participants reported living with arterial hypertension, 54.7% were active or passive smokers, 45.3% had diabetes mellitus, 40% dyslipidemia, and 12% were obese. This way, all the indicators of DCTP were evidenced in this sample. The NDS and their elements are shown in Table 2.

Related to RAI diagnosis, all the risk factors of this condition were present in more than 50% of the individuals in the sample, except for the immobility risk factor, which was not identified in any of the participants. As for the risk of DCTP, 96% of the participants had insufficient knowledge about modifiable risk factors (e.g., smoking, sedentary lifestyle, obesity) and 94.7% had artery spasms associated with this coronary ND.

Table 3 shows the significant diagnostic inferences between risk factors, associated conditions, population at risk, and the presence or absence of the corresponding nursing diagnosis. Table 4, in turn, presents the multiple Poisson regression models.

It was observed that patients admitted at risk of activity intolerance and dyspnea were 1.32 times more likely to develop a diagnosis of activity intolerance. For the risk of decreased cardiac perfusion, those who did not have dyspnea and pain duration for less than ten hours were more likely to develop such diagnosis. Finally, in those who did not report pain irradiation the probability of having this diagnosis decreased by 47%, compared to those who had this symptom.

Table 2. Frequency of nursing diagnosis of risk for activity intolerance and risk for decreased cardiac tissue perfusion

Variables	No.	%
Risk for activity intolerance	63	84
Risk factors		
Inexperience with activity	70	93.3
Sedentary lifestyle	67	89.3
Physical deconditioning	65	86.7
Imbalance between oxygen supply/demand	48	64.5
At risk population		
History of previous activity intolerance	56	75.7
Associated condition		
Circulatory problem	51	68
Respiratory problem	16	21.3
Risk for decreased cardiac tissue perfusion	60	80
Risk factors		
Insufficient knowledge of modifiable factors	72	96
Substance misuse	10	13.5
At risk population		
Family history of cardiovascular disease	40	53
Associated condition		
Coronary artery spasm	71	94.7
Pharmaceutical agents	63	84
Diabetes mellitus	34	45.3
Hyperlipidemia	29	38.7
Cardiovascular surgery	14	18.7
Hypertension	14	18.7
Hypoxemia	9	12
Hypovolemia	7	9.3
Cardiac tamponade	1	1.3

Source: authors, based on research data.

Table 3. Analysis of associations of nursing diagnoses of risk for activity intolerance and risk for decreased cardiac tissue perfusion with risk factors, associated conditions, and at-risk population

Variables	Risk for activity intolerance						p-value
	Present		Absent				
	n	%	n	%			
Physical deconditioning (RF)	Yes	58	89.2	7	10.8	.0074*	
	No	5	50.0	5	50.0		
Inexperience with activity (RF)	Yes	63	90.0	7	10.0	.0001*	
	No	0	.0	5	100.0		
History of previous activity intolerance (RP)	Yes	56	100.0	0	.0	.0001*	
	No	7	38.9	11	61.1		
Circulatory problem (AC)	Yes	51	100.0	0	.0	.0001*	
	No	12	50.0	12	50.0		

Variables	Risk for decreased cardiac tissue perfusion						p-value
	Present		Absent				
	n	%	n	%			
Family history of cardiovascular disease (RP)	Yes	36	90.0	4	10.0	.0206**	
	No	24	68.6	11	31.4		
Pharmaceutical agent (AC)	Yes	54	85.7	9	14.3	.0111*	
	No	6	50.0	6	50.0		
Diabetes mellitus (AC)	Yes	34	100.0	0	.0	.0186*	
	No	26	63.4	15	36.6		

*Fisher test. **Chi-square test.

RF = Risk factor; AC = Associated conditions; RP = At-risk population.

Source: authors, based on research data.

Table 4. Multiple Poisson regression models

Dependent variable	Independent variable	PR*	CI 95%		p-value
			LL	UL	
Risk for activity intolerance	Age (≥ 60)	1.05	.83	1.33	.6713
	Sex (Male)**	.98	.79	1.22	.8712
	Dyspnea (No)	1.32	1.01	1.72	.0429
	Radiation to the shoulder and jaw (No)	1.14	.90	1.43	.2786
	Nausea (No)	.83	.58	1.20	.3286
	Vomiting (No)	1.24	.81	1.88	.3238
	Dizziness (No)	.76	.57	1.02	.0679
	Altered blood pressure (No)	.95	.77	1.18	.6354
	Pulmonary rales (No)	.97	.76	1.25	.8279
	Pain score greater than 7 (No)	1.07	.81	1.41	.6318
	Pain time less than 10 hours (No)	.98	.80	1.19	.8114
	Altered cardiac enzymes (No)	1.06	.80	1.41	.6840
	Risk for decreased cardiac tissue perfusion	Age (≥ 60)	.92	.71	1.18
Sex (Male)**		1.08	.83	1.42	.5600
Dyspnea (No)		1.54	1.15	2.06	.0042
Radiation to the shoulder and jaw (No)		.53	.36	.77	.0009
Nausea (No)		.84	.66	1.07	.1575
Vomiting (No)		.86	.61	1.20	.3628
Dizziness (No)		1.12	.75	1.65	.5836
Altered blood pressure (No)		1.17	.85	1.62	.3287
Pulmonary rales (No)		.88	.67	1.16	.3787
Pain score greater than 7 (No)		.84	.61	1.17	.3034
Pain time less than 10 hours (No)		2.28	1.42	3.64	.0006
Altered cardiac enzymes (No)		.65	.42	1.03	.0673

* PR = Prevalence ratio, the probability of presenting the result "Yes" was estimated. ** In relation to the set of variables, the evaluation of the multiple Poisson regression models only considered the male sex as the reference for the variable, since there is a higher prevalence of ACS among the participating male population.

CI = Confidence interval; LL = Lower limit; UL = Upper limit.

Bold indicates the significance at $p < 0.05$.

Source: authors, based on research data.

Discussion

Population aging and the increase in chronic diseases are important factors in the growing prevalence of ACS, as reflected in the sociodemographic characteristics collected by this study (10-12). Corroborating the findings of other studies, participating patients showed hypertension, diabetes mellitus, dyslipidemia, and other cardiovascular diseases. We also found that smoking and obesity are associated risk factors (13-16). Due to the absence of studies that analyzed the elements of the two NDS studied in the population chosen for this study, we made comparisons with studies that performed an analysis in patients with a phenomenon often installed, which demands the importance of early assessment.

Research with stable heart disease patients identified 13 NDS, where lifestyle (00168) and activity intolerance (00092) stand out. However, RDCTP (00200) and RAI (00094) presented less agreement between specialist nurses (15), which shows the difference between NDS of the acute, subacute and chronic faces. The evidence of activity intolerance (00092) shows that the presence of dyspnea on exertion, fatigue, the imbalance between oxygen supply and demand, stress discomfort, and the presence of angina is a common feature (15).

The insufficient knowledge of modifiable factors was identified in a more significant percentage of the patients in this sample who were diagnosed with RDCTP. The correct adherence to treatment and the rapid conduction in emergency cases are affected by the lack of knowledge about ACS risk factors, as well as by failure in identifying the signs of the ischemic event in heart diseases (17, 18).

In this context, knowledge about risk factors needs to be addressed in primary and secondary care services. The prioritization of the approach to drug therapy in consultations results in a lack of knowledge about heart disease as a whole by the population, a fact observed in a study with patients with ACS (18).

In nursing appointments, RDTP increases when there is a family history of cardiovascular disease or diabetes mellitus, accompanied by the use of drugs that act on the heart pump. Concomitant with these factors, early non-adherence to treatment and lack of therapeutic continuity increases the chances of the patient developing again DCTP.

In this study, the defining characteristics of DCTP were observed in patients at the time of admission to the emergency department, showing a relationship with the determination of risk diagnoses. In other words, the accurate and precise identification of clinical parameters is essential for making diagnostic and therapeutic decisions with potential impacts on the evolution of the patient.

As previously stated (3), patients with ACS showed angina (100%), pain score higher than seven (81.3%), electrocardiographic changes (82.4%), arrhythmia (78%), altered cardiac enzymes (76.4%), dyspnea (61.6%), radiation to shoulder and jaw (55.4%), pain for less than ten hours (53.3%), nausea (45.3%), altered blood pressure (30.7%), dizziness (29.3%), vomiting (26.7%), pulmonary rales (18.9%), altered heart rate (14.7%), and third heart sound (4%).

Another point to be explored is the relationship between the presence of ACS and increased risk of functional decline, which leads to an increased chance of hospital readmissions and death in elderly patients, as well as impacts on their quality of life (19, 20). Thus, it is evident the gap that needs to be addressed during early stages, through intervention studies aimed at promotion, disease prevention, and rehabilitation of patients at risk, thus favoring the non-development of problem-focused NDS (15).

In the context of hospital discharge of patients with RDCTP and RAI diagnoses, it is necessary to continue with rehabilitation mediated by a multidisciplinary team in patients' home environment. Therefore, the complete evaluation, with the introduction of elements of the measure of functionality, should be prioritized, especially when considering individuals with multiple morbidities and the elderly (20). Based on the data collected, the importance of investigating the patient's previous and current functional status during hospitalization is a remarkable finding that will contribute to reducing the impacts on rehabilitation.

Linked to the elements of physical deconditioning and inexperience with activity, the ability to walk is a crucial point to be evaluated.

A possible explanation for this fact would be the relationship between the loss of microvascular perfusion after ACS, which is responsible for the supply of oxygen and nutrients to the tissues, and the continuity of ischemia and the development of adverse clinical results (21).

In another study, it was observed that the limitations in performing basic activities of daily living started eight years before the acute event (22). That is the period in which interventions for RDCTP and RAI could have started, considering the presence of risk factors, thus corroborating the findings of our study regarding the association of NDS and the risk of activity intolerance and the history of previous activity intolerance element.

However, faced with the evidence of low adherence to health promotion and disease prevention activities, the possibility of a decline that may affect the area of cardiac rehabilitation (CR) becomes a concern. At home and in the primary care setting, CR could contribute to improving and maintaining physical activity habits and increasing cardiorespiratory fitness and survival, thus reducing the risk of mortality (23).

It should be noted that in the current study sample all risk factors for RAI were identified, although only in association with inexperience with activity, and imbalance between oxygen supply/demand. On this regard, a study whose objective was to verify the prevalence of ND sedentary lifestyle (SL) and the defining characteristic of physical deconditioning in patients with ACS reported a frequency of 56.1 and 46.8%, respectively (24).

In the case of appearance of these two elements, there is a higher chance of systemic, muscular, respiratory, cardiovascular, and articular changes, and, consequently, activity intolerance, which might cause a new episode of DCTP (15). As for RDCTP, all the constituent elements of NDS were identified in the sample studied. According to the research study by Moreira *et al.* (25), in which 1,542 adult patients were monitored for seven years after percutaneous coronary intervention (PCI), 12.1% of the sample developed DCTP, listing advanced age, multivessel disease, and complications of the intra-procedure injury as predictor of such a condition. These outcomes reinforce the need for continuous monitoring of the population at risk.

The care of patients with DCTP should be focused on health promotion and risk prevention in light of their basic psychosocial and psychobiological human needs. The results found in this study reinforce this claim. The importance of early identification of risk diagnosis and risk factors was also observed through this research. Therefore, diagnosing these risk factors significantly contributes to the care and prevention of ACS complications.

We should mention as a limitation of this research that it is a cross-sectional study, leading to the observation of some characteristics only in a single moment, which could compromise the reading of the phenomenon. Besides, due to the use of a sample for convenience in a single hospital, our findings may not be generalized.

Conclusion

The nursing diagnoses of risk of activity intolerance and risk of decreased cardiac tissue perfusion were present in 84 and 80% of patients with DCTP, respectively.

When verifying the relationship between the risk factors and the NDS studied, we found an association of the use of pharmacological agents, the presence of diabetes mellitus, and family history of coronary disease with RPCD. In the case of RAI, this reports an association with lack of physical conditioning, previous history of activity intolerance, inexperience with activity, and circulatory problems. Finally, RPCP is associated with pharmacological agents and dyslipidemia.

The absence of evidence of DCTP such as dyspnea, radiation to shoulder and jaw, and arterial pressure, and pain for less than ten hours acted as protective factors for the development of the studied risk diagnoses.

To conclude, considering the high cardiovascular vulnerability of patients with ACS to nursing diagnoses through the NANDA-I domain Class 4, we recommend further studies to determine the predictive power of the risk diagnoses evaluated by ND from a problem-centered approach.

Financial support

This study did not receive any funding.

References

- (1) Herdman T; Kamitsuru S. NANDA International Nursing Diagnoses: Definitions & Classification 2018-2020. 11th ed. New York: Thieme; 2017.
- (2) Santos VB; Atallah AN; Lopes CT; Lopes JL; Bottura ABL. Defining characteristics and related factors of decreased cardiac tissue perfusion: Proposal of a new nursing diagnosis. *Int J Nurs Knowl.* 2016;27(3):175-180. <http://doi.org/10.1111/2047-3095.12095>

- (3) Gonçalves JL; Giovanini LP; Lopes JL; Cornélio ME; Lopes MVO; Oliveira-Kumakura ARS. Clinical validation of the nursing diagnosis, decreased cardiac tissue perfusion, in patients with coronary artery obstruction. *Int J Nurs Knowl.* 2020;31(3):205-209. <http://doi.org/10.1111/2047-3095.12269>
- (4) Souza TSD; Campos JDC; Villela SF; Lins GAI; Melo MC. Nursing diagnoses and outcomes for children with nutritional anomalies: A descriptive study. *Rev Bras Enferm.* 2020;73(3):e20180809. <http://doi.org/10.1590/0034-7167-2018-0809>
- (5) Souza-Neto VLD; Costa RTDS; Santos WN; Fernandes SF; Lima DM; Silva RAR. Validation of the definitions of nursing diagnoses for individuals with Aids. *Rev Bras Enferm.* 2020;73(4):e20180915. <http://doi.org/10.1590/0034-7167-2018-0915>
- (6) Melguizo-Herrera E; Acosta-López A; Gómez-Palencia IP; Manrique-Anaya Y; Hueso-Montoro C. The design and validation of a nursing plan for elderly patients with postoperative delirium. *Int J Environ Res Public Health.* 2019;16(22):4504. <http://doi.org/10.3390/ijerph16224504>
- (7) Lopes MV; Silva VM; Araujo TL. Validation of nursing diagnosis: Challenges and alternatives. *Rev Bras Enferm.* 2013;66(5):649-655. <http://doi.org/10.1590/S0034-71672013000500002>
- (8) Oliveira AR; Rodrigues RC; Sousa VE; Costa AG; Lopes MV; Araujo TL. Clinical indicators of 'caregiver role strain' in caregivers of stroke patients. *Contemp Nurse.* 2013; 44(2):215-224. <http://doi.org/10.5172/conu.2013.44.2.215>
- (9) Zou G. A modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol.* 2004;159(7):702-706. <http://doi.org/10.1093/aje/kwh090>
- (10) Al-Rasadi K; Al-Zakwani I; Alsheikh-Ali AA; Almahmeed W; Rashed W; Ridha M *et al.* Prevalence, management, and outcomes of familial hypercholesterolemia in patients with acute coronary syndromes in the Arabian Gulf. *J Clin Lipidol.* 2018;12(3):685-92.e2. <http://doi.org/10.1016/j.jacl.2018.02.003>
- (11) Abreu M; Vensentini N; Mariani J; Gagliardi J; Doval H; Tajer C. Acute coronary syndromes in public and non-public institutions of Argentina. *Medicina (B Aires).* 2019;79(6):461-467.
- (12) Lindholm D; James SK; Gabrysch K; Storey RF; Himmelmann A; Cannon CP *et al.* Association of multiple biomarkers with risk of all-cause and cause-specific mortality after acute coronary syndromes: A secondary analysis of the PLATO Biomarker Study. *JAMA Cardiol.* 2018;3(12):1160-1166. <http://doi.org/10.1001/jamacardio.2018.3811>
- (13) Cavalcante AM; Brunori EH; Lopes CT; Silva AB; Herdman TH. Nursing diagnoses and interventions for a child after cardiac surgery in an intensive care unit. *Rev Bras Enferm.* 2015;68(1):155-160. <http://doi.org/10.1590/0034-7167.20156801211>
- (14) Bugiardini R; Badimon L. The International Survey of Acute Coronary Syndromes in Transitional Countries (ISACS-TC): 2010-2015. *Int J Cardiol.* 2016;217(Suppl:S1-6). <http://doi.org/10.1016/j.ijcard.2016.06.219>
- (15) Cardoso PC; Caballero LG; Ruschel KB; Moraes MAP; Silva ERR. Profile of the nursing diagnoses in stable heart disease patients. *Invest Educ Enferm.* 2019;37(2):234-251. <http://doi.org/10.17533/udea.iee.v37n2e08>
- (16) Chang HJ; Lin FY; Lee SE; Andreini D; Bax J; Cademartiri F *et al.* Coronary atherosclerotic precursors of acute coronary syndromes. *J Am Coll Cardiol.* 2018;71(22):2511-2522. <http://doi.org/10.1016/j.jacc.2018.02.079>
- (17) Duber HC; McNellan CR; Wollum A; Phillips B; Allen K; Brown JC *et al.* Public knowledge of cardiovascular disease and response to acute cardiac events in three cities in China and India. *Heart.* 2018;104(1):67-72. <http://doi.org/10.1136/heartjnl-2017-311388>
- (18) Fernandes AC; McIntyre T; Coelho R; Prata J; Maciel MJ. Impact of a brief psychological intervention on lifestyle, risk factors and disease knowledge during phase I of cardiac rehabilitation after acute coronary syndrome. *Rev Port Cardiol.* 2019;38(5):361-368. <http://doi.org/10.1016/j.repc.2018.09.009>
- (19) Flint K; Kennedy K; Arnold SV; Dodson JA; Cresci S; Alexander KP. Slow gait speed and cardiac rehabilitation participation in older adults after acute myocardial infarction. *J Am Heart Assoc.* 2018;7(5):e008296. <http://doi.org/10.1161/JAHA.117.008296>

(20) Dodson JA; Hajduk AM; Murphy TE; Geda M; Krumholz HM; Tsang S *et al.* Thirty-day readmission risk model for older adults hospitalized with acute myocardial infarction. *Circ Cardiovasc Qual Outcomes.* 2019;12(5):e005320.

<http://doi.org/10.1161/CIRCOUTCOMES.118.005320>

(21) Gkontra P; El-Bouri WK; Norton KA; Santos A; Popel AS; Payne SJ *et al.* Dynamic changes in microvascular flow conductivity and perfusion after myocardial infarction shown by image-based modeling. *J Am Heart Assoc.* 2019;8(7):e011058.

<http://doi.org/10.1161/JAHA.118.011058>

(22) Kucharska-Newton A; Griswold M; Yao ZH; Foraker R; Rose K; Rosamond W *et al.* Cardiovascular disease and patterns of change in functional status over 15 years: Findings from the Atherosclerosis Risk in Communities (ARIC) study. *J Am Heart Assoc.* 2017;6(3).

<http://doi.org/doi.org/10.1161/JAHA.116.004144>

(23) Chernomordik F; Sabbag A; Tzur B; Kopel E; Goldkorn R; Matetzky S *et al.* Cardiac rehabilitation following an acute coronary syndrome: Trends in referral, predictors and mortality outcome in a multicenter national registry between years 2006-2013: Report from the Working Group on Cardiac Rehabilitation, the Israeli Heart Society. *Eur J Prev Cardiol.* 2017;24(2):123-132.

<http://doi.org/10.1177/2047487316680692>

(24) Costa C; Costa-Linch GF; Souza EN. Nursing diagnosis based on signs and symptoms of patients with heart disease. *Int J Nurs Knowl.* 2016;27(4):210-214.

<http://doi.org/10.1111/2047-3095.12132>

(25) Moreira FM; Nascimento TCD; Murakami BM; Bergamasco EC; Lopes CT; Santos ER. Clinical predictors of risk for decreased cardiac tissue perfusion after percutaneous coronary intervention: A retrospective cohort study. *Int J Nurs Knowl.* 2019;30(3):162-167.

<http://doi.org/10.1111/2047-3095.12212>