

Association between self-assessment and clinical oral health status in adults, Paraguay.

Asociación entre la autoevaluación y el estado clínico de salud oral en adultos, Paraguay.

Clarisse Díaz-Reissner.¹
Juan Roldán-Merino.²⁻⁵
Irma Casas-García.⁶⁻⁸

Affiliations:

¹Facultad de Odontología. Universidad Nacional de Asunción. Asunción, Paraguay.

²Sant Joan de Déu-Fundació Privada. Barcelona, Spain.

³Universitat Autònoma de Barcelona. Barcelona, Spain.

⁴Rovira i Virgili University of Tarragona. Tarragona, Spain.

⁵Research Group GEIMAC, Barcelona, Spain.

⁶Universitat Autònoma de Barcelona. Departamento de Pediatría, Obstetricia y Ginecología y Medicina Preventiva. Barcelona, Spain.

⁷Preventive Medicine Service. Hospital Germans Trias i Pujol. Barcelona, Spain.

⁸Research Group Innovation in Respiratory Infections and Tuberculosis Diagnosis (Group Consolidat 2017 SGR 494).

Corresponding author: Clarisse Díaz-Reissner. Facultad de Odontología UNA. Avenida España c/ Brasil N° 430. Asunción, Paraguay. **E-mail:** cdiazr@odo.una.py

Receipt : 04/22/2020 **Revised:** 08/12/2020
Acceptance: 12/28/2020

Cite as: Díaz-Reissner C, Roldán-Merino J & Casas-García I.

Association between self-assessment and clinical oral health status in adults, Paraguay.

J Oral Res 2020; 9(6):483-489.

Doi:[10.17126/joralres.2020.094](https://doi.org/10.17126/joralres.2020.094)

Abstract: Introduction: Oral health can be defined as the absence of pathologies and disorders that affect the stomatognathic system. **Objectives:** This study aimed to determine the characteristics of self-assessment oral health status, in the clinical experience of dental caries, periodontal status, periodontal fixation loss and to investigate the association between self-reported and clinical oral health status among Paraguayans adults during early 2017. **Material and Methods:** The design is cross-sectional. Two dentists carried out the oral examinations following the recommendations of the World Health Organization (WHO). **Results:** There were 333 adult participants with a mean age of 35 ± 13 years. Most (77.2%) of the participants were female. Missing teeth (5.32 ± 6), filled teeth (3.56 ± 4), and decayed teeth (2.55 ± 3) were also detected. Almost half (48.0%) of participants had dental calculus, while 5.8% had a periodontal pocket and 48.6% periodontal fixation loss. The self-perception of oral health was poor in 12.3% of participants, fair in 29.8%, normal in 31.8%, good in 16.2%, and excellent in 9.9%. Oral health self-assessment was positive in 58.0% and negative in 42.0%. A statistically significant difference was found when comparing the DMFT index according to self-perception of oral health, the score being higher in those who had negative self-perception; obtaining similar results in the decayed component. **Conclusion:** Negative oral health self-perception was associated with a high DMFT index, of this, the decayed component was the only one that presented statistically significant differences.

Keywords: Public health; oral health; adults; Paraguay; diagnosis, oral; self-assessment.

Resumen: Introducción: La salud bucodental puede ser definida como la ausencia de patologías y trastornos que afectan el sistema estomatognático. **Objetivos:** Este estudio tuvo como objetivo determinar las características de la autoevaluación del estado de salud bucal, en la experiencia clínica de la caries dental, el estado periodontal, la fijación de la pérdida periodontal e investigar la asociación entre el estado de salud bucal autoinformado y clínico entre los adultos paraguayos a principios de 2017. **Material y Métodos:** El diseño es transversal. Dos odontólogos fueron calibrados para el examen bucal, siguiendo

las recomendaciones de la Organización Mundial de la Salud (OMS). **Resultados:** Participaron 333 sujetos, el promedio de edad fue de 34.93 (DE=12.64) años, 77.2% eran mujeres. El 21.1% de las piezas dentarias estaban perdidas, el 12.2% obturadas y el 9.2% cariadas. El 48.0% presentó cálculo dental, 5.8% el bolsa periodontal y el 48.6% pérdida de fijación. La autopercepción de salud bucal fue para el 12.3% pobre, el 29.8% regular, el 31.8% normal, el 16.2% buena y el 9.9% excelente. Tuvieron una autoevaluación de su salud oral positiva el 58.0% y negativa el 42.0%. Se encontró

diferencia estadísticamente significativa al comparar el DMFT index según la autopercepción de salud oral, siendo mayor el puntaje en los negativos; obteniéndose resultados similares con el componente cariado. **Conclusión:** La autopercepción de salud oral negativa se asoció con un elevado DMFT index, de este, el componente cariado fue el único que presentó diferencias estadísticamente significativas.

Palabra Clave: Salud pública; salud bucal; adulto; Paraguay; diagnóstico bucal; autoevaluación.

INTRODUCTION.

In developed countries, the prevalence of dental caries has decreased in recent years. However, in developing countries, the prevalence of dental caries has increased or stagnated due to high carbohydrate intake, as well as an absence of regular programs of fluoride incorporation, and preventive and integral programs in dental health services.¹

In the Republic of Paraguay, 62.5% of the population is between 15-and 64-years-olds, with a demographic dividend,² whereas in Europe there is an aging population and a decrease in the birth rate.³ When measuring the dental caries experience in the Paraguayan population in 1974 the DMFT Index was consistently higher for females than males.⁴ When measuring the dental caries experience in the Lingua Indian Paraguayan Chaco population in 1984, the DMFT Index was 8.67 in 20–24-year-olds,^{9,10} in 25–29-year-olds, 8.67 in 30–34-year-olds, 9.64 in 35–40-year-olds, and 9.84 in those >40-year-olds.⁵ According to a national oral health inquiry carried out in 2017, the DMFT Index was 0.17 in 5- to 6-year-old children, 2.07 in 12-year-olds, and 4.14 in 15-year-olds.⁶

Periodontal illness can lead to chronic systemic illnesses, such as diabetes and cardiovascular pathologies, especially in older patients.⁷ It has also been associated with type 2 diabetes mellitus, atherosclerosis and atherosclerotic complications, kidney disease and adverse results of maintenance therapy of renal hemodialysis, premature delivery, some forms of pneumonia, Alzheimer's disease and rheumatoid arthritis.⁸

Only a few studies have addressed oral epidemiological oral health in the Republic of Paraguay. These

studies have reported a poor oral health condition in certain groups of the population, but there are no published data in adults in the Republic of Paraguay.

Here we aimed to determine the characteristics of self-assessment oral health status, in the clinical experience of dental caries, periodontal status, periodontal loss fixation and to investigate the association between self-reported and clinical oral health status among Paraguayans adults.

MATERIALS AND METHODS.

The design of the study was cross-sectional. The sample included adults attending the Ministry of Public Health in the Republic of Paraguay (MSPyBS) establishments (Asunción and Paraguari cities) at the Dentistry School of the National University of Asunción (UNA) and Community Center "Chacarita", during the first trimester of 2017, who were included in the study with sampling of consecutive cases. Participants were between 18- and 59-years-old and included both sexes, as well as naturalized Paraguayans and foreign nationals with more than 14 years of residence in the country.

For the calculation of the sample size, an infinite population was considered with 58% of patients who had a negative self-perception in Belarusian adults,⁸ with a precision of 7% and a confidence level of 95% a minimum of 191 subjects is required for the study. Considering a 10% loss of information, it is suggested to recruit 212 patients.

Following random sampling in the clinics of the university, and sampling of consecutive cases in the clinics of the ministry and the community center, the final study sample consisted of 333 participants. The

patients who met the inclusion criteria were invited to participate in the study after their care in the dental clinics, previously explaining the objectives of the study. Those who agreed to participate signed the informed consent and subsequently the survey and oral examination were carried out.

Two dentists were trained and calibrated (Cohen's kappa=0.84; $p=0.001$). The recommended gauge by the World Health Organization (WHO) for epidemiological studies was applied. The experience of caries was evaluated by counting the number of decayed teeth (DT), filled teeth (FT), and missing teeth (MT), and this data was used to calculate the DFMT Index score. The Periodontal Community Index (CPI) was calculated using the highest value in the Ramfjört teeth test.⁹ We used PCPUNC¹⁵ periodontal probes¹⁰ (HLW-Germany).

The variables considered were sociodemographic (age, sex, education), self-assessment oral health status, dental caries experience, periodontal health state and periodontal loss fixation.

To evaluate the self-assessment oral health status, patients were asked how they rated their oral health status, choosing one of the following based in a Likert-scale to 1 at 5 points: Excellent=1; Good=2; Normal=3; Fair=4; and Poor=5. Was categorized into two groups: positive (excellent, good or normal) and negative (fair or poor).⁸

Descriptive and inferential statistics were applied. The results were presented in Tables, for qualitative variables frequency and percentage were used, for quantitative variables measures of central tendency and dispersion were used. The Chi-square test was used to analyze the associations of oral health self-evaluation with sociodemographic variables. The relationships between the self-assessment of oral health and the clinical status of oral health were used using the Chi-square test and the Mann-Whitney U test.

All tests were used with a confidence level of 95%. Analyses were done using IBM SPSS program (Statistical Package social for Science) for Macintosh, version 22.0. The participants were informed about the aim of the investigation and who agreed to sign written informed consent. The participation was voluntary and the data collection was made confidentially. The autonomy of the person was respected, being free to participate or being able to withdraw from the study without this affecting their care in the institution or that of the person in their charge.

Thus, sufficient information about the study and the researcher's data were also provided for any subsequent consultation. No subject was discriminated against for any reason. Permission was requested from the corresponding authorities for its execution following the usual procedures of the country.

Table 1. Self-assessment according to demographic factors.

Demographic factors		Self-assessment of oral health				p-value
		Positive		Negative		
		n	%	n	%	
Sex	Male	39	20.2	36	25.7	0.235
	Female	154	79.8	104	74.3	
Education	None	11	5.7	7	5.0	0.156
	Primary	46	23.8	35	25.0	
	Secondary	79	40.9	71	50.7	
	Tertiary	57	29.5	27	19.3	
Ages	18 - 29	81	42.0	56	40.0	0.927
	30 - 39	38	27.1	38	27.1	
	40 - 49	16	11.4	16	11.4	
	50 - 59	30	21.4	30	21.4	
Total		193	58.0	140	42.0	

Table 2. Clinical oral health status of participants categorized by self-assessment oral health.

Clinical oral health		Self-assessment of oral health		p-value
		Positive	Negative	
Number of decayed teeth	Mean ± SD	2.0 ± 2.5	3.3 ± 3.3	0.001*
Adjusted mean to 5%		1.7	3.0	
Number of missing teeth	Mean ± SD	5.0 ± 6.1	4.8 ± 5.7	0.051
Adjusted mean to 5%		4.3	5.3	
Number of filled teeth	Mean ± SD	3.7 ± 3.8	3.3 ± 3.2	0.558
Adjusted mean to 5%		3.4	3.0	
DMFT index		10.71 ± 6.8	12.43 ± 6.5	0.016*

*: Statistically significant.

Table 3. Periodontal status of participants categorized by self-assessment oral health (n=329)

Clinical oral health		Self-assessment of oral health				p-value
		Positive		Negative		
		n	%	n	%	
Teeth with calculus	Yes	90	27.4	76	23.1	p=0.155
	No	101	30.7	62	18.8	
Teeth with bleeding on probing	Yes	34	10.3	24	7.3	0.923
	No	157	47.7	114	34.7	
Teeth with periodontal pockets	Yes	15	4.6	10	3.0	0.838
	No	176	53.5	128	38.9	
Teeth with loss fixation	Yes	44	13.4	40	12.2	0.222
	No	147	44.7	98	29.8	

RESULTS.

Three-hundred-thirty-three participants were included in the study, with a mean age of 35 ± 13 years. Most (77.2%) participants were female and 25.2% had tertiary education.

Among our participants, the self-assessment of oral health condition categories were valued as poor by 12.3%, as fair by 29.7%, as normal by 31.8%, as good by 16.2%, and as excellent by 9.9%. Oral health self-assessment was positive in 58.0% and negative in 42.0%. When comparing self-reported oral health status with sociodemographic factors, no statistically significant differences were found. (Table 1).

The general DMFT Index was 11.43 ± 6.7 . Among the participants, the mean (\pm standard deviation) number of decayed teeth (DT) was 2.55 ± 2.90 , missing teeth (MT) was 5.32 ± 5.95 , and filled teeth (FT) was 3.56 ± 3.63 . There is a statistically significant difference by sex for MT ($p=0.007$) being higher in women, but not for FT ($p=0.288$) and DT ($p=0.832$). A statistically significant difference was found when comparing the DMFT index according to the self-perception of oral health, the score being higher in the negatives; obtaining similar results for the decayed component (Table 2).

Among the participants, 1.2% were edentulous. Of the 329 patients who were evaluated, 40.4% had

a healthy periodontal state, 5.7% had bleeding with probing and needing oral hygiene instructions, 48.0% presented dental calculus, 5.2% had pockets of 4–5mm, and 0.6% had pockets of 6–7mm, for which they needed root scaling and radicular polishing. Regarding periodontal fixation loss, 50.2% were healthy, 40.8% had losses of 4–5mm, and 7.8% losses of 6–8mm. No association was found between periodontal health status and oral health self-assessment (Table 3).

DISCUSSION.

Here we aimed to determine the characteristics of self-assessment overall oral health status, the clinical dental caries experience, periodontal status, periodontal loss fixation and to investigate the association between self-reported and clinical oral health status among Paraguayans adults.

Negative oral health self-perception was associated with a high DMFT index, of which the decayed component was the only one that presented statistically significant differences. This could be because the presence of caries in the patients of this study is accompanied by pain. In another study, a negative self-evaluation has been associated with the presence of pain and difficulty in pronouncing words as an important trigger factor.¹¹

The DMFT Index in this study (11.43) was higher than that reported for Lingua Indian Paraguayan Chaco (20-to 40-years-old) in 19845; higher (7.33) than patients attending the School of Dental Medicine in Case Western Reserve University,¹² higher (6.38) than Syrian and Iranian refugees in Germany,¹³ and higher (6.83) than refugees in a region of the federal state of Schleswig-Holstein, Germany,¹⁴ and lower to that reported in adults between 20 and 64 years old in Piracicaba, Saint Paul, Brasil,¹⁵ and in Chilean adults(14.6).¹⁶

Regarding dental caries experience, Paraguayan adults have lower FT (3.56) than Latino immigrants groups in Los Angeles, United States (6.29), but higher than an adult female population in Nigeria and Syrian, and Iranian (1.5) refugees in Germany. Paraguayans adults have a mean of DT (2.55) similar to Latino adult immigrants (2.47), lower than refugees in Germany (2.55), and higher than the female population in (1.8) Nigeria.^{13,17,18} The mean edentulous value in this study (1.2) was lower than the 3.5 reported for adults in Southern Vietnam.¹⁹ This could be due to the fact that Paraguay is a developing country where access to

health services continues to be limited, therefore, this is reflected by seeing almost half of filled teeth compared to Latinos in another country where there is greater coverage in Health. But this country still maintains an intermediate level of caries experience.

In our study, the proportion of patients with good, very good and excellent in their self-assessment (58.8%) was lower (64%) than of immigrants. Slightly higher than the 51% reported for adults in Yemeni.²⁰ A lower percentage in our study were rated as having poor oral health (12%) compared than the 27% of the elderly reported as having poor oral health among institutionalized adults aged 65 years or older in Norway.

The diversity of the results could be explained due to the low access to dental health services, since limited access has been reported in the Itapúa Department as reported by Caballero *et al.*,²¹ That is not alien to the reality of other departments of the country, that is, because the number of dentists varies considerably by health region according to data from the Directorate of Surveys, Statistics and Census of Paraguay.²²

This study included patients from the department of Paraguari. In addition, there are other influencing factors such as the wide diversity of ages of study participants, their level of knowledge about oral health and their awareness about the importance of oral health. Studies have reported that there is concordance between what is reported and the dental clinic.^{8,23}

In our study, no differences in perception were found by periodontal health status, while a study carried out in Turkish adults did find a statistically significant difference carried out by Beşiroğlu *et al.*,²⁴ Although close to half of the patients had dental calculus and periodontal loss fixation, we observed many patients with gingival retraction but relatively few with a periodontal pocket, which could explain this finding. In a Chinese study including patients less than 18-years-old, 93.5% presented with dental calculus,²⁵ which is almost double the value recorded in our study.

The limitations of our study include its small sample and the non-inclusion of indigenous populations, as well as the absence of data on other determinants of oral health status. The sample only covers two health regions and the non-inclusion of indigenous populations, which is why it would not be prudent to extrapolate to the entire population. However, the sample could be considered representative in

terms of social strata, since it includes a very socio-economically disadvantaged sector such as the Barrio Ricardo Brugada known as “La Chacarita”. The patients who attended FOUNA are from different parts of the country, who go for affordable costs and good referrals, as well as the patients who came from the Pirayú health center who reside in the interior of the country, where there are very few private dental practices. There is inequity in access to health services.

CONCLUSION.

Negative oral health self-perception was associated with a high DMFT index, of which the carious component was the only one that presented statistically significant differences.

Conflict of interests: The authors report no current or potential conflicts of interest. This article is part of the thesis of CDR to obtain the PhD in Methodology of Biomedical and Public Health Research at the Universitat Autònoma de Barcelona, Spain.

Ethics approval: All the authors declare no conflict of interest.

Funding: No funding.

Authors' contributions: Contributed to conception and design, data collection, analysis and interpretation, and drafted the manuscript: CDR. Contributed to design and critically revised the manuscript: JRM and IC. All authors gave final approval and agreed to be accountable for all aspects of work.

Acknowledgements: The authors wish to acknowledge the contribution to DDS. Milner Morel, DDS, and Leticia Cataldi, DDS regarding data collection.

REFERENCES.

1. Pan American Health Organization. Health in the Americas+, 2017 Edition. Summary: Regional Outlook and Country Profiles. Washington, D.C. PAHO, 2017.
2. Dirección General de Encuestas Estadísticas y Censos. Aprovechar ahora la juventud: un análisis del bono demográfico del Paraguay. Período 2000 - 2025. Asunción: DGEEC; 2016.
3. Coale AJ. The Decline of Fertility in Europe Since the Eighteenth Century as a Chapter in Human Demographic History. En: Watkins SC, Coale AJ, editores. The Decline of Fertility in Europe. United States of America: Princeton University Press; 2017.
4. Fischman S. Oral health in the republic of Paraguay. *Community Dent Oral Epidemiol.* 1974;2(4):176-81.
5. Kieser JA, Preston CB. Oral health status of the Lengua Indians of Paraguay. *Community Dent Oral Epidemiol.* 1984;12(6):406-7.
6. Ministerio de Salud Pública y Bienestar Social. Encuesta nacional de salud bucal del Paraguay, ENSABUD PY 2017: del diagnóstico al diseño y la ejecución de políticas públicas de salud bucodental. Asunción: OPS/OMS; 2018.
7. Lira R, Åkerman S, Klinge B, Boström EA, Gustafsson A. Salivary microbial profiles in relation to age, periodontal, and systemic diseases. *PLoS One.* 2018;13(3):1-15.
8. Blizniuk A, Ueno M, Zaitso T, Kawaguchi Y. Association between self-reported and clinical oral health status in Belarusian adults. *J Investig Clin Dent.* 2017;8(2):1-6.
9. World Health Organization. Oral health surveys: basic methods. 5th Ed. Geneva: WHO; 2013.
10. Holtfreter B, Alte D, Schwahn C, Desvarieux M, Kocher T. Effects of different manual periodontal probes on periodontal measurements. *J Clin Periodontol.* 2012;39(11):1032-41.
11. Arantes R, Frazão P. Subjective oral symptoms associated with self-rated oral health among Indigenous groups in Central-West Brazil. *Community Dent Oral Epidemiol.* 2018;46(4):352-9.
12. Baskaradoss JK. Relationship between oral health literacy and oral health status. *BMC Oral Health.* 2018;18(1):172.
13. Solyman M, Schmidt-Westhausen AM. Oral health status among newly arrived refugees in Germany: A cross-sectional study. *BMC Oral Health.* 2018;18(1):1-9.
14. Goetz K, Winkelmann W, Steinhäuser J. Assessment of oral health and cost of care for a group of refugees in Germany: A cross-sectional study. *BMC Oral Health.* 2018;18(1):1-8.
15. Manoelito Ferreira S-J, Correia de Sousa CA, Batista MJ, Rosário de Sousa LM. Condição de saúde bucal e motivos para extração dentária entre uma população de adultos (20-64 años). *Cien Saude Colet.* 2017;22(8):2693-702.
16. Quinteros ME, Cáceres DD, Soto A, Mariño RJ, Giacaman RA. Caries experience and use of dental services in rural and urban adults and older adults from central Chile. *Int Dent J.* 2014;64(5):260-8.
17. Spolsky VW, Marcus M, Der-Martirosian C, Coulter ID, Maida CA. Oral health status and the epidemiologic paradox within latino immigrant groups. *BMC Oral Health.* 2012;12(1).
18. Lawal F, Alade O. Dental caries experience and treatment needs of an adult female population in Nigeria. *Afr Health Sci.* 2017;17(3):905-11.
19. Nguyen TC, Witter DJ, Bronkhorst EM, Truong NB, Creugers NHJ. Oral health status of adults in Southern Vietnam - a cross-sectional epidemiological study. *BMC Oral Health.* 2010;10(2).
20. Alhajj MN, Halboub E, Amran AG, Alkheraif AA, Al-Sanabani FA, Al-Makramani BM, Al-Basmi AA, Al-Ghabri FA. Link between perceived oral and general health status among Yemeni adult dental patients. *BMC Oral Health.* 2019;19(1):1-9.
21. Caballero-García CR, Espínola-Verdún PA, Domínguez-González DD, Martínez-Benítez GG, Figueredo-Palacios S, Fernández-Cáceres AM, et al. Salud bucodental y utilización de servicios odontológicos. *Memorias del Inst Investig en Ciencias de la Salud.* 2018;15(3):57-63.
22. Dirección General de Encuestas Estadísticas y Censos. Compendio Estadístico 2014. Fernando de la Mora: DGEEC; 2015.
23. Kim SY, Kim JE, Kim HN, Jun EJ, Lee JH, Kim JS, Kim JB. Association of Self-Perceived Oral Health and Function with Clinically Determined Oral Health Status among Adults Aged 35-54 Years: A Cross-Sectional Study. *Int J Environ Res Public Health.* 2018;15(8):1681.
24. Beşiroğlu E, Lütüoğlu M. Relations between periodontal status, oral health-related quality of life and perceived oral health and oral health consciousness levels in a Turkish population. *Int J Dent Hyg.* 2020;18(3):251-60.
25. Lu H-X, Wong MCM, Lo ECM, McGrath C. Oral Health Related Quality of Life Among Young Adults. *Appl Res Qual Life.* 2015;10(1):37-47.