

Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics



www.renhyd.org



RESEARCH ARTICLE

Development and characterization of a ready-to-eat vegetable *millefeuille* enriched with polyphenols

Rosa A Abalos^{a,*}, M Victoria Aviles^{a,b}, Elisa F Naef^b, M Beatriz Gómez^{a,b}

^aFacultad de Bromatología, Universidad Nacional de Entre Ríos, Gualeguaychú, Argentina.

^bInstituto de Ciencia y Tecnología de los Alimentos, CONICET-UNER, Gualeguaychú, Argentina.

*rosa.abalos@uner.edu.ar

Assigned Editor: Édgar Pérez Esteve, Universitat Politècnica de València, Valencia, España.

Received: 12/08/2022; accepted: 03/08/2023; published: 04/11/2023.

KEYWORDS

Entry Term(s)

Food design;
Vacuum
impregnation;
Sous vide;
Commercial life.

Development and characterization of a ready-to-eat vegetable *millefeuille* enriched with polyphenols

ABSTRACT

Introduction: The development of high-quality plant-based ready-to-eat food options can respond to today's dietary demands. The trend of consumers opting for a plant-based diet is on the rise; however, they have few options for ready-to-eat products. The aim of this study was to develop a ready-to-eat plant product enriched with polyphenols using vacuum technologies.

Methodology: The physicochemical, sensory and microbiological characteristics of the product were evaluated. A lasagna was designed (layers of vegetables of regional origin –sweet potato (*Ipomea batata* L.), zucchini (*Curcubita pepo* L.) and dried tomato– interspersed with a binder and cheese) ready to eat, which were vacuum impregnated with an extract of polyphenols (95% oligomeric proanthocyanidins) from grape seeds. The concentration of polyphenols was measured by the Folin-Ciocalteu colorimetric method. The commercial life in refrigeration (0-3 °C) of the culinary preparation was determined, evaluating pH, color, texture and microbial count of aerobic and anaerobic psychrotrophic bacteria, lactic acid bacteria, molds and yeasts following techniques established in ISO standards. The sensory evaluation was carried out with regular consumers of ready-to-eat products. They evaluated the appreciation of sensory attributes and the general acceptability of the sample on a 9-point hedonic scale.

Results: In this study, a ready-to-consume preparation with 220 mg of GAE/100 was designed, which could contribute to the daily intake of polyphenols. The values of weight variation and pH (from 5.2 to 5.5) did not present significant variations during the 21 days of storage studied. The surface color parameters (L*, a* and b*) did not change until day 7. The firmness of the culinary preparation decreased over the storage time (from 4.2 to 2.9 N/mm). The microbial counts were consistent with those of an innocuous product (less than 10 CFU/g). This study demonstrated that the *sous vide* cooking method improved the refrigerated storage time of the preparation since the microbial load was maintained at safe levels. The sensory evaluation showed a positive acceptance by consumers for most of the sensory characteristics of the designed product.

Conclusions: The technologies used, *sous vide* and vacuum impregnation, could enhance the gastronomic quality of the product, preserve the organoleptic and nutritional properties and guarantee its safety. The designed product could be an option to incorporate ready-to-eat vegetable dish preparations into the market and thus contribute to increasing the consumption of vegetables.

Funding: The work was carried out with the financial support of the University of Entre Ríos (UNER).



PALABRAS CLAVE

Entry Term(s)

Diseño de alimentos;

Impregnación al vacío;

Sous vide;

Vida comercial.

Desarrollo y caracterización de una milhoja de verdura lista para consumir enriquecida con polifenoles

RESUMEN

Introducción: El desarrollo de opciones de alimentos listos para el consumo a base de plantas de alta calidad puede responder a las demandas dietéticas actuales. La tendencia de los consumidores que optan por una dieta basada en plantas va en aumento; sin embargo, tienen pocas opciones de este tipo de productos. El objetivo de este estudio fue desarrollar un producto vegetal listo para consumir enriquecido con polifenoles utilizando tecnologías de vacío.

Metodología: Se evaluaron las características fisicoquímicas, sensoriales y microbiológicas del producto. Se diseñó una milhoja (capas de vegetales de origen regional –batata (*Ipomea batata* L.), zucchini (*Cucurbita pepo* L.) y tomate seco– intercaladas con un aglutinante y queso) lista para comer, las cuales fueron impregnadas al vacío con un extracto de polifenoles (95% de proantocianidinas oligoméricas) de semillas de uva. La concentración de polifenoles se midió por el método colorimétrico de Folin-Ciocalteu. Se determinó la vida comercial en refrigeración (0-3 °C), evaluando pH, color, textura y conteo microbiano de bacterias psicrótroficas aerobias y anaerobias, bacterias ácido lácticas, mohos y levaduras siguiendo técnicas establecidas en normas ISO. La evaluación sensorial se realizó con consumidores habituales de productos listos para el consumo. Evaluaron la apreciación de los atributos sensoriales y la aceptabilidad general de la muestra en una escala hedónica de 9 puntos.

Resultados: Se logró diseñar una preparación lista para el consumo con 220 mg of GAE/100, que podría contribuir a la ingesta diaria de polifenoles. Durante 21 días de almacenamiento no hubo variaciones en lo referente al cambio de peso ni de pH, que se mantuvo entre 5,2 y 5,5. En cuanto al color, se pudo observar que los parámetros L*, a* y b* se mantuvieron constantes hasta el día 7. La firmeza de la preparación culinaria fue disminuyendo a medida que transcurrió este tiempo de almacenamiento (de 4,2 a 2,9 N/mm). Los recuentos microbianos resultaron con valores que se corresponden con un producto inocuo (menos de 10 UFC/g). Este estudio demostró que el método de cocción *sous vide* mejoró el tiempo de almacenamiento refrigerado de la preparación ya que la carga microbiana se mantuvo en niveles seguros. La evaluación sensorial mostró una aceptación positiva por parte de los consumidores para la mayoría de las características sensoriales del producto diseñado.

Conclusiones: Las tecnologías utilizadas, *sous vide* e impregnación al vacío, podrían potenciar la calidad gastronómica del producto, conservar las propiedades organolépticas y nutricionales y garantizar su inocuidad. El producto diseñado podría ser una opción para incorporar al mercado preparaciones de platos de verduras listas para comer y así contribuir a incrementar el consumo de verduras.

Financiación: La obra se realizó con el apoyo financiero de la Universidad de Entre Ríos (UNER).

KEY MESSAGES

1. Vacuum technologies allowed the design of a ready-to-eat vegetable-based dish with 200.23 mg EAG/100.
2. The *sous vide* cooking method used allowed to improve the conservation time in refrigeration.
3. The *sous vide* cooking kept the microbial load at safe levels for 21 days.
4. Sensory evaluation showed that texture and flavor significantly influenced the overall acceptability of *millefeuille*.

CITATION

Abalos RA, Aviles MV, Naef EF, Gómez MB. Development and characterization of a ready-to-eat vegetable *millefeuille* enriched with polyphenols. Rev Esp Nutr Hum Diet. 2023; 27(2): 163-72. doi: <https://doi.org/10.14306/renhyd.27.2.1839>

INTRODUCTION

Vegetables are important components of a healthy diet as they are a source of vitamins, minerals, dietary fiber, and phytonutrients. There is a consensus that low consumption of fruits and vegetables is associated with poor health and an increased risk of non-communicable diseases¹. People are changing their eating habits by increasing their daily intake of vegetables. However, the consumer who spends less time cooking finds practical problems such as lack of cooking skills, the effort required to prepare vegetable dishes, or a lack of available ready-to-eat plant-based foods, which makes it difficult to incorporate healthy products that contain vegetables².

Several studies suggest that facilitating consumer access to fruits and vegetables through the design of refrigerated precooked products increases the consumption of this particular food group³. Therefore, the design of precooked plant foods of high nutritional quality can contribute to improving the quality of the diet by helping to overcome these barriers in the preparation of vegetable dishes⁴. Ready-to-eat plant-based products are attractive to consumers looking for healthy and convenient meals, especially if they contain ingredients that play a specific role in the physiological functions of the human body⁵.

Vacuum impregnation is a methodology that has been studied to incorporate physiologically active compounds to improve the nutritional quality of vegetables. This technique consists in the exchange of the gas present in the food pores for the external liquid due to the action of hydrodynamic mechanisms (HDM) promoted by pressure changes⁶. In this study, this methodology was used to incorporate polyphenols into the preparation and increase its nutritional quality. These compounds are currently of great nutritional interest because they contribute to preventing various diseases related to oxidative stress⁷. Several studies have corroborated the beneficial effect of consuming polyphenols added to a diet rich in vegetables⁸.

The *sous vide* (SV) cooking technique facilitates the preparation of ready-to-eat vacuum-packed products and consists in using temperatures below 100 °C for long periods, immediate cooling, and refrigeration at 3 °C until the dish is served. This technique preserves natural sensory qualities and nutritional value and prolongs shelf life. The precise temperature control in this cooking method provides more options for cooking and texture than traditional methods⁹. Chiavaro *et al.*¹⁰ demonstrated that *sous vide* cooking of carrots can preserve and/or improve nutritional quality; Iborra-Bernard *et al.*¹¹ corroborated that the samples of *sous vide* red cabbage were tastier and more purple and that they retained more anthocyanins than those cooked by conventional treatments. Alcusón *et al.*¹² reported that *sous vide*

cooking improved the total phenolic content, antioxidant activity, color, and texture of borage (*Borago officinalis* L.) stems.

Studies on the mechanisms of spoilage that occur in food during storage are the subject of continuous and exhaustive research. An important aspect of food development is knowing its stability during storage under the conditions determined for commercialization¹³. In this sense, *sous vide* technology is capable of increasing the useful life of products in comparison to other traditional methods¹⁴. Few published studies have focused on ready-to-eat vegetable-based products cooked by *sous vide*. However, this cooking technique has been reported for other food matrices, especially products of animal origin¹⁵.

In view of these considerations, the aims of this study were to: 1) study were to design a ready-to-eat *millefeuille* with functional characteristics by applying vacuum technologies, 2) analyze the effect of refrigerated storage on its physicochemical parameters, 3) evaluate the main sensory parameters.

METHODOLOGY


The experimental design was divided into three phases, as shown in the scheme in Figure 1.

Materials

In the design of the preparation, the following ingredients were used: sweet potato (*Ipomoea batatas* L.), zucchini (*Cucurbita pepo* L.), dried tomatoes, eggs, powdered milk, cornstarch and Danbo cheese. All the ingredients were bought at the local market. The vegetables were selected considering their freshness, firmness and absence of microbial or mechanical damage. They were sanitized with 0.1% sodium hypochlorite (1 mL/L) and chopped. Sweet potato slices 3 mm thick and 50 mm in diameter were vacuum impregnated (VI) with polyphenols following the technique proposed by Abalos *et al.*¹⁶. The polyphenol solution used was Vitolol® from the national company Nialtec S.A. This polyphenol extract contains 95% oligomeric proanthocyanidins. The formulation of the 3.5% VI solution was made based on the study carried out by Rózek *et al.*¹⁷. For the VI experiments, the Gastrovac® vacuum cooking equipment was used. The sweet potatoes were immersed in the VI solution in the equipment and subjected to vacuum for 25 minutes at 20 °C. Pressure was restored at the same time intervals as that of the LV.

Millefeuille preparation

To prepare the *millefeuille*, the following proportion of ingredients was used: sweet potatoes vacuum impregnated with polyphenols

Figure 1. Experimental design.


(35%) and Zucchini (24%) in slices 3 mm in thickness and 40 mm in diameter, dried tomatoes (11%), Danbo cheese (21%), and a binder—milk, cornstarch, and egg—(9%). For the assembly, the ingredients were interspersed in layers (Figure 2), resulting in 285 g servings of *millefeuille*. The preparation was vacuum packed (VACPACK, ICC, 80016™, Spain) in polyamide-polyethylene bags (O₂ permeability, 25 to 30 cm³/m²/day; water vapor permeability of 5 g/m²/day). The heat treatment was by *sous vide* cooking in a constant water-circulation bath with temperature and time regulation (RONER COMPACT, ICC, 80060, Spain) at 80 °C for 45 minutes. It was subsequently rapidly cooled in a water bath at 0 °C for 5 minutes. The cooking temperature was monitored with a thermocouple (HANNA Instruments, HY 93530N, Italy). The cooked samples were refrigerated at 2 °C, until further study.

Physicochemical characterization

The ready-to-eat *millefeuille* with the addition of polyphenols was characterized physicochemically, in triplicate, as follows, the compositional properties of the SRB were determined according to AOAC (Association of Official Analytical Chemist)¹⁸. Moisture content was determined after heating in an infrared drying oven (RADWAG, Mag. 50/WH, Poland) according to 7.003-8421.

Protein was estimated by the Kjeldahl method described in AOAC 928.08, with a conversion factor of 6.25 for protein. Total fat was determined by the Soxhlet solvent extraction gravimetric method, following method official AOAC 965.33. Total carbohydrates were estimated by subtraction of the percentage of protein, lipid, ash from the total dry weight.

For polyphenols, total phenol content was determined by the reagent Folin-Ciocalteu method, using the test Heimer, Vignolini, Dini and Romani¹⁹. Samples of sweet potatoes of 16 g were processed in a blender with 50 mL of methanol (Sintorgan, Lot. N° 38931). The homogenates obtained were placed in a beaker and heated at 40 °C with continuous stirring (Barnstead Thermoline) for 15 minutes before centrifugation (Rolco, CM 2036.4, Argentina) for 5 minutes at 11.200 g force and filtration. A 0.5-mL aliquot was added to a test tube together with 1 mL of Folin-Ciocalteu reagent (Biokar) and 10 mL of distilled water, left to stand for 2 minutes. Then, 4 mL of a 20% (w/v) sodium carbonate solution (Merck, Argentina) was added, brought up to a final volume of 25 mL, and heated at 50 °C for 5 minutes in a thermostatic bath (Vicking SRL, Masson, Argentina). The optical density was finally measured in a spectrophotometer (Jenway Mod. 6505 UV/VIS™) at 765 nm¹⁶.

Figure 2. Ready-to-eat vegetable-based dish enriched with polyphenols preparation.

The pH was measured using a pH-meter with a combined Ag/AgCl electrode (ORION, S.A. 720, United States). The pH measurement was performed by immersing the electrode in a homogenate of 10 g of *millefeuille* in 100 mL of distilled water. Soluble solids (°Bx) were estimated using a table refractometer (A. KrussOptronic, Germany) according to AOAC 932.1221. Acidity was determined by the volumetric method. Water activity was measured with a hygrometer (Rotronic, Hygrolab C1, Switzerland) according to AOAC method 978.18.

Sensory evaluation

To determine the consumer acceptability of the product, the smell, color, texture, and taste parameters were evaluated. The *millefeuille*, cooked the day before tasting and kept refrigerated, was heated to the usual temperature for consumption (65 °C), cut into 5 cm/5 cm portions, and served on white plates. The sensory evaluation was carried out by a panel of 105 people, selected for being regular consumers of ready-to-eat products, in the laboratory facilities of the Faculty of Bromatology, following the ISO 8589 standard²⁰. Consumers were asked to indicate their level of appreciation for the sensory attributes and the overall acceptability of the sample on a 9-point hedonic scale ranging from 1 = "I don't like it very much" to 9 = "I like it very much." The study was approved by an Institutional review board of the National University of Entre Ríos (RESOLUTION "C.D." N° 017/17).

Evolution of physicochemical and microbiological parameters during refrigerated storage

Twelve servings of *sous vide* cooked *millefeuille* were prepared and stored in a refrigerator (2 °C). The temperature was monitored throughout the storage period with a digital thermometer (Elitech, D-T1, China). The samples were analyzed physicochemically and microbiologically at different times (0, 7, 14, and 21 days) by the determinations reported below.

Physicochemical analysis: Polyphenols and pH were determined following the techniques previously mentioned. Weight variation: samples were weighed using a precision balance (OHAUS, ADVENTURES, Canada), and the weight variation (ΔP) was calculated as the percentage of variation between the weight of the samples on days 7, 14, and 21 with respect to the weight of samples on day 0.

Color was determined using a colorimeter (HUNTER ASSOCIATES LABORATORY, Inc., MiniScan EZ Model, EE UU) in the CIE L* a* b* color space. The variation was evaluated in the parameters L* (black 0, white 100), a* (red-green), and b* (yellow-blue). The color parameters were determined from triplicate measurements in each sample. Color intensity C*ab (Equation 1) and color change ΔE (Equation 2) were calculated, where L₀, a₀, and b₀ represent the readings of the non-impregnated samples.

$$\text{Equation 1: } C^*ab = \sqrt{a^2 + b^2}$$

$$\text{Equation 2: } \Delta E = \sqrt{(L_0 - L)^2 + (a_0 - a)^2 + (b_0 - b)^2}$$

Texture was determined by a compression test using a texturometer (Instron 3342™, Massachusetts, USA). The values obtained for the maximum compression force were expressed in Newtons (N). The tests were carried out in quintuplicate, with 30% deformation and at a test speed of 0.5 mm/s. The firmness (strength) of the material was calculated from the relationship between the maximum force (N) and the maximum deformation (mm). The elastic modulus was calculated from the relationship between the maximum force (N) and the initial module (mm).

Microbiological analysis: To perform the microbiological analysis of the *millefeuille*, a 10-g sample was weighed in a sterile bag and diluted with 90 mL of buffered peptone water (Acumedia, Neogen Corporation, Michigan, USA). A 1/10 dilution was obtained after homogenization. Counts were done in duplicate.

To estimate molds and yeasts, a culture medium composed of oxytetracycline, gentamicin, yeast extract, and glucose (OGY) (Merck KGaA, Germany) was used. The inoculated plates were incubated at 25 °C for 5 days, according to ISO 17410²¹.

To determine total aerobic psychrotrophic bacteria, the Plate Count Agar (PCA) culture medium (Britania, Argentina) was used. The inoculated plates were incubated for 7 days in a cold room at 4 °C.

Determination of total anaerobic psychrotrophic bacteria was carried out in Tryptone Soya Agar (TSA) medium (Oxoid, Basingstoke, United Kingdom). The inoculated plates were placed in an anaerobic jar with a capacity of 2.5 L and incubated in a cold room at 4 °C for 7 days according to ISO 21527-2²².

At the end of the incubation period, agar plates with dilutions that allowed the best Colony Forming Unit (CFU) counts were selected. The counts were expressed as log CFU/g of product for the different groups of microorganisms analyzed.

Statistical analysis

The data of the test results are expressed as means ± standard deviations, as computed by the software XLSTAT (New York, USA) program for Windows, version 2018.7.525²³. The significance of differences between the means was determined by the analysis of variance ANOVA followed by the Tukey multiple-comparison test with a limit of significance at $p < 0.05$.

Multiple regression was performed to analyze the overall acceptability by considering the influence of the specific sensory attributes (color, taste, texture, and smell), using the Minitab 18 Statistical Software (LLC, United States).

RESULTS

Ready-to-eat *millefeuille* enriched with polyphenols

This work was based on the design of a ready-to-eat vegetable-based preparation using vacuum impregnation (VI) to incorporate polyphenols and *sous vide* cooking as the heat treatment (Figure 3).

The physicochemical characterization of the designed culinary preparation is shown in Table 1. The ready-to-eat product obtained provides 200.23 mg GAE/100 g per serving, through the VI.

Figure 3. Ready-to-eat vegetable-based dish enriched with polyphenols.



Table 1. Physicochemical and nutritional parameters of the culinary preparation: ready-to-eat *millefeuille* with the addition of polyphenols.

Portion of 285 g	Amount per proportion Mean (SD) (n=3)
Protein (g/100 g)	18 g (0.32)
Total fat (g/100 g)	15 g (0.24)
Polyphenols (mg GAE/100 g)	200.23 (0.85)
Moisture (g/100 g)	89.79 (0.24)
pH	5.38 (0.03)
Soluble solids (° Brix)	7.10 (0.10)
Acidity (g/mL)	0.19 (0.15)
Water activity	0.99 (0.01)

Sensory evaluation

For the sensory evaluation, a group of 105 consumers participated in the study, of which 65% were women and 35% men. The age ranges were 18-30 years (54%), 31-59 years (35%), and 60 years or older (12%).

The results presented in Table 2 show the level of satisfaction of each parameter studied. It was observed that the general acceptability, taste, smell and color were indicated in the "I like" category, with color obtaining the highest score. The texture was identified as different from the rest corresponding to the category "I am indifferent about it".

Table 2. Descriptive statistics.

Sensory characteristics	Vegetable <i>millefeuille</i> Mean (SD)
Global acceptability	3.56 (0.67) ^a
Taste	3.90 (0.91) ^a
Smell	3.96 (0.71) ^a
Color	4.12 (0.70) ^a
Texture	3.27 (1.09) ^b

9-point hedonic scale evaluation.

*Scores with different superscripts are significantly different according to Tukey's test at a confidence level of 95%.

Table 3 shows that texture and taste significantly influence the other sensory characteristics (smell and color) as indicated by the significant differences between the two means ($p < 0.05$). The main sensory characteristics that influence the consumer in the global acceptability was the texture and flavor of the product.

Table 3. Multiple regression values corresponding to the sensory attributes of vegetable *millefeuille*.

Sensory characteristics	Global acceptability P
Smell	0.74
Color	0.31
Texture	0.02
Taste	0.03

$P < 0.05$.

Physicochemical and microbiological parameters during refrigerated storage

These physicochemical and microbiological parameters were analyzed over 21 days, estimated useful lifetime for vacuum-packed vegetables cooked by *sous vide*. The parameters evaluated are presented in Table 4.

It was observed that the polyphenol content remained constant until day 14. It was observed that the studied product did not experience changes in pH or weight ($p > 0.05$) during the storage period. Regarding the color parameters, it was observed that the values of a^* and b^* remained constant until day 7. The color intensity and luminosity during storage did not show a significant difference ($p > 0.05$) in the evaluated time. The ΔE^* remained constant until day 14. Taking into account the evolution of the values, color was observed to decrease over time.

The uniaxial compression test, it was observed that the values of firmness and maximum force remained constant until day 14 and then began to decrease.

The microbiological parameters studied showed that there was no proliferation of the groups of microorganisms evaluated, possibly due to the absence of oxygen, the low refrigeration temperature, and the appropriate conditions of the *sous vide* cooking process. In this way, the product could be considered suitable for human consumption during the period studied.

DISCUSSION

The ready-to-eat product obtained provides 200.23 mg GAE/100 g per serving, through the VI. In previous studies, the times and conditions of VI (vacuum and restoration, times and concentration of the impregnation solution) were determined to impregnate the sweet potato slices used in the design of the culinary preparation¹⁶. The literature shows that vegetables rich in polyphenols have a concentration ranging between 170-930 mg of GAE/100 g¹⁷. Data analysis indicated that the impregnated sweet potato represents an increase in phenolic-compound concentration over the control sample of 473%. Thus, this method could be considered as suitable for incorporating biologically active substances into vegetables, so as to facilitate the improvement of the nutritional quality by those supplements.

Although the recommended values for the population have not yet been stipulated for polyphenols, various studies state that it is necessary to increase daily consumption. Many polyphenols have free radical scavenging properties, giving them

antioxidant activity, which could be related to the prevention of non-communicable diseases⁸. Some authors have verified the feasibility of incorporating polyphenols using VI. Moreira and Almohaimeed²⁴ added polyphenols to healthy potato chip snacks using VI and vacuum frying technology. Tappi *et al.*²⁵ used a green tea extract to enrich the apples minimally processed by VI to obtain a nutritionally fortified product. In this way, VI can be considered an effective technique to design new products with functional components.

In the study by Aviles *et al.*²⁶, two products cooked by *sous vide*, a vegetable lasagna and a hamburger, obtained the rating "I like it slightly", results similar to those observed in this study regarding general acceptability of the product. The sensory evaluation of the main parameters that consumers judge for the acceptability of products –smell, texture, color, and taste– was carried out²⁷. Visual appearance and color are usually the first sensory stimuli that are presented to consumers and they influence expectations about food. The visual properties can produce positive or negative sensations that lead to the acceptance or rejection of food, respectively²⁸. Firmness is one of the main factors that consumers use to define whether a vegetable is properly

cooked²⁷. Consumers evaluated the acceptability of the texture by expressing "I do not like or dislike it", and comments such as "it feels crunchy, but some parts seemed to be raw", "it could be tenderer", "it seemed tough to me, kind of raw" were highlighted. This perception of a raw product with a tough texture could be attributed to the *sous vide* method, characterized by better preserving the texture of vegetables, unlike traditional cooking methods, in which the toughness constantly decreases. The acceptability of new products is determined by conditions related to food and consumers. The appearance, texture, and taste of the products are fundamental characteristics to establish favorable sensory and hedonic responses for the consumer²⁹.

To determine the refrigerated storage time of the preparation, the parameters that determine the quality of food and affect the sensory perception and acceptability of the consumer (variation of weight, pH, color, and texture) were evaluated³⁰. The parameters that indicate the quality and safety (microbial count) of food, as well as the compound added through VI (polyphenols), were also measured. These physicochemical and microbiological parameters were analyzed over 21 days, estimated useful lifetime for vacuum-packed vegetables cooked by *sous vide*³¹.

Tabla 4. Parameters evaluated in ready-to-eat *millefeuille* with the addition of polyphenols.

Parameters	Day 0	Day 7	Day 14	Day 21
Polyphenols (mg EAG/100 g)	200.23 (0.85) ^a	199.96 (0.69) ^a	199.55 (2.03) ^a	189.69 (1.40) ^b
pH	5.25 (0.05) ^a	5.42 (0.03) ^a	5.40 (0.10) ^a	5.50 (0.20) ^a
Color				
L*	48.37 (1.97) ^a	47.81 (0.32) ^a	47.50 (0.52) ^a	47.87 (1.43) ^a
a*	21.07 (1.93) ^a	19.73 (0.03) ^a	18.60 (0.82) ^b	16.97 (0.45) ^c
b*	49.59 (13.18) ^a	46.65 (5.20) ^a	45.20 (1.23) ^b	40.64 (0.94) ^c
ΔE*	-	15.690 (9.85) ^a	15.86 (11.06) ^a	12.29 (9.02) ^b
C*ab	47.82 (1.28) ^a	41.62 (0.53) ^a	41.53 (1.28) ^a	44.03 (1.12) ^a
Texture				
Initial module (mm/mm)	25.94 (2.68) ^a	23.57 (0.05) ^a	20.77 (0.10) ^b	18.30 (0.61) ^b
Hardness (N)	6.16 (0.05) ^a	6.05 (0.06) ^a	6.16 (0.06) ^a	6.11 (0.08) ^a
Maximum deformation (mm)	4.21 (0.40) ^a	3.89 (0.26)	3.69 (0.36) ^b	2.99 (0.11) ^b
Mohos y levaduras (UFC/g)	< 10	< 10	< 10	< 10
Lactobacilos (UFC/g)	< 10	< 10	< 10	< 10
Aerobic psychrotrophic (UFC/g)	< 10	< 10	< 10	< 10

*Average value ± standard deviation of the results (n=3). Samples with the same letter in the same column were not significantly different by the Tukey test (p<0.05).

In this study it was observed that the polyphenol content remained constant until day 14, while Tappi *et al.*²⁵ studied the quality and stability of minimally processed apples impregnated with green tea polyphenols, finding polyphenol values without significant modification until day 7 of storage.

The pH affects the properties of food and can be an indicator of growth or mortality of microorganisms, inactivation of bacterial spores, and chemical reactions such as Maillard's³⁰. In this study, the pH obtained did not present changes during the storage period studied. Therefore, control of effects during storage is necessary to safely produce quality and value-added products.

CONCLUSIONS

In conclusion a ready-to-eat *millefeuille* was designed with 220 mg of GAE/100 g of polyphenols using vacuum technologies. The developed product maintained, in general, the initial quality and safety characteristics with a commercial life of at least 14 days stored in refrigerated conditions at 3 °C. The *sous vide* cooking method used allowed us to improve the conservation time in refrigeration of the preparation since the microbial load was kept at safe levels. The sensory attributes of taste, smell, texture and color analyzed were accepted by consumers, with texture being the most significant parameter that influences the acceptability of the product. It would be interesting to complement these results with sensory tests and the qualitative determination of polyphenols during storage and their relationship with antioxidant activity, in addition to other bioavailability experiments. Products of nutritional quality, such as the one studied in the present work, could be an option to incorporate ready-to-eat vegetable dish preparations into the market and thus contribute to increasing the consumption of vegetables.

ACKNOWLEDGMENTS

The authors acknowledge the financial support from the University of Entre Ríos (UNER). Dr. Donald F. Haggerty, a retired academic career investigator and native English speaker, edited the final version of the manuscript.

AUTHORS' CONTRIBUTIONS

R.A.A.: idea design, methodology, investigation, writing. E.F.N.: formal analysis, investigation, writing. M.V.A.: formal analysis, investigation, writing. M.B.G.: writing-review and editing, supervision.

FUNDING

The work was carried out with the financial support of the University of Entre Ríos (UNER).

COMPETING INTERESTS

The authors declare that there are no known conflicts of interest associated with this publication.

REFERENCES

- (1) World Health Organization. Prevention of cardiovascular disease: Guideline for assessment and management of cardiovascular risk. https://www.who.int/cardiovascular_diseases/publications/Prevention_of_Cardiovascular_Disease/en/. (July 7, 2021).
- (2) Reipurth MFS, Hørby L, Gregersen CG, Bonke A, Perez Cueto FJA. Barrier and facilitator toward adopting a more plant-based diet in a sample of Danish consumers. *Food Qual Prefer.* 2019; 73: 288-92. doi: 10.1016/j.foodqual.2018.10.012.
- (3) Bach Hyldelund N, Worck S, Olsen A. Convenience may increase vegetable intake among young consumers. *Food Qual Prefer.* 2020. doi: 103925. j.foodqual.2020.103925.
- (4) Contini C, Boncinelli F, Marone E, Scozzafava G, Casini L. Drivers of plant-based convenience foods consumption: Results of a multicomponent extension of the theory of planned behaviour. *Food Qual Prefer.* 2020; 84: 103931. doi: 10.1016/j.foodqual.2020.103931.
- (5) European Commission Community Research. Project Report: Functional food science in Europe. EUR-18591, in: Office for Official Publications of the European Communities, L-2985, Luxemb, 2000.
- (6) Fito P, Chiralt A, Barat JM, Andrés A, Martínez-Monzó J, Martínez-Navarrete N. Vacuum impregnation for development of new dehydrated products. *J Food Eng.* 2001; 49(49): 297-302. doi: 10.1016/S0260-8774(00)00226-0.
- (7) Godos J, Rapisarda G, Marventano S, Mistretta A, Grosso G. Association between polyphenol intake and adherence to the Mediterranean diet in Sicily, southern Italy. *NFS Journal.* 2017; 8: 1-7.

- (8) Grosso G, Stepaniak U, Micek A, Stefler D, Bobak M, Pająk A. Dietary polyphenols are inversely associated with metabolic syndrome in Polish adults of the hapiee study. *Eur J Nutr*. 2016; 56(4): 1409-20.
- (9) Baldwin E. Sous vide cooking: A review. *Int J Gastron Food Sci*. 2012; 1: 15-30.
- (10) Chiavaro E, Mazzeo T, Visconti A, Manzi C, Fogliano V, Pellegrini N. Nutritional Quality of Sous Vide Cooked Carrots and Brussels Sprouts. *J Agric Food Chem*. 2012; 60(23): 6019-25. doi: 10.1021/jf300692a.
- (11) Iborra-Bernad C, Tárrega A, García-Segovia P, Martínez-Monzó J. Advantages of sous-vide cooked red cabbage: Structural, nutritional and sensory aspects. *LWT - Food Science and Technology*. 2014; 56(2): 451-60. doi: 10.1016/j.lwt.2013.12.027.
- (12) Alcusón G, Remón S, Salvador ML. Quality related aspects of sous-vide processing of borage (*Borago officinalis* L.) stems. *LWT - Food Science and Technology*. 2017; 85: 104-9. doi: 10.1016/j.lwt.2017.07.012.
- (13) Wibowo S, Buvé C, Hendrickx M, Van Loey A, Grauwet T. Integrated science-based approach to study quality changes of shelf-stable food products during storage: A proof of concept on orange and mango juices. *Trends Food Sci Technol*. 2018; 73: 76-86. doi: 10.1016/j.tifs.2018.01.006
- (14) Ayub H, Ahmad A. Physicochemical changes in sous-vide and conventionally cooked meat. *Int J Gastron Food Sci*. 2019; 100145.
- (15) Diaz Molins P. Calidad y deterioro de platos "sous vide" preparados a base de carne y pescado y almacenados en refrigeración. Tesis de Doctorado. Universidad de Murcia. Departamento de Tecnología de Alimentos, Nutrición y Bromatología. Murcia, 2009.
- (16) Abalos RA, Naef EF, Aviles MV, Gómez B. Vacuum impregnation: A methodology for the preparation of a ready-to-eat sweet potato enriched in polyphenols. *LWT- Food Science and Technology*. 2020; 109773. doi: 10.1016/j.lwt.2020.109773
- (17) Rózek A, Achaerandio I, Güell C, López F, Ferrando M. Efecto del secado convectivo en la estabilidad de compuestos fenólicos añadidos a alimentos sólidos mediante deshidratación osmótica. *Memorias del II Congreso Iberoamericano sobre Seguridad Alimentaria y el V Congreso Español de Ingeniería en Alimentos*. Barcelona, España. 2018. <https://upcommons.upc.edu/handle/2117/2523>.
- (18) AOAC. Official methods of analysis, 18th Ed. Gaithersburg, Maryland. Association of Official Analytical Chemists, 20877e2417. USA, 2005, 270-310.
- (19) Heimer D, Vignolini P, Dini M, Romani A. Rapid test to assess the antioxidant activity of *Phaseolus vulgaris* L. Dry beans. *Journal of Agriculture*. 2005; 53(8): 3053-6.
- (20) ISO 8589:2007 Sensory analysis — General guidance for the design of test rooms. <https://www.iso.org/standard/36385.html>. (July 7, 2021).
- (21) ISO 17410. Microbiology of food and animal feedingstuffs. Horizontal method for the enumeration of psychotropic microorganisms. <https://www.iso.org/standard/30649.html>. (July 7, 2021).
- (22) ISO 21527-2. Microbiology of food and animal feedingstuffs - Horizontal method for the enumeration of yeasts and moulds. <https://www.iso.org/standard/38276.html>. (July 7, 2021).
- (23) Addinsoft. XLSTAT Statistical and Data Analysis Solutions. New York, USA. 2020. <http://www.xlstat.com>.
- (24) Moreira R, Almohaimeed S. Technology for processing of potato chips impregnated with red root beet phenolic compounds. *J Food Eng*. 2018; 228: 57-68. doi: 10.1016/j.jfoodeng.2018.02.010.
- (25) Tappi S, Tylewicz U, Romani S, Dalla RM, Rizzi F, Rocculi P. Study on the quality and stability of minimally processed apples impregnated with green tea polyphenols during storage. *Innov Food Sci Emerg Technol*. 2017; 39: 148-55. doi: 10.1016/j.ifset.2016.12.007.
- (26) Aviles MV, Naef E, Gómez B, Abalos RA. Consumer-consumption characteristics of ready-to-eat sous vide food products within the habitual context of the household. *Braz. Journal of Food Technology*. 2022; 25: e2021051. doi: 10.1590/1981-6723.05121.
- (27) Iborra-Bernad C, Tárrega A, García-Segovia P, Martínez-Monzó J. Comparison of Vacuum Treatments and Traditional Cooking Using Instrumental and Sensory Analysis. *Food Anal Methods*. 2013; 7(2): 400-8. doi: 10.1007/s12161-013-9638-0
- (28) Mielby LH, Kildegaard H, Gabrielsen G, Edelenbos M, Thybo AK. Adolescent and adult visual preferences for pictures of fruit and vegetable mixes – Effect of complexity. *Food Qual Prefer*. 2012; 26(2): 188-95. doi: 10.1016/j.foodqual.2012.04.014.
- (29) Santagiuliana M, Bhaskaran V, Scholten E, Piqueras Fiszman B, Stieger M. Don't judge new foods by their appearance! How visual and oral sensory cues affect sensory perception and liking of novel, heterogeneous foods. *Food Qual Prefer*. 2019; 77: 64-77. doi: 10.1016/j.foodqual.2019.05.005.
- (30) Andrés-Bello A, Barreto-Palacios V, García-Segovia P. Effect of pH on Color and Texture of Food Products. *Food Engineering Reviews*. 2013; 5: 158-70. doi: 10.1007/s12393-013-9067-2.
- (31) Tansey F, Gormley R, Butler F. The effect of freezing compared with chilling on selected physico-chemical and sensory properties of sous vide cooked carrots. *Innov Food Sci Emerg Technol*. 2010; 11(1): 137-45. doi: 10.1016/j.ifset.2009.11.001.