

# Record of partial albino in the scalloped hammerhead *Sphyrna lewini* (Carcharhiniformes, Sphyrnidae) in the Mexican Pacific Ocean

Registro de albinismo parcial en el martillo común *Sphyrna lewini* (Carcharhiniformes, Sphyrnidae) en el océano Pacífico mexicano

Jesús Leonardo Hernández-Corona<sup>1</sup>, Armando T. Wakida-Kusunoki<sup>2\*</sup> and Vicente Anislado-Tolentino<sup>3</sup>

<sup>1</sup>Estación de Investigación Acuícola y Pesquera (ESIAP-Pto. Chiapas), Instituto Nacional de Pesca y Acuicultura, calle Dos Norte S/N, Parque Industrial Francisco I. Madero, Puerto Chiapas 30830, Tapachula, Chiapas, México

<sup>2</sup>Centro Regional de Investigación Acuícola y Pesquera de Yucatán, Instituto Nacional de Pesca y Acuicultura, Boulevard del Pescador s/n, esquina Antigua Carretera a Chelem 97320, Progreso, Yucatán, México

<sup>3</sup>Grupo de Investigadores Libres Sphyrna (GILS), Boulevard del Cimatario 439, Constelación, 76087 Santiago de Querétaro, Querétaro, México

\*Corresponding author: [armando.wakida@inapesca.gob.mx](mailto:armando.wakida@inapesca.gob.mx)

**Abstract.**- This study reports the capture of a partial albino scalloped hammerhead *Sphyrna lewini* in Puerto Madero, Chiapas, Mexico. The specimen was a neonate female that measured 491 mm total length and had a total weight of 800 g. This is the first report of an albino scalloped hammerhead *Sphyrna lewini* in the Pacific Ocean and the second in the world. The presence of albinism could be the result of multifactorial events, although anthropogenic stressors such as excessive fishing pressure and marine pollution might be involved.

**Key words:** Elasmobranch, abnormalities, Gulf of Tehuantepec, Mexico

## INTRODUCTION

Coloration of each living being plays a role in basic functions such as crypsis and aposematism and even becomes an indicator of physiological and ethological functions such as sexual selection during the mating season (Protas & Pastel 2008, Hubbard *et al.* 2010). Loss of color has been suggested as one of the disadvantages that wild animals present due to the loss of crypsis, resulting in increased predation and diminished hunting abilities (Sandoval-Castillo *et al.* 2006).

Albinism is a genetically inherited condition in which melanin is absent or nonfunctional (Reum *et al.* 2008), caused by an autosomal recessive gene that affects normal pigmentation in animals (Oliveira & Foresti 1996). There are two types of albinism: a) total albinism is total absence of melanin, it involves the entire body and is characterized by the whitish body and red eyes (Sazima & Pombal 1986), and b) partial (or leucistic) albinism is phenotypically characterized by the absence of melanin in affected parts of the body or reduction of melanin in the entire body or parts of it (Lutz 2001). Albinism has been reported in at least twenty-three shark and fifteen ray species (Wakida-Kusunoki 2015).

The scalloped hammerhead, *Sphyrna lewini* (Griffith & Smith, 1834) is considered a warm-water coastal-pelagic species (Compagno 1984) able to tolerate wide temperature ranges (23-31 °C), low salinity values including freshwater, depths up to 1,000 m, and hypoxic conditions (Castro-Aguirre *et al.* 1999, Shane 2001, Jorgensen *et al.* 2009, Hoyos-Padilla *et al.* 2014, Moore & Gates 2015, Spaet *et al.* 2017, Iqbal *et al.* 2020). It has a circumtropical distribution (Castro 1983), which in the Eastern Pacific Ocean includes southern California, from the Gulf of California to Panama, Ecuador, and northern Peru (Compagno 1984). It can be found near beaches, river deltas, estuaries, and even enters rivers (Compagno 1984, Castro-Aguirre *et al.* 1999, Iqbal *et al.* 2020). Neonates and juveniles are found in shallow water off the coast. They are viviparous placental sharks with a gestation period from 10 months to one year. Neonates are born in breeding areas that are typically shallow protected waters such as bays, inlets, or shoals. In the Mexican Pacific (MP), the reported breeding areas are found at the mouth of the Gulf of California, and on the coast of Jalisco, Michoacán, Oaxaca, and Chiapas (Campuzano-Caballero 2002, Anislado-Tolentino & Robinson-Mendoza 2001, Torres-Huerta *et al.* 2008, Bejarano-Álvarez *et al.* 2011, Alejo-Plata *et al.* 2018,



Anislado-Tolentino 2018, Corgos & Rosende-Pereiro 2022). In the MP, three genetically divergent populations have been described: the North Pacific (Baja California and Baja California Sur), the Central Pacific (Nayarit-Michoacán), and the South Pacific (Oaxaca-Chiapas). Gene flow depends mainly on males, as females have more pronounced philopatry, a trait that helps them conserve energy during gestation and is also one of the factors that shape population structure (Anislado-Tolentino & Robinson-Mendoza 2001, Castillo-Olguín *et al.* 2012).

The scalloped hammerhead is one of the most important species in the catches of the artisanal fleet of different regions of the Mexican Pacific (Furlong-Estrada *et al.* 2015, Coiraton *et al.* 2017). In the Gulf of Tehuantepec, *S. lewini* is the second most captured species with 31% of the total catch. 65% of *S. lewini* catch are obtained from gillnetting for targeted finfish (Morales-Pacheco *et al.* 2016). This zone has been suggested as a birth, breeding, and nursery area for *S. lewini* (Soriano-Velásquez *et al.* 2001, Alejo-Plata *et al.* 2006).

In the last decade, the significant decline (by almost 20%) in scalloped hammerhead catch has been taken as an indirect indicator of the decline in their populations. It is also an international warning sign given their categorization as endangered species since 2010 (CITES 2022)<sup>1</sup>, and critically endangered species in 2018 by IUCN (Rigby *et al.* 2019). In Mexico the protection work on this species began in 1996 and there are currently laws and regulations that regulate shark fishing (DOF 2004<sup>2</sup>, 2007<sup>3</sup>, 2018<sup>4</sup>). However, the recovery and resilience of hammerhead populations have not yet been assessed, and populations may still be stressed by fishing and habitat loss.

This note describes the first report of partial albinism in scalloped hammerhead shark *S. lewini* in the Pacific Ocean.

## MATERIALS AND METHODS

An albino specimen of scalloped hammerhead *Sphyrna lewini* was found during routine shark sampling in artisanal fishing camps. The specimen was caught on June 26, 2018, at a depth of 25 m, 12.5 km off the coast of Puerto Madero, Chiapas (14°40.059'N; 92°31.677'W), using a drifting gillnets mesh with a stretched size of 3.5 inches. The specimen was collected and transported to Estación de Investigación Acuícola y Pesquera (ESIAP-Pto. Chiapas) of Instituto Nacional de Pesca y Acuicultura (INAPESCA), where it was identified using the criteria described by Compagno (1984) and McEachran & Fechhelm (1998). Morphometric data of the albino specimen were obtained. These characteristics were measured with a measuring tape, and total wet weight (W) was measured with an Ohaus® Scout® Pro 4000 (±0.1 g) analytical balance. The neonate state was estimated using the proposal of Duncan & Holland (2006), which is based on characteristics of the healing of the umbilical wound. The specimen was deposited in the fish collection of College of La Frontera Sur, Unidad San Cristóbal de las Casas (ECOSC), under the catalog number ECOSC 14763.

<sup>1</sup>CITES. 2022. History of CITES listing of sharks (Elasmobranchii). Convention on International Trade in Endangered Species of Wild Fauna and Flora. <<https://cites.org/eng/prog/shark/history.php>>

<sup>2</sup>DOF. 2004. Decreto por el que se reforman, adicionan y derogan diversas disposiciones del Reglamento de la Ley de Pesca. 28 de enero del 2004. Diario Oficial de la Federación, Secretaría de Gobernación, Ciudad de México. <[https://dof.gob.mx/nota\\_detalle.php?codigo=677049&fecha=28/01/2004#gsc.tab=0](https://dof.gob.mx/nota_detalle.php?codigo=677049&fecha=28/01/2004#gsc.tab=0)>

<sup>3</sup>DOF. 2004. Decreto por el que se reforman, adicionan y derogan diversas disposiciones del Reglamento de la Ley de Pesca. 28 de enero del 2004. Diario Oficial de la Federación, Secretaría de Gobernación, Ciudad de México. <[https://dof.gob.mx/nota\\_detalle.php?codigo=677049&fecha=28/01/2004#gsc.tab=0](https://dof.gob.mx/nota_detalle.php?codigo=677049&fecha=28/01/2004#gsc.tab=0)>

<sup>4</sup>DOF. 2018. Decreto por el que se adicionan diversas disposiciones de la Ley General de Pesca y Acuicultura Sustentables. 24 de abril del 2018. Diario Oficial de la Federación, Secretaría de Gobernación, Ciudad de México. <[https://dof.gob.mx/nota\\_detalle.php?codigo=5520483&fecha=24/04/2018#gsc.tab=0](https://dof.gob.mx/nota_detalle.php?codigo=5520483&fecha=24/04/2018#gsc.tab=0)>

## RESULTS AND DISCUSSION

The material examined corresponded to a neonate female of *Sphyrna lewini* measuring 491 mm total length and weighing 800 g. Identification of the species was made according to the following observations: head is moderately expanded, width about 30% of total body length. Its anterior margin has shallow undulations that form three lobes, naris groove does not extend to the middle of the anterior margin of the head. Teeth are triangular, wavy base, smooth edges. First dorsal fin straight, not sloping backwards, its free margin never reaching the origin of the pelvic fins, second dorsal fin shorter than anal, the base of anal fin longer than the base of second dorsal, caudal peduncle with a crescent-shaped precaudal pit on the dorsal side. Main morphometric measurements and counts are presented in Table 1. Specimen was considered to have partial albinism due to the tip of the lower lobe of the caudal fin having a small black spot; all the remaining skin was white and eyes were red (Fig. 1a). In pigmented specimens, as observed in the comparison with a 600 mm TL female of approximately three months old, the upper part of the body normally varies between olive to dark gray, white ventrally, with ventral tips of the pectoral fin dark gray to black (Fig. 1b).


According to Gilbert (1967) and Ebert *et al.* (2021), *S. lewini* neonates present a light gray coloration on the back and whitish on the belly with dark marks on the tips of the fins. As they grow, they acquire a darker coloration and lose the marks on the fins. In this work, the analyzed specimen presents a whitish coloration both on the back and on the belly, with the addition that the eyes are red. Since the lower caudal lobule has a black spot it can be considered to have partial albinism (Sazima & Pombal 1986).

Currently, 31 species of sharks with albinism have been reported around the world (Table 2), however, only one report for *S. lewini* (McKenzie 1970) was known before this work. Of the five reported cases of albinism on the Mexican coast, two were elasmobranchii found in the Gulf of Tehuantepec: *Carcharhinus limbatus* (Sancho-Vazquez *et al.* 2015) and *Urotrygon nana* (Anislado-Tolentino *et al.* 2016).

Albinism and alterations in coloration have been related to exposure to heavy metals (Oliveira & Foresti 1996, Irigoyen-Arredondo *et al.* 2017), genetic alteration in melanin production (Bejarano-Álvarez & Galván-Magaña 2013), inbreeding within an isolated population (Sanabria *et al.* 2010), environmental stress, and heavy fishing pressure (Gervais *et al.* 2016, Bruckner & Coward 2018).

**Table 1. Morphometric measurements (mm) of the reported *Sphyrna lewini* partial albino female (ECOSC 14763) /**  
Medidas morfológicas (mm) del espécimen hembra de *Sphyrna lewini* reportado con albinismo parcial (ECOSC 14763)

Measurements	(mm)	Measurements	(mm)
Total length	491	Anal base	30
Fork length	371	Anal length	45
Pre-caudal length	342	Anal height	20
Inter-dorsal space	113	Ventral base	39
Pre-first dorsal length	145	Ventral height	47
Pre-second dorsal length	301	Caudal length	148
Snout length	40	Terminal caudal margin	25
Head length	131	Sub-terminal caudal margin	16
First dorsal base	50	Space between pectoral and pelvic fins	91
First dorsal length	65	Space between the vent and caudal length	98
First dorsal height	62	Eye length	15
Second dorsal base	19	Eye height	15
Second dorsal length	44	Intergill length	33
Second dorsal height	13	Pre-oral length	37
Pectoral base	28	Pre-orbital length	31
Pectoral length	47	Space between pelvic and anal fins	39
Pectoral height	52	Head width	164
Pelvic base	30		
Pelvic length	41		
Pelvic height	24		



**Figure 1.** *Sphyrna lewini*. A) Head dorsal view of the reported partial albino female with a close-up of the right eye; B) Body dorsal view with a close-up of tail lower lobe showing a black spot; C) Pectoral area ventral view, where fresh umbilical opening is observed; D) Head dorsal view of a normal female; E) Body dorsal view of a normal female; F) Pectoral area ventral view of a normal female with black spotted fin tips and healed umbilical opening. Partial albino specimen photo by JL Hernández-Corona and normal specimen photo by V Anislado-Tolentino / *Sphyrna lewini*. A) Vista dorsal de la cabeza de la hembra albina parcial con un acercamiento al ojo derecho; B) Vista dorsal del cuerpo con un acercamiento al lóbulo inferior de la aleta caudal que muestra una mancha oscura; C) Vista ventral del área pectoral de la hembra albina donde la abertura umbilical se nota fresca; D) Vista dorsal de la cabeza de una hembra con coloración normal; E) Vista dorsal del cuerpo de una hembra con coloración normal; F) Vista ventral del área pectoral de una hembra con coloración normal, donde se notan las manchas oscuras en las puntas de las aleta y a la abertura umbilical cicatrizada. Foto de ejemplar con albinismo parcial por JL Hernández-Corona y foto del ejemplar normal por V Anislado-Tolentino

Although this is the second reported case of albinism for *S. lewini* worldwide, inbreeding must be ruled out as a cause because this is a migratory species, exchange of genetic material is unrestricted, and therefore this factor does not have weight. However, it must be considered that in the northern Pacific states, up to 95% of wetlands have been lost in the last decade, mostly affecting Baja California, Baja California Sur, Sonora and Sinaloa; central Pacific states have lost up to 50% (Landgrave & Moreno-Casasola 2012). This suggests a correlation between the decreasing populations of large coastal predators and the loss and deterioration of the coastal habitat where the breeding areas are found, being detrimental to gene transmission in the three scalloped hammerhead populations from MP. Additionally, organochlorines and polychlorinated benzyl are persistent organic pollutant compounds (OPCs) that are known to cause teratogens and

carcinogens (Martínez-Villa *et al.* 2014), and have been shown to cause joint albinism with cyclopia in sea turtles (Bárcenas & Maldonado 2009). Chiapas has been reported as one of the states with the greatest problems derived from OPCs due to the presentation of persistent organic pesticides (POPs) from agriculture and the use of pesticides against malaria (Albert 1996, Ruiz-Suárez *et al.* 2014). In the coastal areas of Chiapas, a large number of POPs have been detected in sediments (Botello *et al.* 2002), as well as in the flesh of fish for human consumption (Herrera *et al.* 2013). For this reason, the high POP levels (DDT-derived heptachlorides) found by Martínez-Villa *et al.* (2014) for *S. lewini* in the Gulf of Tehuantepec might suggest a correlation with reported albinism, however, as Escobar-Sánchez *et al.* (2014) mention, this and other abiotic factors can be ruled out since the presented specimen is the only one of the litter affected.

**Table 2. Reported cases of albinism and partial albinism in shark species. Species in the order suggested by Compagno (2005) and Cornejo *et al.* (2015) / Casos reportados de albinismo y albinismo parcial en especies de tiburones. Especies ordenadas de acuerdo a la propuesta de Compagno (2005) y Cornejo *et al.* (2015)**

Family	Species	Albinism	Capture site	Reference
Hexanchidae	<i>Notorynchus cepedianus</i>	Partial	California, USA	Herald (1953)
	<i>Notorynchus cepedianus</i>	Partial	California, USA	Ebert (1975)
Squalidae	<i>Squalus acanthias</i>	Partial	Georgia, USA	Froiland (1975)
	<i>Squalus acanthias</i>	Partial	Nova Scotia, Canada	Coad & Gilhen (2002)
	<i>Squalus megalops</i>	Total	Black Sea, Romania	Sanda & De Maddalena (2003)
Etmopteridae	<i>Etmopterus lucifer</i>	Total	New Zealand	Finucci (2020)
Somniosidae	<i>Centroscymnus coelolepis</i>	Partial	NE Atlantic	Deynat (2003)
Dalatiidae	<i>Dalatis licha</i>	Partial	Italy	Bottaro <i>et al.</i> (2005)
Squatinae	<i>Squatina californica</i>	Total	Baja California Sur, Mexico	Escobar-Sánchez <i>et al.</i> (2014)
Hemiscylliidae	<i>Chiloscyllium plagiosum</i>	Total	USA	Clark (2002)
Orectolobidae	<i>Orectolobus japonicus</i>	Total	Japan	Iwamasa & Okano (1982)
Ginglymostomatidae	<i>Nebrius ferrugineus</i>	Total	Japan	Taniuchi & Yanagisawa (1987)
Stegostomatidae	<i>Stegostoma fasciatum</i>	Total	Indian Ocean	Nayaka (1973)
Odontaspidae	<i>Odontaspis ferox</i>	Partial	New South Wales Australia	Fergusson <i>et al.</i> (2008)
Cetorhinidae	<i>Cetorhinus maximus</i>	NA	Norway	Froiland (1975)
Lamnidae	<i>Carcharodon carcharias</i>	Partial	South Africa	Smale & Heemstra (1997)
	<i>Lamna nasus</i>	Total	Norway	Froiland (1975)
Scyliorhinidae	<i>Cephaloscyllium ventriosum</i>	Total	Baja California Sur, Mexico	Becerril-García <i>et al.</i> (2017)
	<i>Scyliorhinus canicula</i>	Total	Ireland	Hoare (2009)
Triakidae	<i>Galeorhinus galeus</i>	Total	English Channel	Deynat (2003)
	<i>Hemitriakis japonica</i>	Partial	Japan	Furuta (1985)
	<i>Mustelus californicus</i>	NA	Monterey Bay, USA	Herald <i>et al.</i> (1960)
	<i>Mustelus californicus</i>	Total	California, USA	Talent (1973)
	<i>Mustelus californicus</i>	Total	California, USA	Cohen (1973)
	<i>Mustelus schmitti</i>	Total	Brazil	Ferreira-Teixeira & Góes de Araújo (2002)
Carcharhinidae	<i>Triakis semifasciata</i>	Total	California, USA	Follet (1976)
	<i>Carcharhinus amboinensis</i>	Partial	Australia	McKay & Beinssen (1988)
	<i>Carcharhinus isodon</i>	Partial	Alabama, USA	Jones <i>et al.</i> (2006)
	<i>Carcharhinus limbatus</i>	Partial	Chiapas, Mexico	Sancho-Vasquez- <i>et al.</i> (2015)
	<i>Carcharhinus melanopterus</i>	Total	India	Manojkumar (2011)
	<i>Carcharhinus melanopterus</i>	Partial	Maldives Islands	Bruckner & Coward (2018)
	<i>Carcharhinus obscurus</i>	Total	Baja California Sur, Mexico	Bejarano-Álvarez & Galván-Magaña (2013)
	<i>Carcharhinus plumbeus</i>	Partial	Gulf of Gabés, Tunisia	Saïdi <i>et al.</i> (2006)
	<i>Galeocerdo cuvier</i>	Total	Baja California Sur, Mexico	Sandoval-Castillo <i>et al.</i> (2006)
	<i>Galeocerdo cuvier</i>	Total	Gulf of Mexico	Rider <i>et al.</i> (2002)
Sphyrnidae	<i>Scoliodon laticaudus</i>	Partial	Arabian Sea	Veena <i>et al.</i> (2011)
	<i>Sphyrna lewini</i>	Total	Georgia, USA	McKenzie (1970)
	<i>Sphyrna lewini</i>	Partial	Chiapas, Mexico	This work

NA= not available

Cárdenas & Franco (2021) suggest that in addition to inbreeding, hybridization and nutritional deficiencies can also cause albinism. Rodríguez-Ruíz *et al.* (2015) report that albinism in city birds is caused by anthropogenic stressors that cause congenital malformations in embryonic development. Irigoyen-Arredondo *et al.* (2018), postulate that the anthropogenic stressors affecting marine organisms can range from fishing, extensive navigation on commercial and tourist routes, industrial waste that is dumped on the coast, and loss of habitat due to human population growth on the coast, causing a failure in the genetic interconnectivity between populations. Despite the fact that for decades hammerhead fishing on the Chiapas coast have been reported as intensive (Soriano-Velásquez *et al.* 2001, Alejo-Plata *et al.* 2006), and the introduction of regulatory fishing laws in 2007 (DOF 2007)<sup>3</sup>, there has been no evaluation of population recovery. The absence of other reports of albinism in *S. lewini* within the area or in other localities with similar levels of exploitation prevents the generalization of anthropogenic stress as the cause of albinism in the specimen. On the other hand, and in agreement with Irigoyen-Arredondo *et al.* (2018), the malformation in the *S. lewini* specimen is of multifactorial origin, which at macroscopic level will remain only as assumption, and may be clarified through genetic, nutritional and biochemical analysis of the specimens in the future.

## ACKNOWLEDGMENTS

The authors would like to thank to José Francisco Valle Estrada and Alejandro Valle Centeno, administrator and owner of the “Manuel Valle” fish reception center, for donating the specimen.

## LITERATURE CITED

- Albert LA. 1996.** Persistent pesticides in Mexico. Reviews of Environmental Contamination and Toxicology 147: 1-44. <doi:10.1007/978-1-4612-4058-7\_1>
- Alejo-Plata MC, S Ramos & JL Cruz. 2006.** La pesquería artesanal del tiburón en Salina Cruz, Oaxaca, México. Ciencia y Mar México 30: 37-51.
- Alejo-Plata MC, MAhumada-Sempool, GC Cerdaneres & J Gómez-Márquez. 2018.** Population structure and reproductive biology of the scalloped hammerhead *Sphyrna lewini* (Carcharhiniformes: Sphyrnidae) caught in artisanal fisheries of Oaxaca, Mexico. Hidrobiológica 28(3): 265-275. <https://doi.org/10.24275/uam/izt/dcbs/hidro/2018v28n3/alejo>
- Anislado-Tolentino V. 2018.** Aspectos reproductivos y área de crianza del tiburón martillo *Sphyrna lewini*, en la costa de Michoacán, México. Ciencia Pesquera 26(2): 13-28.
- Anislado-Tolentino V & C Robinson-Mendoza. 2001.** Age and growth for the scalloped hammerhead shark, *Sphyrna lewini* (Griffith and Smith, 1834) along the Central Pacific coast of Mexico. Ciencias Marinas 27(4): 501-520. <https://doi.org/10.7773/cm.v27i4.503>
- Anislado-Tolentino V, MG Medina-Lerma & LF Del Moral-Flores. 2016.** Primer registro de albinismo en la raya redonda enana, *Urotrygon nana* (Urotrygonidae: Elasmobranchii), en el Pacífico oriental. Revista de Ciencias Marinas y Costeras 8(2): 93-100. <https://doi.org/10.15359/revmar.8-2.6>
- Bárceñas IA & G Maldonado. 2009.** Malformaciones en embriones y neonatos de tortuga golfina (*Lepidochelys olivacea*) en Nuevo Vallarta, Nayarit, México. Veterinaria México 40(4): 371-380.
- Becerril-García EE, E Tamburin, R González-Armas & F Galván-Magaña. 2017.** First record of albinism in the swell shark, *Cephaloscyllium ventriosum* (Elasmobranchii: Carcharhiniformes: Sphyrnidae). Acta Ichthyologica et Piscatoria 47(2): 201-204. <https://doi.org/10.3750/AIEP/02175>
- Bejarano-Álvarez M, F Galván-Magaña & RI Ochoa-Báez. 2011.** Reproductive biology of the scalloped hammerhead shark *Sphyrna lewini* (Chondrichthyes: Sphyrnidae) off south-west Mexico. Aqua: International Journal of Ichthyology 17(1): 11-23.
- Bejarano-Álvarez OM & F Galván-Magaña. 2013.** First report of an embryonic dusky shark (*Carcharhinus obscurus*) with cyclopia and other abnormalities. Marine Biodiversity Records 6. <https://doi.org/10.1017/S1755267212001236>
- Botello AV, L Rueda-Quintana, G Díaz-González & A Toledo. 2002.** Persistent Organochlorine Pesticides (POPs) in Coastal Lagoons of the Subtropical Mexican Pacific. Bulletin of Environmental Contamination and Toxicology 64(3): 390-397. <https://doi.org/10.1007/s001280000013>
- Bottaro M, S Ferrando, L Gallus, L Girosi & M Vacchi. 2005.** First record of albinism in the deep-water shark *Dalatias licha*. Marine Biodiversity Records 1: 1-4. <https://doi.org/10.1017/S1755267205001156>
- Bruckner AW & G Coward. 2018.** The unusual occurrence of abnormal skin pigmentation in blacktip reef sharks (*Carcharhinus melanopterus*). Coral Reefs 37: 389. <https://doi.org/10.1007/s00338-018-1663-4>
- Campuzano-Caballero JC. 2002.** Biología y pesquería del tiburón martillo *Sphyrna lewini* (Griffith y Smith, 1834), en Puerto Madero, Chiapas, Estados Unidos Mexicanos. Tesis de Licenciatura en Biología, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad de México, 196 pp. <http://132.248.9.195/ppt2002/0307482/0307482.pdf>
- Cárdenas G & JC Franco. 2021.** New records of pigmentary abnormalities in two species of birds in Cali, Colombia. Biota Colombiana 22(2): 147-154.
- Castillo-Olguín E, M Uribe-Alcoce & P Díaz-Jaimes. 2012.** Assessment of the population genetic structure of *Sphyrna lewini* to identify conservation units in the Mexican Pacific. Ciencias Marinas 38(4): 635-652. <https://doi.org/10.7773/cm.v38i4.2110>
- Castro JI. 1983.** The sharks of North American waters, 194 pp. Texas A&M University Press, College Station.
- Castro-Aguirre JL, HS Espinosa-Pérez & JJ Schmitter-Soto. 1999.** Ictiofauna estuarino-lagunar y vicaria de México, 711 pp. Limusa, Ciudad de México.

- Clark S. 2002.** First report of albinism in the white-spotted bamboo shark, *Chiloscyllium plagiosum* (Orectolobiformes: Hemiscyllidae), with a review of reported color aberrations in elasmobranchs. *Zoo Biology* 21: 519-524. <<https://doi.org/10.1002/zoo.10068>>
- Coad B & J Gilhen. 2002.** A rare leucistic spiny dogfish, *Squalus acanthias*, from the Bay of Fundy, Nova Scotia. *The Canadian Field-Naturalist* 116(1): 120-121.
- Cohen JL. 1973.** An albino grey smoothhound *Mustelus californicus*, Gill. *California Fish and Game* 59: 210-211.
- Coiratón C, F Amezcua & IH Salgado-Ugarte. 2017.** Estructura de longitudes de las capturas del tiburón martillo común (*Sphyrna lewini*) en el Pacífico mexicano. *Ciencia Pesquera* 25(1): 27-40.
- Compagno LJV. 1984.** FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2. FAO Fisheries Synopsis 125 4(2): 251-655.
- Compagno LJV. 2005.** Appendix 1: Global checklist of living chondrichthyan fishes. In: Fowler SL, RD Cavanagh, M Camhi, GH Burgess, GM Cailliet, SV Fordham, CA Simpfendorfer & JA Musick (eds). *Sharks, rays and chimaeras: The status of the Chondrichthyan fishes. Status survey*, pp. 401-423. IUCN/ SSC Shark Specialist Group. IUCN, Gland and Cambridge.
- Corgos A & A Rosende-Pereiro. 2022.** Nursery habitat use patterns of the scalloped hammerhead shark, *Sphyrna lewini*, in coastal areas of the central Mexican Pacific. *Journal of Fish Biology* 100(1): 117-133. <<https://doi.org/10.1111/jfb.14925>>
- Cornejo R, X Vélez-Zuazo, A González-Pestana, CJ Kouri & G Mucientes. 2015.** An updated checklist of Chondrichthyes from the southeast Pacific off Peru. *Check List* 11(6). <<http://dx.doi.org/10.15560/11.6.1809>>
- Deynat PP. 2003.** Albinisme partiel chez le pailona commun, *Centroscymnus coelolepis* (Elasmobranchii, Somniosidae). *Cybium* 27: 233-236. <<https://doi.org/10.26028/cybium/2004-273-006>>
- Duncan KM & KN Holland. 2006.** Habitat use, growth rates and dispersal patterns of juvenile scalloped hammerhead sharks *Sphyrna lewini* in a nursery habitat. *Marine Ecology Progress Series* 312: 211-221. <<https://doi.org/10.3354/meps312211>>
- Ebert DA. 1975.** Color variation in the sevengill shark, *Notorynchus maculatus* Ayres along the California Coast. *California Fish and Game* 71: 53-59.
- Ebert DA, S Fowler & M Dando. 2021.** *Sharks of the world - A fully illustrated guide*, 607 pp. Wild Nature Press, Princeton.
- Escobar-Sánchez O, XG Moreno-Sánchez, CA Aguilar-Cruz & LA Abitia-Cárdenas. 2014.** First case of synophthalmia and albinism in the Pacific angel shark *Squatina californica*. *Journal of Fish Biology* 85(2): 494-501. <<https://doi.org/10.1111/jfb.12412>>
- Fergusson IK, KJ Graham & LJV Compagno. 2008.** Distribution, abundance and biology of the smalltooth sandtiger shark, *Odontaspis ferox* (Risso 1810) (Lamniformes: Odontaspidae). *Environmental Biology of Fishes* 81: 207-228.
- Ferreira-Teixeira S & ML Góes de Araújo. 2002.** First record of albinism in the Smooth Dogfish *Mustelus schmitti* Springer, 1939 (Carcharhiniformes- Triakidae) from Southern Brazil. *Brazilian Archives of Biology and Technology* 45: 241-243. <<https://doi.org/10.1590/S1516-89132002000200017>>
- Finucci B. 2020.** First record of albinism in the lanternshark family, Etmopteridae. *Journal of Fish Biology* 96: 1512-1515. <<https://doi.org/10.1111/jfb.14329>>
- Follet WI. 1976.** First record of albinism in the leopard *Triakis semifasciata* Girard. *California Fish and Game* 62: 163-164.
- Frøiland Ø. 1975.** Albinisme hos hai. *Fauna, Oslo* 28: 170-173.
- Furlong-Estrada E, J Tovar-Ávila, JC Pérez-Jiménez & E Ríos-Jara. 2015.** Resilience of *Sphyrna lewini*, *Rhizoprionodon longurio*, and *Carcharhinus falciformis* at the entrance to the Gulf of California after three decades of exploitation. *Ciencias Marinas* 41(1): 49-63. <<http://dx.doi.org/10.7773/cm.v41i1.2442>>
- Furuta M. 1985.** Sharks of Mie Prefecture. Report of the Mie Zoological Society 7: 10-13. [In Japanese]
- Gervais C, J Mourier & J Rummer. 2016.** Developing in warm water: irregular colouration and patterns of a neonate elasmobranch. *Marine Biodiversity* 46: 743-744. <<http://dx.doi.org/10.1007/s12526-015-0429-2>>
- Gilbert CR. 1967.** A revision of the Hammerhead sharks (family Sphyrnidae). *Proceedings of the United States National Museum* 119(3539): 1-88.
- Herald ES. 1953.** The 1952 shark derbies at Elkhorn Slough, Monterey Bay, California. *California Fish Game* 39: 237-243.
- Herald ES, W Schneebeil, N Green & K Innes. 1960.** Catch records for seventeen shark derbies held at Elkhorn Slough, Monterey Bay. *California Fish and Game* 46: 59-67.
- Herrera PC, G Franco, G Bermúdez & HB Schlottfeldt. 2013.** Niveles de DDT y metabolitos (DDE y DDD) en peces de consumo humano en una comunidad endémica de paludismo en Chiapas, México. *Higiene y Sanidad Ambiental* 13(4): 1080-1085.
- Hoare D. 2009.** First record of albinism in the Lesser-spotted Dogfish (*Scyliorhinus canicula* (L., 1758)) (Chondrichthyes: Scyliorhinidae) and a review of albinism in other elasmobranchs. *The Irish Naturalist's Journal* 30: 115-118.
- Hoyos-Padilla EM, JT Ketchum, AP Klimley & F Galván-Magaña. 2014.** Ontogenetic migration of a female scalloped hammerhead shark *Sphyrna lewini* in the Gulf of California. *Animal Biotelemetry* 2: 17. <<https://doi.org/10.1186/2050-3385-2-17>>
- Hubbard JK, JAC Uy, ME Hauber, HE Hoekstra & RJ Safran. 2010.** Vertebrate pigmentation: from underlying genes to adaptive function. *Trends in Ecology and Evolution* 26: 231-239. <<http://dx.doi.org/10.1016/j.tig.2010.02.002>>
- Iqbal M, D Setiawan, A Setiawan & I Yustian. 2020.** First record of scalloped hammerhead shark *Sphyrna lewini* (Carcharhiniformes: Sphyrnidae) in freshwater habitat. *Ecological Montenegrina* 30: 113-118. <<http://dx.doi.org/10.37828/em2020.30.10>>

- Irigoyen-Arredondo MS, O Escobar-Sánchez, LA Abitia-Cárdenas, XG Moreno-Sánchez & DS Palacios-Salgado. 2017.** Incidence of xanthism in the leopard grouper *Mycteroperca rosacea* (Perciformes: Serranidae) in the Gulf of California. *Marine Biodiversity* 48: 2255-2258. <<http://dx.doi.org/10.1007/s12526-017-0753-9>>
- Iwamasa H & S Okano. 1982.** Albino embryo of *Orectolobus japonicus*. Report of Japanese Group for Elasmobranch Studies 9: 56-57.
- Jones L, M Shipp & L Robert. 2006.** Oculocutaneous albinism in a finetooth shark, *Carcharhinus isodon*, from Mobile Bay, Alabama. *Gulf of Mexico Science* 24(1/2): 81-82. <<http://dx.doi.org/10.18785/goms.2401.10>>
- Jorgensen SJ, AP Klimley & AF Muhlia-Melo. 2009.** Scalloped hammerhead shark *Sphyrna lewini*, utilizes deep-water, hypoxic zone in the Gulf of California. *Journal of Fish Biology* 74: 1682-1687. <<https://doi.org/10.1111/j.1095-8649.2009.02230.x>>
- Landgrave R & P Moreno-Casasola. 2012.** Evaluación cuantitativa de la pérdida de humedales en México. *Investigación Ambiental* 4: 19-35.
- Lutz GC. 2001.** Practical genetics for aquaculture, 235 pp. Blackwell Science, Oxford.
- Manojkumar PP. 2011.** First record of albinism in the blacktip reef shark *Carcharhinus melanopterus* from Malabar Coastal. *Marine Fisheries Information Service T&E Series* 208: 36. <<http://eprints.cmfri.org.in/8907/1/208-26.pdf>>
- Martínez-Villa G, M Betancourt-Lozano, G Aguilar-Zárate, J Ruelas-Inzunza, V Anislado-Tolentino, G Cerdaneres & G González. 2014.** Contenido de plaguicidas organoclorados en varios peces depredadores de la costa de Oaxaca y evaluación del riesgo de exposición por consumo en la salud humana. En: Botello AV, F Páez-Osuna, L Méndez-Rodríguez, M Betancourt-Lozano, S Álvarez-Borrego & R Lara-Lara (eds). *Pacífico mexicano. Contaminación e impacto ambiental: diagnóstico y tendencias*, pp. 169-209. UAC, UNAM-ICMYL, CIAD-MAZATLAN, CIBNOR, CICESE, Ciudad de México.
- McEachran J & JD Fechhelm. 1998.** Fishes of the Gulf of Mexico, Vol. 1: Myxiniiformes to Gasterosteiformes, 1112 pp. University of Texas Press, Austin.
- McKay RJ & K Beinssen. 1988.** Albinism in the pigeye whaler shark *Carcharhinus amboinensis* Mueller and Henle from Queensland Australia. *Memoirs of the Queensland Museum* 25(2): 463-464.
- McKenzie MD. 1970.** First record of albinism in the Hammerhead Shark, *Sphyrna lewini* (Pisces: Sphyrnidae). *Journal of the Elisha Mitchell Scientific Society* 86(1): 35-37.
- Moore ABM & AR Gates. 2015.** Deep-water observation of scalloped hammerhead *Sphyrna lewini* in the western Indian Ocean off Tanzania. *Marine Biodiversity Records* 8: E91. <[doi:10.1017/S1755267215000627](https://doi.org/10.1017/S1755267215000627)>
- Morales-Pacheco O, JL Hernández-Corona, AA Lizárraga-Rodríguez & AJ Alatorre-Alba. 2016.** Golfo de Tehuantepec. In: Castillo-Géniz L & J Tovar-Ávila (eds). *Tiburones mexicanos de importancia pesquera en la CITES*, pp. 53-55. Instituto Nacional de la Pesca, Ciudad de México.
- Nayaka K. 1973.** An albino zebra shark *Stegostoma fasciatum* from the Indian Ocean, with comments on albinism in elasmobranchs. *Japanese Journal of Ichthyology* 20: 120-122.
- Oliveira C & F Foresti. 1996.** Albinism in the banded knifefish, *Gymnotus carapo*. *Tropical Fish Hobbyist Magazine* 44: 92-96.
- Protas ME & NH Pastel. 2008.** Evolution of coloration patterns. *Annual Review Cell and Developmental Biology* 24: 425-446. <<https://doi.org/10.1146/annurev.cellbio.24.110707.175302>>
- Reum J, CE Paulsen, TW Pietsch & SL Parker-Stetter. 2008.** First record of an albino chimaeriform fish, *Hydrolagus colliei*. *Northwestern Naturalist* 8: 60-62. <[https://doi.org/10.1898/1051-1733\(2008\)89\[60:FROAAC\]2.0.CO;2](https://doi.org/10.1898/1051-1733(2008)89[60:FROAAC]2.0.CO;2)>
- Rider S, M Athorn & G Bailey. 2002.** First record of a white tiger shark *Galeocerdo cuvier*, from the northeastern Gulf of Mexico. *Florida Scientist* 65: 13-15.
- Rigby CL, NK Dulvy, R Barreto, J Carlson, D Fernando, S Fordham, MP Francis, K Herman, RW Jabado, KM Liu, A Marshall, N Pacoureau, E Romanov, RB Sherley & H Winker. 2019.** *Sphyrna lewini*. The IUCN Red List of Threatened Species 2019: e.T39385A2918526. <<https://www.iucnredlist.org/es/species/39385/2918526>>
- Rodríguez-Ruiz ER, I Martínez-Sánchez & J Treviño-Carreón. 2015.** Nuevos registros de aberraciones cromáticas en el plumaje de dos especies de aves en zonas urbanas de Hidalgo y Tamaulipas, México. *Acta Zoológica Mexicana* 31(3): 466-469.
- Ruiz-Suárez LE, RA Castro-Chan, NE Rivero-Pérez, A Trejo-Acevedo, GK Guillén-Navarro, V Geissen & R Bello-Mendoza. 2014.** Levels of organochlorine pesticides in blood plasma from residents of malaria-endemic communities in Chiapas, Mexico. *International Journal of Environmental Research and Public Health* 11(10): 10444-10460. <<http://dx.doi.org/10.3390/ijerph111010444>>
- Saïdi B, MN Bradai, S Marouani, O Guélorget & C Capapé. 2006.** Atypical characteristics of an albino embryo of *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae) from the Gulf of Gabés southern Tunisia, central Mediterranean. *Acta Adriatica* 47: 167-174.
- Sanabria EA, LB Quiroga & A Laspiur. 2010.** First record of partial albinism and scoliosis in *Odontophrynus occidentalis* tadpoles (Anura: Cycloramphidae). *Brazilian Archives of Biology and Technology* 53(3): 641-642. <<https://doi.org/10.1590/S1516-89132010000300019>>
- Sancho-Vazquez E, LF Del Moral-Flores & OU Mendoza-Vargas. 2015.** Albinismo parcial en un embrión de tiburón puntas negras, *Carcharhinus limbatus* (Carcharhinidae: Chondrichthyes) del Pacífico oriental. *Latin American Journal of Aquatic Research* 43: 274-276. <<https://doi.org/10.3856/vol43-issue1-fulltext-25>>
- Sanda R & A De Maddalena. 2003.** Collection of sharks of the National Museum in Prague - Part 1. Complete taxiderms and liquid preservations. *Journal of the National Museum (Prague), Natural History Series* 172: 61-70.



- Sandoval-Castillo J, E Mariano-Melendez & C Villavicencio-Garayzar. 2006.** New records of albinism in two elasmobranchs: the tiger shark *Galeocerdo cuvier* and the giant electric ray *Narcine entemedor*. *Cybium* 30: 191-192. <<https://doi.org/10.26028/cybium/2006-302-014>>
- Sazima I & JP Pombal. 1986.** Um albino de *Rhambdella minuta*, com notas sobre comportamento (Osteichthyes, Pimelodidae). *Revista Brasileira de Biologia* 46: 377-381.
- Shane MA. 2001.** Records of Mexican barracuda, *Sphyrna ensis*, and scalloped hammerhead, *Sphyrna lewini*, from southern California associated with elevated water temperatures. *Bulletin of the Southern California Academy of Sciences* 100(3): 160-166.
- Smale MJ & PC Heemstra. 1997.** First record of albinism in the great white shark, *Carcharodon carcharias* (Linnaeus, 1758). *South African Journal of Science* 93: 243-245.
- Soriano-Velásquez SR, A Solís-Nava, C Ramírez-Santiago, A Cid del Prado-Vera & JL Castillo-Géniz. 2001.** Tiburones del Golfo de Tehuantepec. En: Cisneros-Mata MA & A Díaz de León (eds). *Sustentabilidad y pesca responsable en México 1999-2000*, pp. 211-227. Instituto Nacional de la Pesca, Ciudad de México.
- Spaet JLY, CH Lam, CD Braun & ML Berumen. 2017.** Extensive use of mesopelagic waters by a scalloped hammerhead shark (*Sphyrna lewini*) in the Red Sea. *Animal Biotelemetry* 5: 20. <<https://doi.org/10.1186/s40317-017-0135-x>>
- Talent LG. 1973.** Albinism in embryo grey smoothhound shark, *Mustelus californicus*, from Elkhorn Slough, Monterey Bay, California. *Copeia* 1973(3): 595-597. <<https://doi.org/10.2307/1443129>>
- Taniuchi T & F Yanagisawa. 1987.** Albinism and a lack of second dorsal fin in an adult tawny nurse shark, *Nebrius concolor*, from Japan. *Japanese Journal of Ichthyology* 34: 393-395.
- Torres-Huerta AM, C Villavicencio-Garayzar & D Corro-Espinosa. 2008.** Biología reproductiva de la cornuda común *Sphyrna lewini* Griffith & Smith (Sphyrnidae) en el Golfo de California. *Hidrobiológica* 18(3): 227-238.
- Veena S, S Thomas, S Raje, G Durgekar & N Raveendra. 2011.** Case of leucism in the spadenose shark, *Scoliodon laticaudus* (Müller and Henle, 1838) from Mangalore, Karnataka. *Indian Journal of Fisheries* 58(1): 109-112.
- Wakida-Kutsunoki AT. 2015.** First record of total albinism in southern stingray *Dasyatis americana*. *Revista de Biología Marina y Oceanografía* 50(1): 135-139.

---

Received 15 October 2021

Accepted 24 October 2022