Transanal endoscopic microsurgery: indications and results after 100 cases

P. Palma / S. Frendenberg / S. Samel / S. Post

Objective Transanal endoscopic microsurgery (TEM), a minimally invasive technique has been employed in the excision of benign and well-selected malignant rectal tumours since June 1998. We present a prospective descriptive study and analyse the currently accepted indications.

Patients and methods Over a 4-year period 100 patients underwent TEM for treatment of rectal tumours located between 4 and 18 cm from the anal verge.

Results TEM was performed in 71 cases for adenomas, 20 potentially curative excisions for pre-operative staged low-grade carcinoma, 3 palliative procedures for advance carcinoma, 5 carcinoids and 1 solitary ulcer. The local complication rate included wound breakdown in 7 patients, three of them requiring

ileostomy. Conversion to laparotomy was performed in two patients. Five adenomas recurred and were successfully treated by TEM. Of the cancers, four patients required immediate salvage therapy by means of total mesorectal excision.

Three patients underwent palliative TEM procedures combined with radiotherapy. A single cancer recurrence was treated by means of abdomino-perineal resection after radiotherapy.

Conclusions TEM appears to be an effective method of excising benign tumours and selected T1 carcinomas of the rectum. The superior exposure of tumours higher in the rectum combined with the greater precision of excision make this minimally invasive technique an attractive surgical approach.

Introduction

Colorectal cancer represent the second leading cause of cancer-related mortality in the Western world. It is widely accepted that the adenoma-carcinoma sequence represents the process by which most, if not all, rectal carcinomas arise [1]. The appropriate management of individuals with precursor adenomatous polyps in the

rectum, is removal by means of polypectomy with snare diathermy, or complete local resection depending on the size and morphology. Additionally, local resection is also indicated for selected 'low-risk' rectal carcinomas. However tumour location may limit local removal per se leading to more aggressive posterior approaches or to an anterior resection via laparotomy-laparoscopy.

Palabras clave: Transanal endoscopic microsurgery, rectal adenoma, rectal carcinoma, local excision.

Fecha de recepción: Enero 2004.

Seminario Médico

Año 2004. Volumen 56, N.º 3. Págs. 11-18

Transanal endoscopic microsurgery (TEM) was introduced by Buess *et al.* in 1984 [2] and is currently undergoing widespread evaluation for treatment of both adenomas and selected carcinomas [3-10]. TEM is a minimally invasive method that allows precise resection of tumours located 4-18 cm from the anal verge using an operative rectoscope.

The aim of the present study was to report our single Institution experience. The major parameters of the study are the analysis of the currently accepted indications, as well as practicability, morbidity and recurrence rates of the method since its introduction in our coloproctology unit.

Materials and methods

Patients

Between June 1998 and June 2002, all patients with rectal tumours were examined in our interdisciplinary surgical-endoscopy unit. Following clinical examination, a biopsy was obtained in all patients, and endorectal sonography was performed using a 360-degree endoprobe with an inflatable balloon at a frequency of 7.5 MHz for accurate pre-operative staging (Fig. 1). An abdominal ultrasound and a colonoscopy were also performed in all patients in order to rule out synchronous more proximal pathology.

Figure 1.—Endorectal sonography showing a T1 rectal tumour.

Pre-operative staging resulted in three patient groups: The largest group in our experience had benign rectal tumours not suitable for removal by endoscopic diathermy snare. The second group had proven rectal cancer. The third and smallest group comprised neuroendocrine, carcinoid and miscellaneous tumours.

A total of 100 patients presenting with tumours located 4-18 cm from the anal verge were treated by TEM. No other surgical approach was used or evaluated for local excision in our department since the beginning of the study. Only three patients, presenting high risk features (lymphovascular or muscular invasion, or poor differentiation) were operated using TEM with a palliative intention. The bowel was prepared as for a formal laparotomy by lavage over 5 h with 3-4 litres of polyethylene glycol solution. Antibiotic prophylaxis consisted of a single dose of a cephalosporin plus metronidazole and was given at the time of anaesthetic induction.

Methods

Following the original description by Buess *et al.* [2], we used an operative rectoscope of 40 mm diameter and 120 or 200 mm length, with a sixfold magnified stereoscopic view (Fig. 2). For better visualization, CO₂ is insufflated to enlarge the intrarectal space and therefore facilitate dissection. This is combined with a nonstop suction

Chirurgische Universitätsklinik Mannheim

Figure 2.—Bluess TEM operative rectoscope with a resected adenoma.

unit to ensure constant, high flow of gas and to evacuate the smoke due to coagulation. The operation itself was performed as described originally by Buess *et al.* [2].

Postoperatively, patients are allowed to sit and walk as soon as they have fully recovered from general anaesthesia. A liquid diet is maintained for 48 h, and patients are discharged on postoperative day 4 after a clinical examination. Initial clinical follow-up occurs at three months postoperatively by means of clinical examination, endoscopy and endorectal ultrasound. A standard follow up occurs once a year for the following years.

Patient data were prospectively collected on a personal computer and managed using Microsoft Excel. Descriptive statistics were used, and results are expressed as the absolute value, mean, standard deviation, range and percent.

Results

Over the 4-year period of study, 100 patients satisfied the indications for TEM and were included in the report. Of these 58 were male, and 42 female, with a median age of 63 years (range 35-91 years) (Table 1). Rectal blood loss was the leading symptom in 68 cases. All other patients presented with nonspecific symptoms. Two patients presented two synchronous rectal adenomas. Mean tumour location from the anal verge was 9.4 ± 2.3 cm (range 4-18 cm).

Tumours were classified in three groups according to the size of the rectal wall defect after resection. Group 1 ($n = 25$), when the defect was semicircular, group 2 ($n = 56$) includes defects representing a quarter of the circumference, and group 3 ($n = 19$) includes those tumours with a defect up to 1.5 cm.

Pre-operative endorectal sonography was performed in 97 cases. In only three the technique was not possible due to high location. Sonography correctly staged the tumour in 96 cases. The case misclassified

Table 1. Patients and tumour characteristics

Characteristic	n	Range
Gender (M/F)	58/42	
Mean age (years)	63	(35-91)
Median follow-up (months)	30	
Histology		
Adenomas	71	
Carcinomas	23	
Carcinoids	5	
Solitary ulcer	1	
Tumor distance from anal verge (cm)	9.4 ± 2.3	(4-18)

demonstrated understaging of T2 tumour assessed as T1.

All cases were performed under general anaesthesia with intra-operative repositioning of the patient for efficient surgical access with the TEM rectoscope. According to the lesion location, patients were placed in 25 cases in a jackknife position, 18 in a right lateral decubitus, 15 in a left lateral decubitus and 42 in the lithotomy position. The average operation time was 98 ± 24 min.

Full-thickness resection (mucosa and serosa down to the perirectal fat) was performed in 94 (94%) cases. In the remaining 6 (6%) cases a mucosectomy was performed in order to avoid peritoneal entry. Adenomas were resected with a 0.5 cm safety margin; in cases of carcinoma the margin was at least 1 cm. After ensuring haemostasis, the defect was closed by a 3-0 running suture in 92 cases. In the remaining 8 adenoma-cases, the defect, localized below the peritoneal reflection, was left open due to closure suture difficulties, without any postoperative morbidity. In 6 patients presenting with distal lesions, TEM resection was complemented with a conventional suture using a Park's retractor because of gas lost and impaired vision.

In two cases, we decided to 'convert' intra-operatively to laparotomy and low anterior rectal resection; the first one after tumour removal due to closure difficulties of the peritoneum, and the second after endoscopic visualization before starting transanal resection. There were eight cases of inadvertent peritoneal entry (8/100; 8%) associated with the resection of high rectal adenomas. Two of these eight patients had a postoperative clinical and radiological pneumoperitoneum but no further therapy was necessary.

Wound breakdown was discovered postoperatively in 7 cases. In two of them, due to retrorectal abscess formation and sepsis, a diverting loop ileostomy was performed. A third case in which we performed a loop ileostomy, developed a postoperative rectovaginal fistula. There was no mortality. The other 4 patients had no further complications and were managed with oral antibiotics and bowel confinement.

Immediate postoperative salvage therapy by means of low anterior rectum resection and total mesorectal excision (TME) was required in four cases (4%) after assessing high-risk features (poor differentiation and/or lymphovascular invasion) in T1 tumours ($n + 3$) or T2 ($n + 1$) invasion in the definitive histology study (Table 2). Definitive histopathology after TME confirmed T1G2 N0 ($n + 3$) and T2G2 N0 ($n + 1$) stages.

Median hospitalization was 5.5 days (range 3-21 days). A clinical examination was performed in all patients before hospital discharge.

Clinical assessment of sphincter function revealed continence impairment in 68 patients during the first three weeks after surgery. We did not perform manometric studies but only clinical assessment. Follow-up at three months showed that 98 patients had fully recovered. The two others, with already impaired continence pre-operatively, showed injury at the internal

sphincter on ultrasound examination with permanent incontinence for liquids and gas. The lesions were benign, tubulo-villous adenomas with well to moderately differentiated dysplasia, in 71 (71%) cases, 9 of them presenting as recurrence following previous open transanal resection. Malignant lesions (adenocarcinoma) were found in 23 (23%) subjects. The histological study showed 16 cases of pT1 low risk carcinoma and 5 cases of pT1 high risk tumour and 2 cases of pT2 invasion. As already mentioned in four cases salvage therapy was indicated. Two patients refused salvage therapy upon our recommendation and underwent adjuvant chemoradiotherapy. Surgical palliation decided upon pre-operatively was the indication in the other patient presenting with a T2 adenocarcinoma. The remaining pathology includes carcinoids in 5 (5%) patients and a solitary rectal ulcer in 1 (1%) case (Table 1).

Over the follow-up period of 30 months (range 6-54 months) the recurrence rate was 5%. Four cases of adenomas and one case of T1 low risk carcinoma. All four adenomas recurrences were treated with a new TEM resection. The patient with the rectal carcinoma recurrence after three years, was managed by abdominoperineal resection (APR) after radiochemotherapy (Table 2). Definitive histopathology showing a T1G2 N0 tumour.

Discussion

Over the past decade our understanding of rectal tumorigenesis has advanced rapidly: the knowledge gained has strengthened the concept of the adenomacarcinoma sequence, and has reinforced the practice of excision and surveillance [1].

Benign rectal tumours that are not suitable for snare diathermy excision are often amenable to removal by the transanal route by means of anal retractors using the method of Parks. However, this method is limited to the lower third of the rectum. In addition, it

Table 2. Complications

Results	Adenomas (n = 71)	Carcinomas (n = 23)
Wound breakdown	7	0
Ileostomy	3	0
Postop. bleeding	4	0
Pneumoperitoneum	2	0
Conversion to laparotomy	2	0
Early salvage therapy by TIME	0	4
Local recurrence	5	1
Resection with TEM	5	0
Resection with APR	0	1

may be difficult to visualize the margins of flat adenomas. Conventional management of higher rectal lesions usually involves transabdominal procedures, posterior trans-sacral (Kraske), or sphincter-splitting (York-Mason) approaches. However, these are major surgical procedures with associated complications and may be unsuitable and unnecessary if the purpose of the intervention is to resect benign lesions or low risk carcinomas [11].

With the development of TEM by Buess *et al.* [2], it is now possible to carry out formal transanal resection by means of a rectoscope, which affords excellent access and vision in the entire rectum. As a minimally invasive technique, TEM might be expected not only to benefit a small, specific population of patients in terms of morbidity but also to improve results in terms of completeness of excision and recurrence rates when compared to conventional transanal resection.

A full-thickness resection, as performed in almost all of our patients (94%), is recommended to ensure an appropriate margin of safety. In addition, we found that this standard procedure is technically easier to perform than mucosectomy and that it decreases the risk of missing a small rectal cancer that may be located inside the

villous adenoma. Such 'intraepithelial neoplasia / dysplasia' or 'early invasion' has been reported in up to 31% of cases [12].

Although it is often difficult to be sure about the precise techniques (full-thickness or submucosal) used, the rates of complete excision at the time of operation and the exact follow up, large published series report a rate of recurrences in up to 27.3% after transanal excision of rectal adenomas [13]. In contrast the results reported using TEM show a recurrence rate between 3 and 7% [3,4,7,14] with the 5.6% for the adenoma cases of the present series being well in accordance with previous studies.

The overall morbidity rate for conventional transanal surgery in reported series varies from 0 to 14.5% [13], compared with 6% reported in the largest series of a total of 318 adenomas treated by Mentges *et al.* [3]. These results are in accordance with others TEM-series reporting morbidity of up to 4.5% in more than seven hundred patients [4,7,14-16] and with the present study (7%). It is of interest that postoperative bleeding is unusual with TEM, whereas it accounts for over half of the complications reported in other series [13]. Peritoneal entry, on the other hand, as assessed intra-operatively in 8 patient (two of them with postoperative pneumoperitoneum) of our present experience, can be a significant source of morbidity with TEM. Presumably the excellent vision afforded by TEM allows more precise haemostasis, whereas the ability to excise high rectal lesions increases the likelihood of peritoneal entry, especially when the tumour is anterior, and thus more likely to be above the peritoneal reflection. In our experience all perforations were recognized and managed by immediate endorectal suture.

Our results with TEM indicate that this technique is reproducible with suitable surgical training. Interestingly all three protective ileostomies in our series occurred within the first 20 cases indicating a learning curve despite the fact

that all surgeons received a special TEM training.

Regarding the effect of a prolonged anal dilatation with the 40 mm diameter TEM operative rectoscope, manometric studies indicate a decrease in anal sphincter tone ranging from 25% to 37% of pre-operative sphincter pressure, with recovery to clinical continence within 6-16 weeks [17-19]. We observed postoperative transitory grade II incontinence in 65 of the 100 patients, with full recovery in 98% of the patients after 12 weeks. There is also evidence indicating a significantly increased risk of lowering the anal resting pressure by procedures lasting more than two hours [20]. Others found 21% disturbances of rectoanal coordination and rectal perception depending to the extent and type of resection of the tumour [21]. However, when making continence judgement, consideration would need to be given to the risk of incontinence by using the Parks retractor in conventional surgery [22], or after a low restorative rectal excision, if those were to be the alternative procedures.

To date, TEM has been used mainly for excision of sessile adenomas, and the experience with rectal cancers is limited. However, if tumours are carefully selected treatment of rectal cancer by TEM is generally accepted for so-called early low risk cancer [3]. The recurrence rate following this procedure lies between 4% and 8% [3,8,16,23] compared with a local recurrence rate of up to 30% for T1 high risk cancer [9].

The primary factor limiting the effectiveness of local treatment of early rectal cancer is lymph node invasion. The lymph node metastasis rate of T1 rectal tumours occurs between 0% and 15.4%, depending on different features [9, 24, 25]. Other adverse factors in T1 adenocarcinoma identified in the literature [26] are the following: gender, extensive budding, microacinar structures, depth of invasion in the submucosa, flat or depressed lesions, and location in the lower

third of the rectum. In our opinion and experience, positive excisional margins should not be regarded as a risk factor but should be viewed as an insufficient therapy which requires further treatment.

Therefore, careful patient selection is crucial to TEM outcome. Pre-operative staging must be precise and should be done by the surgeon himself. After clinical and endoscopy examination, assessment of histology including grading, and possible lymphovascular invasion, is mandatory. The tumour should be visualized with the rigid rectoscope to determine its distance from the dentate line and localize it precisely in the quadrant in order to plan the patient position for TEM and to avoid conversion to laparotomy due to size or localization. Endorectal sonography has always to be performed in order to assess depth of infiltration and lymph node invasion. Such precise staging is reportedly possible with an accuracy of up to 93% with respect to depth invasion and up to 81% for lymph node involvement [27, 28]. In the present study, pre-operative endosonography staging was incorrect in only one T2-carcinoma case (understage as uT1) of the 97 patients investigated.

Although the reported rate of recurrence for T1 cancers resected by TEM is between 3.8 % and 13 %, follow-up and differentiation between low and high risk features are very variable among the series published [3-6,8,9,16,23,29]. Compared with a recurrence rate of up to 18% after conventional transanal surgery [30,31] TEM seems to achieve better results in the local excision of selected rectal cancer. To date, the only published prospective, randomised study comprised 52 patients with T1 tumours treated by TEM or anterior resection [29]. There were no significant differences in group outcome: The 5-year survival was 96%; the local recurrence rate was 4.1% for TEM and 0% for anterior resection; and the metastasis rate was 0% for TEM and 4.1% for anterior resection. These results suggest that TEM

may offer some advantage relative to anterior resection for T1 rectal cancer, with similar oncological results [29].

Although adjuvant or neoadjuvant chemoradiotherapy seems to improve the prognosis after local excision, the indications for any kind of therapy following local resection of rectal cancer by TEM remain controversial. In fact, local treatment of rectal cancer is limited by the impossibility of removing the potentially positive lymph nodes, supporting the concept of adjuvant radiotherapy, chemotherapy, or both to achieve local control of the lymph nodes. The already published reports on the combined effect of TEM resection for rectal cancer followed by radiotherapy appears to support such a benefit [23]. Preliminary results suggest its reliability, with an improved recurrence-free disease survival for irradiated patients with T2 carcinomas, similar to that obtained by conventional surgery [32]. However, except for T1 low grade tumours, there is currently scant evidence to recommend the use of TEM for

curative treatment of rectal cancer, with or without adjuvant therapy.

The use of TEM for purely palliative treatment of rectal cancer is not recommended [14]. In our limited experience (one case), local resection of certain high risk tumours with TEM but without radiotherapy is possible for compromised patients or those who refuse a laparotomy. However, only a few reports confirm this use of TEM, with a local complication rate of 14% [6,23,33].

Because they are based on currently published indications and criteria for patient selection, our findings, both for adenomas and carcinomas, suggest that these standards remain adequate for identifying appropriate surgical candidates for TEM. ◀

P. Palma, S. Freudenberg, S. Samel and S. Post, Department of Surgery, University Clinic, Mannheim, Germany.

References

1. LESLIE A, CAREY FA, PRATT NR, STEELE RJC. The colorectal adenoma-carcinoma sequence. *Rev Br J Surg* 2002; 89: 845-60.
2. BUSS G, HUTTERER F, THEISS J, BÖBEL M, ISSELHARD W, PICHLMAIER H. Das System für die transanale endoskopische Rectumoperation. *Chirurg* 1984; 55: 677-80.
3. MENTGES B, BUSS G, SCHÄFER D, MANNCKE K, BECKER HD. Indications and results of local treatment of rectal cancer. *Dis Colon Rectum* 1996; 39: 886-92.
4. STEELE RJ, HERSHMAN MJ, MORTENSEN NJ, ARMITAGE NC, SCHOLEFIELD JH. Transanal endoscopic microsurgery - initial experience from three centres in the United Kingdom. *Br J Surg* 1996; 83: 207-10.
5. SMITH LE, KO ST, SACLARIDES T, CAUSHAJ P, ORKIN BA, KHANDUJA KS. Transanal endoscopic microsurgery: initial registry results. *Dis Colon Rectum* 1996; 39: 79-84.
6. LEZOCHE E, GUERRIERI M, PAGANINI A, FELICIOTTI F, DI PIETRANTONI E. Is transanal endoscopic microsurgery (TEM) a valid treatment for rectal tumors? *Surg Endosc* 1996; 10: 736-41.
7. MOERSCHER M, HEINTZ A, BUSSMANN M, JUNGINGER T. Followup after transanal endoscopic microsurgery or transanal excision of large benign rectal polyps. *Langenbecks Arch Surg* 1998; 383: 320-4.
8. WINDE G. Outcome following transanal endoscopic microsurgery. *Dis Colon Rectum* 1998; 41: 526-31.
9. HEINTZ A, MOERSCHER M, JUNGINGER T. Comparison of results after transanal endoscopic microsurgery and radical resection for T1 carcinoma of the rectum. *Surg Endosc* 1998; 12:1145-8.
10. DE GRAAF EJ, DOORNEBOSCH PG, STASSEN LP, DEBETS JM, TETTEROO GW, HOP WC. Transanal

- endoscopic microsurgery for rectal cancer. *Eur J Cancer* 2002; 38: 904-10.
11. SENGUPTA S, TJANDRA JJ. Local excision of rectal cancer. What is the evidence. *Dis Colon Rectum* 2001; 44: 1345-61.
 12. WINBURN GB. Surgical resection of villous adenomas of the rectum. *Am Surg* 1998; 64: 1170-6.
 13. SAKAMOTO GD, MACKEIGAN JM, SENAGORE AJ. Transanal excision of large rectal villous adenomas. *Dis Colon Rectum* 1991; 34: 880-5.
 14. SAID S, STIPPEL D. Transanal endoscopic microsurgery in large, sessile adenomas of the rectum: a 10-year experience. *Surg Endosc* 1995; 9: 1106-12.
 15. SWANSTROM LL, SMILEY P, ZELKO J, CAGLE L. Video endoscopic transanal-rectal tumor excision. *Am J Surg* 1997; 173:383-8.
 16. SACLARIDES TJ. Transanal endoscopic microsurgery: a single surgeon's experience. *Arch Surg* 1998; 133: 595-9.
 17. HEMINGWAY D, FLETT M, MCKEE RF, FINLAY IG. Sphincter function after transanal endoscopic microsurgical excision of rectal tumours. *Br J Surg* 1996; 83: 51-6.
 18. KREIS ME, JEHLE EC, HAUC V *et al*. Functional results after transanal endoscopic microsurgery. *Dis Colon Rectum* 1996; 39: 1116-20.
 19. BANERJEE AK, JEHLE EC, KREIS ME *et al*. Prospective study of the proctographic and functional consequences of transanal endoscopic microsurgery. *Br J Surg* 1996; 83: 211-6.
 20. KENNEDY ML, LUBOWSKI DZ, KING DW. Transanal endoscopic microsurgery excision. Is anorectal function compromised? *Dis Colon Rectum* 2002; 45: 601-4.
 21. HERMAN RM, RICHTER P, WALEGA P, POPIELA T. Anorectal sphincter function and rectal barostat study in patients following transanal endoscopic microsurgery. *Int J Colorectal Dis* 2001; 16: 370-6.
 22. VAN TETS WF, KUIJPERS JH, TRAN K, MOLLEN R, VAN GOOR H. Influence of Parks anal retractor on anal sphincter pressures. *Dis Colon Rectum* 1997; 40: 1042-5.
 23. LEZOCHÉ E, GUERRIERI M, PAGANINI AM, FELICIOTTI F. Transanal endoscopic microsurgical excision of irradiated and nonirradiated rectal cancer: a 5-year experience. *Surg Laparoscendosc* 1998; 8: 249-56.
 24. MORSON BC. Histological criteria for local excision. *Br J Surg* 1985; 72 (Suppl.): 53-4.
 25. HERMANEK P, A. Pathologist's point of view on endoscopically removed polyps of the colon and rectum. *Acta Hepatogastroenterol* 1978; 25: 169-70.
 26. NASCIBENI R, BURGART LJ, NIVATVONGS S, LARSON DR. Risk of lymph node metastasis in T1 carcinoma of the colon and rectum. *Dis Colon Rectum* 2002; 45: 200-6.
 27. MILSOM JW, LAVERY IC, STOLFI VM. The expanding utility of endoluminal ultrasonography in the management of rectal cancer. *Surgery* 1992; 112: 832-41.
 28. MASSARI M, DE SIMONE M, CIOFFI U, ROSSO L, CHIARELLI M, GABRIELLI F. Value and limits of endorectal ultrasonography for preoperative staging of rectal carcinoma. *Surg Laparosc Endosc* 1998; 8: 438-43.
 29. WINDE G, NOTTBORG H, KELLER R, SCHMID KW, BUNTE H. Surgical cure for early rectal carcinomas (T1): transanal endoscopic microsurgery vs. anterior resection. *Dis Colon Rectum* 1996; 39: 969-75.
 30. GARCÍA AGUILAR J, MELLGREN A, SIRIVONGS P, BUIE D, MADOFF RD, Rothenberger DA. Local excision of rectal cancer without adjuvant therapy: a word of caution. *Ann Surg* 2000; 3: 345-51.
 31. MADBOULY KM, REMZI FH, SENAGORE AJ, FAZIO VW, LAVERY IC. Recurrence after trans-anal excision of T1 rectal cancer: Should we be concerned? *Colorectal Dis* 2002; 4 (Suppl 1): 4.
 32. LEZOCHÉ E, GUERRIERI M, PAGANINI AM, FELICIOTTI F. Longterm results of patients with pT2 rectal cancer treated with radiotherapy and transanal endoscopic microsurgical excision. *World J Surg* 2002; 26: 1170-4.
 33. TURLER A, SCHAFER H, PICHLMAIER IL. Role of transanal endoscopic microsurgery in the palliative treatment of rectal cancer. *Scand J Gastroenterol* 1997; 32: 58-64.