

Musschia isambertoii M. Seq., R. Jardim, M. Silva & L. Carvalho (Campanulaceae), a new species from the Madeira Archipelago (Portugal)

by

Miguel Menezes de Sequeira¹, Roberto Jardim², Magda Silva¹ & Lúgia Carvalho¹

¹ Dep. Biologia/CEM, Universidade da Madeira, Campus da Penteada, 9000-390 Funchal, Portugal

² Jardim Botânico da Madeira, Caminho do Meio, 9064-512 Funchal, Portugal
sequeira@uma.pt, robertojardim.sra@gov-madeira.pt

Abstract

Menezes de Sequeira, M., Jardim, R., Silva, M. & Carvalho, L. 2007. *Musschia isambertoii* M. Seq., R. Jardim, M. Silva & L. Carvalho (Campanulaceae), a new species from the Madeira Archipelago (Portugal). *Anales Jard. Bot. Madrid* 64(2): 135-146.

A new species of *Musschia* Dumort. (Campanulaceae), endemic from Madeira Archipelago (Portugal), is described as *Musschia isambertoii* M. Seq., R. Jardim, M. Silva & L. Carvalho. Both vegetative and reproductive structures have been studied and are remarkably distinct from the recognized species [*M. aurea* (L. f.) Dumort. and *M. wollastonii* Lowe]. A full description and a diagnosis are included as well as line art, color drawings and pictures. Diagnostic characters are discussed and compared with those of the closely related species. The new species has a very restricted distribution occurring in two isolated populations in the Deserta Grande Island (Madeira Archipelago, Portugal). Nomenclatural aspects on the genus *Musschia* Dumort. are discussed and a diagnostic key for the species is also included.

Keywords: *Musschia aurea*, *Musschia wollastonii*, Deserta Grande, endemism, taxonomy.

Resumen

Menezes de Sequeira, M., Jardim, R., Silva, M. & Carvalho, L. 2007. *Musschia isambertoii* M. Seq., R. Jardim, M. Silva & L. Carvalho (Campanulaceae), una nueva especie del archipiélago de Madeira (Portugal). *Anales Jard. Bot. Madrid* 64(2): 135-146 (en inglés).

Se describe una nueva especie de *Musschia* Dumort. (Campanulaceae), endémica del archipiélago de Madeira (Portugal), *Musschia isambertoii* M. Seq., R. Jardim, M. Silva & L. Carvalho. La morfología de las estructuras vegetativas y florales estudiadas es claramente distinta de la de las otras dos especies conocidas, *Musschia aurea* (L. f.) Dumort. y *Musschia wollastonii* Lowe. Se incluyen: una descripción y una diagnosis, ilustradas por imágenes, una lámina en blanco y negro y una en color. Se comparan y discuten los caracteres diagnósticos dentro del conjunto de especies del género. La nueva especie es endémica de la isla Deserta Grande (archipiélago de Madeira, Portugal). Se discuten aspectos nomenclaturales y se incluye una clave de las especies conocidas para Madeira.

Palabras clave: *Musschia aurea*, *Musschia wollastonii*, Deserta Grande, endemismo, taxonomía.

Introduction

The Madeira archipelago is composed of three groups of islands, Porto Santo with about 20 My., Madeira and Desertas more recent and sharing a common geological origin. The Madeira island is situated between 32°38' and 32°52'N and 16°39' and 17°16'W, at approximately 600 km northwest of the Western African coast. It is a within-plate volcanic island the tip of a stratovolcano about 6 km high. The emerged part of the island dates back to Post-Miocene times, < 5.6 My. (Ribeiro & al., 2005), and the more recent volcanic

activity took place 6000-7000 years B.P. (Geldmacher & al., 2000). The Madeira island has 737 km² of surface. The maximum altitude is at Pico Ruivo, reaching 1861 m above sea level.

To the southeast the Madeira archipelago is continued by the Desertas sub-archipelago composed by three small islands. The northernmost is Ilhéu Chão (ca. 0.5 km²), which is also the smallest with only 100 m above sea level. The largest, Deserta Grande (ca. 10 km²), has a maximum altitude of 442 m above sea level. Finally Bugio (ca. 3 km²), the southern island has a maximum altitude of 348 m above sea level (Press & Short, 1994).

The vascular flora of Madeira comprehends 1226 species (Press & Short, 1994), including 780 autochthonous. Of these 234 are Macaronesia endemics (Vieira, 1992), 157 of them restricted to Madeira (if hybrids are to be considered the total number rises to 165, Jardim & Francisco, 2000). Six endemic genera occur in the Madeira archipelago: *Chamaemeles* Lindl. (*C. coriacea* Lindl., Rosaceae), *Melanoselinum* Hoffm. [*M. decipiens* (Schrad. & J.C. Wendl.) Hoffm., Apiaceae], *Monizia* Lowe (*M. edulis* Lowe, Apiaceae), *Parafestuca* E.B. Alexeev [*P. albida* (Lowe) E.B. Alexeev, recently ascribed to *Koeleria* as *Koeleria loweana* Quintanar, Catálan & Castrov., Poaceae], *Musschia* Dumort. [until now *M. aurea* (L. f.) Dumort. and *M. wollastonii* Lowe, Campanulaceae] and *Sinapidendron* Lowe [*S. angustifolium* (D.C.) Lowe, *S. frutescens* (Aiton) Lowe, *S. gymnocalyx* (Lowe) Rustan, *S. rupestre* Lowe and *S. sempervivifolium* Menezes, Brassicaceae]. The 15 Macaronesian endemic genera include: *Aichryson* Webb & Berthel. and *Monanthes* Haw. (Crassulaceae), *Argyranthemum* Webb, *Pericallis* Webb & Berthel. and *Schizogyne* Cass. (Asteraceae), *Bencomia* Webb & Berthel. and *Marcetella* Svent. (Rosaceae), *Bystropogon* L'Hér. and *Cedronella* Moench (Lamiaceae), *Drusa* DC. (Apiaceae), *Isoplexis* (Lindl.) Benth. (Scrophulariaceae), *Phyllis* L. (Rubiaceae), *Picconia* DC. (Oleaceae), *Semele* Kunth (Ruscaceae) and *Visnea* L. f. (Theaceae). According to Press & Short (1994), the most important families and also the ones with the most endemic species are: Asteraceae (132 species, 24 endemics), Brassicaceae (47 species, 11 endemics), Crassulaceae (20 species, 7 endemics), Lamiaceae (39 species, 9 endemics), Liliaceae s.l. (24 species, 7 endemics) and Poaceae (139 species, 8 endemics). There are also 75 species of ferns including 14 Madeira and Macaronesian endemics.

The family Campanulaceae includes around 82 genera and 2000 species (Mabberley, 1997). In the Macaronesian region two genera and eight endemic species are recognized, *Azorina* Feer a monospecific genus [*Azorina vidalii* (H.C. Watson) Feer] endemic from the Azores archipelago, *Musschia* Dumort. endemic from the Madeira archipelago with two recognized species *M. aurea* (L. f.) Dumort. and *M. wollastonii* Lowe, the Canary islands endemics *Canarina canariensis* (L.) Vatke and *Campanula occidentalis* Y. Nymann, the Cape Vert endemics *Campanula bravenensis* (Bolle) A. Chev. and *Campanula jacobaea* C. Sm. ex Hook. and *Wahlenbergia lobelioides* (L. f.) Link subsp. *lobelioides* common to Madeira, Canary and Cape Vert archipelagos (Hansen & Sunding, 1993; Tebbs, 1994; Turland, 1994; Acebes Ginovés & al., 2004; Sánchez-Pinto & al., 2005; Silva & al., 2005).

The genus *Musschia* was first isolated from *Campanula* L. by Dumortier (1823) based on *Campanula aurea* L. f. The differential characters of *Musschia* were described by Dumortier (op. cit.) as “*Car. Diff. Calyx quinquepartitus. Corolla basi calycis inserta, quinquepartita. Stamina basi serrato-dillatata inflexa. Stigmata quinque convoluta. Capsula quinquelocularis*”. The name *Musschia* was dedicated to Jean Henri Musshe (1765-1834), Director of the Gent Botanical Garden by Dumortier (1823).

As stated, the first species was described previously by Linnaeus f. (1782) as *Campanula aurea* L. f. as “*Campanula (aurea) capsulis quinquelocularibus, stigmatibus, quinquefidis, caule paniculato, foliis duplicato-serratis; Habitat in Insula Madera. Fr. Masson. Caulis paniculati, rupibus adpressi. Folia lanceolata, glabra. Calyx superus, coloratus. Corollae tubus a calyce distans: laciniis linearibus, reflexis*”.

Later Dumortier (1823) segregated the genus *Musschia* from *Campanula* based on *Campanula aurea* L. f., as *M. aurea* (L. f.) Dumort., and apart from the type specie, he described *Musschia angustifolia* Dumort. Later Lowe (1856) described *M. wollastonii* Lowe.

The early (1777-1830) references to *Musschia* (as *Campanula* L.) are all presumably based on seeds taken to the mainland from Madeira by Masson but also (possibly) by other authors. Ker Gawler (1815) not only includes a beautiful drawing of *Musschia aurea* (as *Campanula aurea*), but also a full and detailed description of this species. The seeds, as stated by Ker Gawler (op. cit.), were taken to England by Masson in 1777, previous to the description by Linnaeus f. (1782). The list of plants collected in Madeira by Francis Masson, including notes, was never published. Two cultivated species at Kew Gardens are referred by Ker Gawler (op. cit.) but the second species corresponds to *Campanula lobelioides* L. f. [later included in the genus *Wahlenbergia* as *W. lobelioides* (L. f.) Link]. *Musschia aurea* had already been referred by Aiton (1789) as *Campanula aurea* L. f. cultivated in Kew Gardens. The early history of Madeira Campanulaceae, as with many other species and families, seems to have an origin in Masson's collections, lists and cultivations.

Ker Gawler (1815) refers to *Campanula aurea* with two varieties, α *latifolia*, which he refers to an icon in Ventenat (1805), and β *angustifolia*, with no description other than a reference to Jacquin (1804) *Plantarum Rariorum Horti Caesari Schoenbrunnensis Descriptiones et Icones*. The line drawing in Jacquin (op. cit) comes with no description or table but clearly corresponds to a narrow leaved plant (Fig. 1). The publi-



Fig. 1. Reproduction of the lamina of *Campanula aurea* L. f. included in Jacquin (1804) [from <http://gallica.bnf.fr>]. This plate was referred by Ker Gawler (1815) as *Campanula aurea* L. f. β *angustifolia*.

cation *Jardin de Malmaison* by Ventenat (op. cit.) includes drawings by Redouté.

Later, De Candolle (1830, 1838) refers to *Musschia aurea* and to the unpublished notes by Masson to cite the β *angustifolia* (which he describes as *foliis angustioribus*). However, Dumortier (1823) had already proposed the name and given a description of these narrow leaved plants as *Musschia angustifolia* Dumort. This name is presumably based on Ker Gawler (1815, as *C. aurea* var. *angustifolia*) while Dumortier makes no reference to previous descriptions or icons of this narrow leaved variety, he does, however, distinguish the habitat of α (*Musschia aurea*) “*in littore*” and β (*Musschia angustifolia*) “*In interioribus insulae*”, suggesting that he based his notes on Ker Gawler (op. cit.) which states the same geographic pattern based on Masson’s unpublished notes.

Dumortier (1823) is in fact the first to give a description and to name correctly the narrow leaved *Musschia aurea* plants. If no plants were to be found in order to name a lectotype, it would be advisable to elect the icon published by Jacquin (1804).

However, *Musschia angustifolia* Dumort. has not been recognized by later authors, in fact only two species, *M. aurea* and *M. wollastonii*, are included in the genus *Musschia* Dumort. by Lowe (1856, 1868), Menezes (1914) and the more recent authors (Hansen & Sunding, 1993; Turland, 1994; Jardim & Francisco, 2000; Costa & al., 2004).

Material and Methods

Plants belonging to the new taxon were collected during spring of 2006 in the Desertas islands and deposited at MA (typus). Studied plants of the genus *Musschia* were examined from MADJ (including MADS) and MADM. Macroscopic analysis was performed using a binocular microscope Zeiss model SV 11 APO.

The distribution map is based on geographic coordinates of the studied specimens converted to Cartesian coordinates using a local *Datum* (Porto Santo) and plotted using ArcView GIS 3.1.

Results and Discussion

During field work in 2006, M. Silva and L. Carvalho collected *Musschia* specimens in Deserta Grande at Fajã Pequena. These specimens present a distinctive morphology not within the variability of the previously described taxa (e.g. leaf size, shape and indumentum, the inflorescence architecture including branching and flower colours). Accordingly a new species is proposed.

Musschia isambertoii M. Seq., R. Jardim, M. Silva & L. Carvalho, **sp. nov.** (Figs. 2, 3)

Speciebus Musschia aurea Dumort. et *M. wollastonii* Lowe similis; ab eis vero differens caule florifero nonnumquam 1.5 m alto atque tantum apice ramoso, floribus in apicibus axis principalis atque ramorum dense congesti, sepalis autem apice atque nervis castaneo-rubentibus, corollis viridibus (tantum basi flavescentibus), antheris maioribus (c. 15.5 mm), lobis denique stigmatibus pariter maioribus (c. 16 mm).

Tall rosetted monocarpic plant (Fig. 3), up to 2 m. Stem stout, ascending and short, up to 40 cm, unbranched (Fig. 3). Leaves 25-33 × 11-13 cm, rigid, dull green with whitish veins, semi-clasping the stem, with indistinct petiole, oblanceolate, herbaceous, abaxial face glabrous except for the pubescent veins with hairs 0.3-0.4 mm, adaxial face sparsely scabrid; margin shallowly undulated, bi-serrate (Figs. 2 b, 3). Inflorescence (Fig. 2 a, 3) up to 150 cm long, unbranched except for the terminal part, with 2-3(4) long (up to 25 cm) branches disposed at a right angle to the main axis; bracts (Fig. 2 g) gradually shorter towards the top, 1.5-2 cm oblanceolate to lanceolate, undulate, clasping the stem and enrolled to their adaxial face (Fig. 2 e); the flowers densely crowded on the apex of the main axis and less so on the branches (Figs. 2 a, 3, 5). Flower (Figs. 2 f, h, 7 a) up to 5 cm, nectariferous; green, yellowish with reddish-brown tones, axillary to short bracts up to 2 cm oblanceolate to ovate, acuminate; with a usually short and puberulous pedicel. Calix (Figs. 2 e, 7 a) with sepals 23-25 × 5-7.4 mm (10 mm at the base), green with reddish brown apex and veins, sometimes yellowish towards the base, oblanceolate to oblong, sub-apiculate to acuminate, sepals with 3 nerves, the central larger, the margin and apex reticulate, puberulous with hairs up to 0.05-0.1 mm. Corolla bright green (Figs. 2 h, 7 a), with tube 13-14.5 mm, yellowish towards the base, lobes ca. 20 × 4.5-5(6) mm, narrowly triangular-lanceolate, acuminate, erect before anthesis, arcuate deflex during anthesis. Stamens (Fig. 2 i) with filament 10-11 mm, winged, 2.4-2.5 mm wide at the base; anthers 15.2-15.5 mm, shortly apiculate, 0.5-0.8 mm, subcordate. Pistil (Figs. 2 j, 7 a) with style 29-30 mm, stigma 15-16.3 × 2.5-3 mm, scabrid towards the base with hairs 0.2-0.4 mm. Hypantial tube (Figs. 2 e, h, j, 7 a) 16-17 × 17 mm, minutely scabrous with hairs 0.2 (0.1-0.3) mm, green with a bright yellow apical ring and reddish-brown and green ribs. Capsule n.v. Plant pollinated by lizards (*Lacerta dugesii maui*).

Type specimens. Portugal, Madeira: Ilhas Desertas, Deserta Grande, perto da Fajã Pequena, Porto das




Fig. 2. *Musschia isambertoi*: **a**, inflorescence; **b**, leaf; **c**, detail of the abaxial leaf surface; **d**, detail of the adaxial leaf surface; **e**, portion of an inflorescence lateral branch; **f**, indumentum of the pedicel; **g**, bract; **h**, flower; **i**, stamen; **j**, flower showing the ovary and ovules [*M. Silva* 868 (MA 751556)].



Fig. 3. *Musschia isambertoi*, colour drawing of the plant habit by Juan Castillo [*M. Silva* 868 (MA 751556)].

Moças, 16-V-2006, *M. Silva, L. Carvalho, C. Viveiros & P. Gouveia* 868 (MA 751556) (holotype). Portugal, Madeira: Deserta Grande: Rocha basáltica por baixo da casa antiga de vigias das baleias do lado oeste. Local exposto ao sol da tarde e aos ventos de oeste, Vertente oeste da Deserta Grande. 350 m, 10-VI-1992, *I. Silva* (MADM) (paratype, sub *M. wollastonii* Lowe).

Etymology. Dedicated and named for its first collector Isamberto Silva, collector of many important plants and arthropods, several of them corresponding to new species to the Madeira archipelago in collaboration with many botanists (and entomologists). Isamberto Silva is also the author of some articles on the flora of Desertas and Selvagens Islands (eg. Costa Neves & al., 1992; Menezes de Sequeira & Silva, 2007).

Distribution. *Musschia isambertoii* is known from only two populations on the Deserta Grande Island, including the population from Fajã Pequena (Fig. 4). *Musschia aurea* is also present (and is much more common) on this island.

Habitat. *Musschia isambertoii* seems to grow preferably at almost sea level at damp, the extraordinary habit of this plant immediately strikes the observer, in fact when in flower, it is the largest macro-

herb of the Desertas. It does not share a common ecology with *Musschia aurea* on the Desertas as on Madeira, where *M. aurea* is a chasmophyte. Several individuals of the Desertas endemic lizard *Lacerta dugesii mauli* Mertens, 1938, were seen pollinating the flowers of *Musschia isambertoii* (Fig. 5 b, c). Elvers (1978) refers to lizard pollination in Madeiran plants of *Musschia aurea* and Olesen & Valido (2003) further stress the role of lizards as pollinators and seed dispersers on islands, referring to *Musschia aurea* and *Lacerta dugesii* Milne-Edwards, 1829. Recently Traveset & Richardson (2006) refer to biological invasions as possible disrupters of mutualism and consequences on conservation, this seems to be the case in the Deserta Grande where goats but also rabbits were responsible for massive vegetation destruction. Goat and rabbit eradication was initiated in 1996. Rabbits were totally eradicated and the reduction of goat numbers was followed by a rapid recovery of the flora. The recent increase in goat population seems to be driving most endemics to a population decrease, both in number and in density. On Madeira, the competition by invading plants, such as *Opuntia tuna* (L.) Mill., may also cause a disruption of mutualism rela-




Fig. 4. Distribution of *Musschia aurea*, yellow squares; *M. wollastonii*, red squares; *M. isambertoii*, green squares.

tions between *Lacerta dugesii dugesii* and *Musschia aurea*. Future works on the biological relations between *Lacerta dugesii dugesii* and *Musschia aurea* as well as *M. isambertoii* and the Deserta endemic *Lacerta dugesii mauli* should enlighten us regarding both mutualism relationships and conservation aspects. Flower colours and inflorescence typical structure are the most important morphological traits that may have evolved from mutualism with reptile pollinators as a co-evolutionary process.

Phenology. Data on flowering is scarce however it seems to flower from May to June.

Conservation status. Due to the scarce number of populations and the reduced occupancy and occurrence area as defined by IUCN (2001), and also due to the grazing effects through the introduction of goats, this new species should be considered as Critically Endangered (CR, C2a(i,ii); D).

Taxonomic remarks. Specimens with the same distinctive morphology had been already collected by Isamberto Silva and included at the local plant collection in Deserta Grande Natural Park facilities, a duplicate of this collection, only a few flowers and a bract, was included at MADM herbarium as *Muss-*

chia wollastonii Lowe, and cited as so in Costa Neves & al. (1992). Our results strongly suggest a taxonomic differentiation at species level. In fact several morphological characters showed a clear discrepancy with the previously known species. Habit (Figs. 5 a, 6 a-c), leaf shape and hairiness (semi-clasping the stem, oblanceolate, rigid herbaceous, dull, abaxially pubescent and adaxial sparsely scabrid), inflorescence remarkable structure and size (Figs. 5 b, 6), long and unbranched except for the terminal part, with branches disposed at a right angle to the main axis, flower characters (Fig. 7) including: the green to yellowish sepals with their reddish brown apex and veins, larger than those of *M. aurea* and wider than in *M. wollastonii*; the green corolla with yellowish base, and wide (4.5-6 mm) lobes; the larger anthers, three times as large as those of *M. aurea* and twice those of *M. wollastonii*; and the larger stigma, all correspond to diagnostic characters. Table 1 summarizes the diagnostic characters of *Musschia isambertoii* and compares them with those of *M. aurea* and *M. wollastonii*. Further studies including chromosome counts and molecular markers may bear light on origin of this new taxon, including a possible hybrid origin that



Fig. 5. Habit and inflorescence of *Musschia isambertoii*: **a**, habit, one of the authors (L. Carvalho) in Deserta Grande (Porto das Moças); **b**, aspect of the inflorescence; **c**, flowers and pollinating lizard, *Lacerta dugesii mauli* (photographies: M. Silva).

Table 1. Diagnostic characters of *Musschia isambertoii* versus *M. aurea* and *M. wollastonii*.

	<i>M. isambertoii</i>	<i>M. aurea</i>	<i>M. wollastonii</i>
Indumentum	Glabrous to shortly scabrid	Glabrous	Pubescent
Habit	Tall rosetted monocarpic, up to 2 m	Short not monocarpic, ca. 1 m	Tall rosetted monocarpic, up to 3 m
Stem	Stout enlarged, unbranched and ascending up to 20-40 cm	Branched stock supporting several leaf rosettes, up to 50 cm	Robust stout ascending and woody, up to 200 cm
Leaves	25-33 × 11-13 cm, semiclasping the stem, with indistinct petiole, oblanceolate, rigid herbaceous, dull, abaxial face glabrous except for the pubescent veins, adaxial face sparsely scabrid	10-35 × 2-5.5 cm, not clasping the stem, ovate to narrow elliptic, coriaceous, shiny, glabrous	14-19 × 3.5-17 cm, clasping the stem, narrowly oblanceolate, soft herbaceous, dull, with pubescent veins
Inflorescence	150 cm, branched at the apex, not pyramidal, flowers densely crowded	40 cm, much branched, pyramidal, flowers not crowded	100 cm, much branched, pyramidal, flowers not crowded
Sepals	23-25 × 5-10 mm, green with reddish brown apex and veins, sometimes yellowish towards the base, oblanceolate to oblong, subapiculate to acuminate	12-18 × 4-10 mm, green, triangular-ovate, cuspidate	20-25 × 6 mm, green to reddish-brown, narrow triangular-lanceolate, cuspidate
Corolla	Green to olive green, yellowish towards the base, lobes 20 × 4.5-6 mm	Bright yellow, lobes 11-16 × 3-4 mm	Reddish brown or sometimes yellowish, lobes 21-28 × 3-5 mm
Stamens	15-16 mm, shortly apiculate, pollen yellow	5-6 mm, shortly apiculate, pollen yellow	8-10 mm, apiculate, pollen white-pinkish
Stigma	15-16 mm	7-11 mm	14-15 mm

could explain some morphological traits (e.g. indumentum and size).

KEY TO *MUSSCHIA* SPECIES

- Plant with leaf rosettes on short stems (less than 50 cm); corolla lobes bright yellow or green. 2
Plant with leaf rosettes on long stems up to 2 meters usually monocarpic; corolla lobes reddish brown or yellowish
..... **2. *M. wollastonii***
- Plant monocarpic; leaf rosettes solitary supported by a non branched stem; leaves dull; corolla-lobes green more than 17 mm long (ca. 20 mm); anthers 15 mm **3. *M. isambertoii***
Plant not monocarpic; leaf rosettes supported by a usually much branched stock; leaves shiny; corolla-lobes bright yellow less than 17 mm; anthers less than 7 mm . **1. *M. aurea***

Nomenclature

1. *Musschia aurea* (L. f.) Dumort., Comment. Bot.: 28. 1822

Campanula aurea L. f., Suppl. Pl. 141. 1782

Musschia angustifolia Dumort., Comment. Bot.: 29. 1822

Campanula aurea (L. f.) Dumort. var. *angustifolia* Ker-Gawl. in Bot. Reg. 1: 57. 1815, nom. nudum;
Musschia aurea (L. f.) Dumort. var. *angustifolia* Ker-Gawl. ex DC., Monogr. Campan. 369. 1830

2. *Musschia wollastonii* Lowe in Hook. Kew Journ. 8: 298. 1856

3. *Musschia isambertoii* M. Seq., R. Jardim, M. Silva & L. Carvalho

Studied Material

Musschia aurea (* correspond to the narrow leaved form). PORTUGAL. **Madeira:** Garajau, 20-VIII-2001, *G. Quinn s.n.* (MADM 484). Ribeiro Frio, X-1973, *C. Pickering 1067485* (MADM 484) (*). Caminho para a Praia do Garajau, VI-2002, *J.G. Quinn s.n.* (MADM 484). Boaventura, XI (MADS 829) (*). Falésias do Cabo Girão no Poio da Manga abaixo do miradouro, 29-IX-1982, *Nóbrega s.n.* (MADJ 4044) (*). Deserta Grande, 12-IV-1983, *Nóbrega & Rui Santos s.n.* (MADJ 6771). Estrada do Curral das Freiras, 25-V-1962, *Rui Santos s.n.* (MADJ 4042) (*). Estrada do Curral das Freiras, entre túnel do Serrado e o túnel do Córrego dos Açougues, 12-VII-1992 *Nóbrega s.n.* (MADJ 7786) (sub *M. aurea* var. *longifolia*, *). Estrada do Curral das Freiras, entre os dois túneis do Serrado e Córrego de Açougues, 14-VII-1992, *Nóbrega s.n.* (MADJ 7785) (*). Fajã da Areia, São Vicente, 6-IX-1994, *Nóbrega s.n.* (MADJ 8374) (*). Fajã dos Asnos do fundo do miradouro do Cabo Girão, 20-IV-1992, *Nóbrega s.n.* (MADJ 7787). Garajau, VII-1921 (MADS 825). Quinta do Bom Sucesso na berm de uma antiga levada, 10-VI-1957, *Beliz & R. Santos s.n.* (MADJ 4040). Levada do Norte, entre a Serra de Água e Ribeira Brava, Rocha Alta do Espigão, 17-VI-1982, *Nóbrega, Pita & Rui Santos s.n.* (MADJ 4046) (sub *M. aurea* f. *angustifolia*, *). Túnel da Ponta Delgada, 1-II-1984, *Nóbrega, Pita, Rui Santos & Duarte s.n.* (MADJ 4045) (*). Entre a Ponta do Sol e Madalena do Mar, próximo dos Anjos, 6-VI-1957, *Beliz & Rui Santos s.n.* (MADJ 4039). Praia Formosa, VIII (MADS 826). Estrada de Ribeira Brava, Lugar de Baixo, numa parede, 2-VIII-1957, *R. Vieira & R. Santos s.n.* (MADJ 4041). Rocha da Escada, falésias marítimas, Santana, 17-XII-1987, *Nóbrega s.n.* (MADJ 5449) (*). São Martinho, muros de areão ao lado da igreja, 19-VIII-1980, *Domingos Noia s.n.* (MADJ



Fig. 6. Habit and inflorescence of *Musschia aurea* and *M. wollastonii*: **a**, *M. aurea*; **b**, *M. aurea*, plant with narrow leaves; **c**, habit of *M. wollastonii*; **d**, inflorescence of *M. wollastonii* (photographies: M. Menezes de Sequeira).




Fig. 7. Detail of a flower, several diagnostic characters are shown, such as sepal colour and veins, green corolla lobes: **a**, *Musschia isambertoi*; **b**, *M. aurea*; **c**, *M. wollastonii* (photographies: a, M. Silva; b, c, M. Menezes de Sequeira).

4043). Volta da Malhada, estrada do Curral das Freiras, vertente da Rib. dos Socorridos, 25-II-1985, *Nóbrega, Paulo & Costa s.n.* (MADJ 4047) (sub *M. aurea* f. *angustifolia*, *). Acima da Praia do Garajau, 14-XII-1989, *Nóbrega s.n.* (MADJ 2992). São Gonçalo, acima do miradouro, num talude rochoso, 18-V-2005, 255 m, *M. Sequeira 4618* (UMad).

Musschia isambertoi. PORTUGAL. **Madeira:** Ilhas Desertas, Deserta Grande, perto da Fajã Pequena, Porto das Moças, 16-V-2006, *M. Silva, L. Carvalho, C. Viveiros & P. Gouveia 868* (MA 751556) (holotype). Deserta Grande, rocha basáltica por baixo da casa antiga de vigias das baleias do lado oeste, local exposto ao sol da tarde e aos ventos de oeste, vertente oeste da Deserta Grande, 10-VI-1992, 350 m, *Isamberto Silva s.n.* (MADM) (paratype, sub *M. wollastonii*). Deserta Grande Porto das Moças, 22-VI-2001, *J.A. Carvalho & T. Pontes s.n.* (MADJ 9586). Deserta Grande, Porto das Moças, 22-VI-2001, *J.A. Carvalho & T. Pontes s.n.* (MADJ 9587).

Musschia wollastonii. PORTUGAL. **Madeira:** Rib. Frio, VIII-1938, *P.T.O. Costa s.n.* (MADM 485). Escarpa sobre a Ribeira do Juncal, Córrego do Sabugueiro (S. Roque do Faial), 4-X-1973, *C. Andrada s.n.* (MADJ 4050). Encumeada, São Vicente, 7-XI-2001, *Olga Baeta & P. Gouveia s.n.* (MADJ 9689). Levada Central da Rib^a da Janela, 28-VI-1984, *Nóbrega & Rui Santos s.n.* (MADJ 4054). Queimadas, 22-IX-1962, *Rui Vieira & Rui Santos s.n.* (MADJ 4049). Queimadas, 20-II-1975, *Rui Vieira s.n.* (MADJ 4057). Rib^a da Ponte dos Ganchos, no Urzal de Boaventura, 28-VIII-1985, *Nóbrega s.n.* (MADJ 4055). Rib^a do Urzal, Boaventura, junto à levada nova, Caldeirão Verde, 18-I-1984, *Nóbrega s.n.* (MADJ 4053). Ribeira das Lages, 2-VIII-1962, *R. Vieira s.n.* (MADJ 4048). Ribeira Funda do Seixal, junto de cascatas que descem do Fanal, 27-VI-1989, *Nóbrega s.n.* (MADJ 6574). Ribeira Grande, São Vicente, no términos da vereda do Chão dos Louros para a Ribeira Grande, 4-X-1988, *Nóbrega s.n.* (MADJ 6460). Ribeiro das Feijocas, ao longo da Encumeada de S. Vicente, lev. no km 13, 18-X-1983, *Nóbrega, Pita & Isidoro s.n.* (MADJ 4052). Parque Florestal do Ribeiro Frio, 5-XI-1982, *Nóbrega & Pita* (MADJ

4051). Rib^o do Velho, Ponta Delgada, junto ao “topo de Ponta Delgada”, 3-X-1986, *Nóbrega s.n.* (MADJ 4056). Ribeira do Juncal, abaixo da levada antiga que dava para o Ribeiro Frio no fundo da Ribeira, 7-VIII-1990, *Nóbrega s.n.* (MADJ 2818).

Acknowledgements

We are grateful to the Natural Park of Madeira especially to the director, Susana Fontinha, and to Dília Menezes for collecting premises and the use of the Natural Park of Madeira facilities during field work. We are also grateful to Carlos Viveiros and Pedro Gouveia who guided the field and sea excursions. The authors would also like to acknowledge the help of Sandra Mesquita for the map and Leopoldo Medina for his literature search at the Real Jardín Botánico de Madrid library. The drawings included are due to Juan Castillo and the authors would like to thank him especially. Finally the authors would also like acknowledge to Father M. Laínz for the Latin version of the diagnosis and comments on plant morphology. This article would not be possible without the encouragement of Santiago Castroviejo, his help and reviews were crucial. Finally the authors would like to express their thanks to an anonymous reviewer, her/his critique and suggestions strongly influenced the final output.

References

- Acebes Ginovés, J.R., Del Arco Aguilar, M., García Gallo, A., León Arencibia, M.C., Pérez De Paz, P.L., Rodríguez Delgado, O., Wildpret De La Torre, W., Martín Osorio, V.E., Marrero Gómez, M.C. & Rodríguez Navarro, M.L. 2004. Pteridophyta & Spermatophyta. In: Izquierdo, I., Martín, J.L., Zurita, N. & Arechavaleta, M. (eds.), *Lista de especies silvestres de Canarias (hongos, plantas y animales terrestres)*. Pp. 96-143. Consejería de Medio Ambiente y Ordenación Territorial. Gobierno de Canarias.
- Aiton, W. 1789. *Hortus Kewensis, or a Catalogue of the Plants cultivated in the Royal Botanic Garden at Kew*. Vol. 1. Lon-

- don. Printed for George Nicol, Bookseller to his Majesty, Pall Mall.
- Costa Neves, H., Silva, I. & Palmeira, C. 1992. Contributions to the knowledge of the flora of Desertas Islands. *Bocagiana* 163: 1-21.
- Costa, J.C., Capelo, J., Jardim, R. & Menezes de Sequeira, M. 2004. Catálogo Florístico da Ilha da Madeira. *Quercetea* 6: 187-200.
- De Candolle, A.P. 1830. *Monographie des Campanulées*. Paris. Veuve Desray.
- De Candolle, A.P. 1838. *Prodromus systematis naturalis regni vegetabilis sive Enumeratio contracta ordinum generum specierum plantarum huc usque cognitarum, juxta methodi naturalis normas digesta*. Vol. 7(1). Parisiis. Sumptibus Sociorum Treuttel et Würtz.
- Dumortier, B.C. 1823. *Commentationes botanicae. Observations botaniques, dédiées à la Société d'Horticulture de Tournay par B.C. Dumortier*. Tournay: Imprimerie Ch. Casterman.
- Elvers, I. 1978. The Madeiran lizard - flower connection observed in a natural habitat. *Botaniska Notiser* 131(1): 159-160.
- Geldmacher, J., Bogaard, P., Hoernle, K. & Schmincke, H. 2000. Ar age dating of the Madeira Archipelago and hotspot track (eastern North Atlantic). *Geochemistry, Geophysics, Geosystems* 1(2) (DOI 10.1029/1999GC000018).
- Hansen, A. & Sunding, P. 1993. Flora of Macaronesia, Checklist of vascular plants, 4. revised edition. *Sommerfeltia* 17: 1-295.
- IUCN. 2001. *IUCN Red List categories and Criteria, version 3.1*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge. United Kingdom.
- Jacquín, N. 1804. *Plantarum Rariorum Horti Caesarei Schoenbrunnensis Descriptiones et Icones*. Vol. 4. Vienna, London.
- Jardim, R. & Francisco, D. 2000. *Flora Endémica da Madeira*, Múchia Publicações. Funchal.
- Ker Gawler, J.B. 1815. *Campanula aurea*. *Botanical Register* 1: 57.
- Linnaeus f., C. 1782. *Supplementum Plantarum Systematis Vegetabilium Editionis Decimae Tertiae, Generum Plantarum Editiones Sextae, et Specierum Plantarum Editionis Secundae*. Editum a Carolo a Linné. Brunsvigae.
- Lowe, R.T. 1856. Species Plantarum Maderensium quaedam Novae, vel hactenus ineditae, breviter descriptae. *Hooker's journal of botany and Kew Garden miscellany* 8: 289-302.
- Lowe, R.T. 1868. *A Manual for Flora of Madeira and the adjacent Islands of Porto Santo and the Deserts*. Vol. 1(5): 574-577. John Van Voorst. London.
- Mabberley, D.J. 1997. *The Plant Book*. Second Edition. Cambridge University Press. United Kingdom.
- Menezes, C.A. 1914. *Flora do Arquipélago da Madeira (Phanerogamicas e Cryptogamicas Vasculares)*. Funchal. Typ. Bazar do Povo.
- Menezes de Sequeira & Silva, I. 2007. Notes on some grasses from the Selvagem Grande (Selvagens archipelago, Portugal). *Boletim do Museu Municipal do Funchal* (submitted).
- Olesen, J.M. & Valido, A. 2003. Lizards as pollinators and seed dispersers: an island phenomenon. *Trends in Ecology and Evolution* 18: 177-181.
- Press, J.R. & Short, M.J. 1994. *Flora of Madeira*. HMSO. London.
- Ribeiro, L., Ech-chakroun, S., Mata, J., Boven, A., Mattielli, N., Hus, J & Maerschalk, C. 2005. Elemental and lead isotopic evidence for coeval heterogeneities at Madeira/Desertas mantle source. In: Anonymous (ed.), *Acts of VIII Congresso de Geológica dos Países de Língua Portuguesa (Portugal)*. Pp. 485-488.
- Tebbs, B.R. 1994. Campanulaceae. In: Press, J.R. & Short, M.J. (eds.), *Flora of Madeira*. Pp. 329-333. HMSO. London.
- Turland, N.J. 1994. *Musschia Dumort.* In: Press, J.R. & Short, M.J. (eds.), *Flora of Madeira*. Pp. 331. HMSO. London.
- Sánchez-Pinto, L., Leticia Rodríguez, M., Rodríguez, S., Martín, K., Cabrera, A. & Carmen Marrero, M. 2005. Pteridophyta, Spermatophyta. In: Arechavaleta, M., Zurita, N., Marrero, M.C. & Martín, J.L. (eds.), *Lista preliminar de especies silvestres de Cabio Verde (hongos, plantas y animals terrestres)*. Pp. 38-57. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias.
- Silva, L., Pinto, N., Press, B., Rumsay, F., Carine, M., Henderson, S. & Sjögren, E. 2005. Lista das plantas vasculares (Pteridophyta e Spermatophyta). In: Borges, P.A.V., Cunha, R., Gabriel, R., Martins, A.F., Silva, L. & Vieira, V. (eds.), *A list of the terrestrial fauna (Mollusca and Arthropoda) and flora (Bryophyta, Pteridophyta and Spermatophyta) from the Azores*. Pp. 131-156. Direcção Regional do Ambiente and Universidade dos Açores, Horta, Angra do Heroísmo and Ponta Delgada.
- Traveset, A. & Richardson, D.M. 2006. Biological invasions as disruptors of plant reproductive mutualisms. *Trends in Ecology and Evolution* 21: 208-216.
- Ventenat, E.P. 1805. *Jardin de la Malmaison*. Vol. 2. Paris.
- Vieira, R. 1992. *Flora da Madeira. O interesse das plantas endémicas macaronésicas*. Serviço Nacional de Parques, Reservas e Conservação da Natureza. Lisboa.

Associate Editor: I. Álvarez

Received: 10-VIII-2007

Accepted: 17-X-2007