

EFFECTIVENESS OF SMALL GROUP TRAINING ON BEHAVIORAL SKILLS OF ICU NURSES

(Recibido 05-06-2017. Aprobado el 07-09-2017)

**Afshin Hazrati
Mernglo**
*MSc student,
intensive care
nursing, faculty of
nursing and
midwifery, Urmia
University of
Medical Science*

Molod Radfar
*PhD of Nursing
education,
faculty of nursing
and midwifery,
Urmia University
of Medical
Science*

**Yousef
Mohammadpour**
*Academic board member of
faculty of nursing and
midwifery, Urmia
University of Medical
Science*

Naser Sheikhi
*Statistics expert,
Urmia University
of Medical
Science*

Abstract. Aim and background: Numerous errors can occur during administration of nutrient-medication orders through feeding tubes. Proper performance of the nurses can enhance the patients' safety and prevent from the future complications. The aim of this study is to evaluate the effect of training in small groups on the performance of nurses in administration of nutrient-medication orders.

Materials and methods: This is an experimental study conducted in Imam Khomeini Hospital of Urmia. 80 nurses were entered to the study by census sampling method. Their knowledge and performance in terms of medication administration were evaluated before and one month after the training program.

Findings: It was observed that before training, the knowledge of the nurses in control group (90.6%) and intervention group (80%) was in weak level. After training, the knowledge state of 97.1% of intervention group's nurses improved to good level. Before training, most of the nurses of both groups showed improper performance. However, training program improved the performance of the nurses in intervention group.

Discussion and conclusion: Some errors may happen during medication administration by feeding tubes which could result in undesirable complications. In this regard, improvement and application of correct medication administration techniques can result in increase of drugs' and diets' effectiveness, patients' safety and reduction of feeding tubes blockage.

Keywords: Small group training, nurse, endoscopic gastric tube, knowledge, gavage, nutrient-drug interaction

INTRODUCCIÓN

Patients are in need of extra care in intensive care unit (ICU) (Heldt T et al., 2013). Patients of ICU often can't fulfill their nutrient needs (Wright-Myrie D et al., 2013). In ICU, the patients are fed by feeding tubes (Zoni MLG et al., 2016).

In addition to food gavage, nurses can administer medication orders by feeding tubes, correct drug delivery by feeding tubes requires the knowledge on different forms of drugs, possibility of their administration by feeding tubes and proper use of the relevant techniques (Mota MLS et al., 2010). Some drug forms, such as coated tablets, can be crushed and administrated by feeding tubes. But the slow-release tablets or entric-coated ones can't be crushed. Crushing of these types of tablets will destroy their properties and can be effective in therapy failure or toxicity (Dashti-Khavidaki S et al., 2012). Crushed tablets are the common reasons of feeding tubes blockage (Seifert C et al., 2005; Belknap D et al., 1997). Such blockage can occur in 8.3% of cases (Matsuba C et al., 2007). Possible interactions can also destroy the medication or food function. Health care team should be cautious and evaluate the possible nutrient-medication interactions (Williams NT et al., 2007).

In the study of Zhu et al, 60% of nurses had adequate knowledge on medication forms and 30% of them were aware of medication crushing or opening. 80% of the nurses administrated all medications of the patient simultaneously through nasogastric tube by the same syringe (Zhu L-L et al., 2012).

ICU staffs should have strong knowledge and scientific background (Mohamadi G et al., 2009). One of the methods to enhance the knowledge and reinforce the skills is training. Small group training is one the effective educational routes (Safari M et al., 2004).

Small group training is an organized and regular dialogue between a limited number of people who have common and mutual objectives (Fischer RL et al., 2004). This educational method let people discuss about their thoughts and experiences and the learners will have a lot of opportunities for thinking, analysis and evaluate their new knowledge and solve the clinical problems. In this regard, this method can enhance the critical thinking of the learners (Sanasuttipun W et al., 2009).

Regarding the effects of incorrect medication administration by feeding tubes and low knowledge of nurses, the present study was conducted with the aim of investigating the effectiveness of small group training on medication administration performance of the nurses through endoscopic gastric tube intubation in ICU of Imam Khomeini Hospital of Urmia in 2014.

MATERIALS AND METHODS

This is a clinical trial by which the effect of training on nurses' performance in nutrient-medication administration by feeding tubes was investigated. After obtaining Urmia Medical Science ethnical committee approval (umsu.rec.1393.36), the researcher referred to Imam Khomeini Hospital of Urmia for coordination with hospital authorities. For lowering the costs and also due to the small population volume, the sampling was done by census sampling and all the ICU nurses of Imam Khomeini Hospital were included in the study. In 2014, 92 nurses were working in ICU of Imam Khomeini Hospital of Urmia city. Regarding the inclusion (clinical nurses of ICU and all nurses with bachelor and master degrees in any nursing branches) and exclusion (fixed morning-shift nurses and those who had passed the medication administration management course in past 6 months) criteria, 80 nurses were selected. Then they were randomly classified into two 40-member groups (control and intervention).

Data collection was done by researcher-made questionnaires and checklists. The questionnaire included two sections. The first section was about demographic data and the second one involved 21 four-option questions about the research goals, which evaluated the nurses' knowledge on nutrient-medication administration. The checklist included 17 items assessing the nurses performance in terms of nutrient-medication administration. Their reliability was determined by content validation method.

For pretesting, the checklist was completed by the researcher in different shifts and then the questionnaires were filled up by the nurses in both groups. Then, for small group training, the intervention group was divided into four 10-member groups and the training was conducted by 4 sessions for each group (overall, 16 sessions were held). The sessions were 1 to 1.5 h long. During

						ranks		ranks	
Before	2.51±0.56	0.09	2.37±0.49	0.08	35.94	1258	31.88	1020	0.32
after	10.45±1.24	0.21	2.53±0.56	0.1	50	1750	16.5	528	0.000***

** P<0.05, 0.001

DISCUSSION

This study first addressed the primary research goal “determination of the nurses’ knowledge on nutrient-medication administration by endoscopic gastric feeding tube before and after small group training among two groups (intervention and control)”. For this purpose, the nurses’ knowledge on medication forms, medication preparation and administration and common nutrient-medication interactions was assessed by means of questionnaires. It was observed that before intervention, the knowledge of the nurses was weak (90.6% in control and 805 in intervention group). After the intervention, the knowledge status of the nurses in intervention group improved and 97.1% of them had good knowledge on the topic.

In the present study, for analysis of the second goal (investigation of the nurses’ performance in terms of nutrient-medication administration by endoscopic gastric feeding tube for ICU patients before and after small group training), the nurses’ performance was evaluated by a checklist. The results revealed improper performance of all the nurses in this term. Before intervention, 100% of the nurses of control and intervention groups administered the medication of the patients in combination with their food by means of a gavage, before intervention, 00% of the control group and 80% of the intervention group did not employ the earth gravity for gavage, these conditions improved by training. In a way that 82.9% of the nurses in intervention group used gavage for medication without combining them with the food.

Studies have shown that inadequate knowledge and improper performance of the nurses in this regard. Study of Dashti et al revealed that 50% of the nurses did not have sufficient knowledge on medication forms and medication administration by feeding tubes. Pharmacist presented an educational program including courses and manuals for the intervention group nurses. Educational program enhanced the knowledge of the nurses about the nutrient and medication administration and nutrient-medication interactions. In the intervention group, before training, 17.6% of the nurses washed

the tube. This percentage reached to 59.37% after training (Dashti-Khavidaki S et al., 2012).Bemt et al conducted a multidisciplinary study to reduce the errors in medication administration by feeding

tubes, this program included daily visit by the drugstore technician, putting the label of “Do not crush” on the drugs, establishment of a database on different forms of oral dosages, and accurate training of the nurses during their work. After the program, a significant decrease was observed in feeding tube blockage and errors of medication administration by the nurses. The main limitation of this study was lacking a control group (Bemt PMLAvd et al., 2006).

Regarding the mentioned reasons, the nurses’ errors could not be reduced only by training, as this study reported any changes in some items of checklist before and after the intervention. Application of alternative medication forms instead of solid tablets or using other routes for medication administration are some of these unchanged items. The responsibility for these items is not by the nurses and the physician is responsible. According to the studies, no proper professional relationship exists between the nurses and physicians. However, proper interaction of nurses and physicians can enhance the patient safety and improve the patient care quality. That’s why in spite of nurse knowledge about this field, the knowledge and skill exchange between the two groups are avoided due to improper nurse-physician interaction (Masror D et al., 2012; Shokri A et al., 2013; Azimi Lolaty et al., 2011). Also, the effective role of pharmacist in helping the physicians in drug prescription, nurses’ training and continuous monitoring of medication administration should not be neglected. Regarding the resources in this field, close collaboration of the therapy team including the physician, pharmacist and nurses, can enhance the drug administration by feeding tubes and prevention from their blockage, reduction of the errors in this regard and therefore increase of patients’ safety and decrease of hospitalization duration and costs.

REFERENCES

- Heldt T, Loss SH. Drug-nutrient interactions in the intensive care unit: literature review and current recommendations. *Rev Bras Ter Intensiva.* 2013;25(2):162-7.
- Wright-Myrie D, Kahwa E, Dover-Roberts D. Nutrition in critical illness: Critical care nurses’ knowledge and skills in the

- nutritional management of adults requiring intensive care – A review of the literature. *Caribbean Journal of Nursing*. 2013;1(1):49-55.
- Zoni MLG, Torre AD, Pires SL. Drugs and feeding tubes. *Rev Assoc Med Bras*. 2010;56(1):17-21.
- Mota MLS, Barbosa IV, Studart RMB, Melo EM, Lima FET, Mariano FA. Evaluation of Intensivist-Nurses' Knowledge Concerning Medication Administration Through Nasogastric and Enteral Tubes. *Rev Latino-Am Enfermagem*. 2010;18(5):888-94.
- Dashti-Khavidaki S, Badri S, Eftekharzadeh S-Z, Keshtkar A, Khalili H. The role of clinical pharmacist to improve medication administration through enteral feeding tubes by nurses. *Int J Clin Pharm*. 2012;34:757–64.
- Seifert C, Johnston B. A nationwide survey of long-term care facilities to determine the characteristics of medication administration through enteral feeding catheters. *Nutr Clin Pract June*. [abstract]. 2005;20(3):354-62.
- Belknap D, Seifert C, Petermann M. Administration of medications through enteral feeding catheters. *Am J Crit Care*. [abstract]. 1997;6(5):382-92
- Matsuba C, Gutiérrez MD, Whitaker I. Development and evaluation of standardized protocol to prevent nasoenteral tube obstruction in cardiac patients requiring enteral nutrition with restricted fluid volumes. *Clin Nurs*. [abstract]. 2007;16(10):1872-7.
- Williams NT. Medication Administration Through Enteral Feeding Tubes. *Am J Health Syst Pharm*. 2008;65(24):2347-57.
- Zhu L-L, Xu L-C, Wang H-Q, Jin J-F, Wang H-F, Zhou Q. Appropriateness of administration of nasogastric medication and preliminary intervention. *Therapeutics and Clinical Risk Management*. 2012;8:393-401.
- Mohamadi G, Ebrahimian A, Mahmoodi H. Evaluating the knowledge of intensive care unit nursing staffs. *IJCCN*. [In Persian]. 2009;2(1):41-6.
- Safari M, Salsali M, Ghofranipour F. The effect of nurses education with method Group discussion on the quality of nursing care in patients with Myocardial infarction. *JAUMS*. [In Persian]. 2004;2(4):437-41.
- Fischer RL, Jacobs SL, Herbert WNP. Small-Group Discussion Versus Lecture Format for Third-Year Students in Obstetrics and Gynecology. *Obstetricians and Gynecologists*. 2004;104(2):349-53.
- Sanasuttipun W, Tungjairob V, Musiksukont S, Lerthamatewe W, Chanwatana B. Effectiveness of Small Group Discussion on Critical Thinking, Self-directed Learning, and Learning Satisfaction of Thai Nursing Students. *J Nurs Sci*. 2009;27(3):8-16.
- Bemt PMLAvd, Cusell MBI, Overbeeke PW, Trommelen M, Dooren Dv, Ophorst WR, et al. Quality improvement of oral medication administration in patients with enteral feeding tubes. *Qual Saf Health Care*. 2006;15:44-7.
- Emami S, Hamishehkar H, Mahmoodpoor A, Mashayekhi S, Asgharian P. Errors of oral medication administration in a patient with enteral feeding tube. *Journal of Research in Pharmacy Practice*. 2012;1(1):37-40.
- Leape LL, Cullen DJ, Clapp MD, Burdick E, Demonaco HJ, Erickson JI, et al. Pharmacist Participation on Physician Rounds and Adverse Drug Events in the Intensive Care Unit. *JAMA*. 1999;282(3):267-71.
- Heydrich J, Heineck I, Bueno D. Observation of preparation and administration of drugs by nursing assistants in patients with enteral feeding tube. *Brazilian Journal of Pharmaceutical Sciences*. 2009;45(1):117-20.
- Bizhani M, Kouhpayeh SA, Abadi R, Tavacool Z. Effective factors on the Incidence of medication errors from the nursing staff perspective in various department of Fasa Hospital. *Journal of Fasa University of Medical Sciences*. [Research]. 2013;3(1):88-93.
- Rahimian Boogar I, Ghodrati Mirkouhi M. Role of workload, sleep, mental health and

individual factors in occurrence of nursing errors. Journal of Gorgan University of Medical Sciences. [Descriptive Study]. 2013;15(3):103-9.

Masror D, Heydarikhayat D, Joolae S. Assessing patient safety events and it's correlation with nurse - physician interaction from nurses' view. Quarterly Journal of Nersing Management. [Research]. 2012;1(2):37-45.

Shokri A, Yazdan Panah A, Vahdat S. The Professional Relationship between the Nurses and Physicians from their Own Point of View. 2. [Research]. 2013;15(1):76-69.

Azimi Lolaty H, Ashktorab T, Bagheri Nesami M, Bagherzadeh Ladari R. Experience of Professional Communication among Nurses Working in Educational Hospitals: A Phenomenological Study. Journal of Mazandaran University of Medical Sciences. [Research(Original)]. 2011;21(85):108-25