

RESEARCH ARTICLE

OPEN ACCESS

Influence of density on growth and survival of freshwater prawn Macrobrachium americanum (Bate, 1868) (Caridea: Palaemonidae) cultured in a cage-pond system

Jesus Ponce-Palafox^{1,2}, Fermín López-Uriostegui¹, Fabiola Lango-Reynoso¹, María Castañeda-Chávez¹, Itzel Galaviz-Villa¹, Jesus Montoya-Mendoza¹ and Héctor Esparza-Leal³

¹Doctoral Program in Aquaculture, National Technology of Mexico /Boca del Río Institute Technology, Road Veracruz-Córdoba Km. 12, Veracruz 94290, Mexico. ²Autonomous Nayarit University, Coastal Bioengineering Laboratory, Nayarit 63000, Mexico. ³National Polytechnic Institute, Interdisciplinary Research Centre for Regional Integral Development, San Joaquin, Guasave, Sinaloa 81101, Mexico.

Abstract

The domestication and culture of freshwater shrimp native of tropical and subtropical American Pacific zone requires the study and development of technologies in open systems where their response to types of farming, feeding and management strategies generate practical information for the production of this species. The aim of this study was to evaluate the effect of stocking density on growth and survival in *M. americanum* juvenile-adult, cage-cultured and to determine the optimal density for grow-out production. The caridean shrimp *Macrobrachium americanum* was cultured for 152 days in a 16 cage-culture (3 m³ each) at densities of 1, 3, 6 and 9 org/m³, respectively, with stocking sizes from 12.1 ± 1.7 to 13.5 ± 2.3 g. The prawns were fed twice daily with Camaronina 35%. Water quality parameters were within standard range for caridean shrimp culture. There was a significant effect of density on final weight, growth rate, specific growth rate, K condition and survival, and an inversely proportional relationship with the feed conversion ratio. The growth was affected by density, resulting to a maximum increase of 6 org/m³ in the asymmetry of the prawn and also increasing the number of small organisms by 9 org/m³. However, it is necessary to study such aspects as the rate and frequency of feeding, the initial size-grade and the implementation of shelter, among others.

Additional keywords: biomass; freshwater prawn; optimal growth; stocking size; shrimp.

Abbreviations used: CV (coefficient of variation); DO (dissolved oxygen); FCR (feed conversion ratio); GR (growth rate); HB (harvest biomass); PSI (production-size index); SD (standard deviation); SGR (specific growth rate); TAN (total ammonia nitrogen).

Authors' contributions: All authors contributed equally to this work: conception; acquisition, analysis, interpretation of data; drafting and critical revision of the manuscript.

Citation: Ponce-Palafox, J.; López-Uriostegui, F.; Lango-Reynoso, F.; Castañeda-Chávez, M.; Galaviz-Villa, I.; Montoya-Mendoza, J.; Esparza-Leal, H. (2018). Influence of density on growth and survival of freshwater prawn *Macrobrachium americanum* (Bate, 1868) (Caridea: Palaemonidae) cultured in a cage-pond system. Spanish Journal of Agricultural Research, Volume 16, Issue 4, e0508. https://doi.org/10.5424/sjar/2018164-13420

Received: 08 May 2018. Accepted: 23 Nov 2018.

Copyright © 2018 INIA. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC-by 4.0) License.

Funding: Instituto Tecnológico de Boca del Río; Universidad Autónoma de Nayarit, México.

Competing interests: The authors have declared that no competing interests exist.

Correspondence should be addressed to Jesus Ponce-Palafox: jesus.ponce@usa.net

Introduction

Problems associated with the deterioration of the environmental conditions of the hydrological basins in Latin America have raised concerns to implement systems for aquaculture production of native populations of prawns. Among the species with gradual decline and more aquaculture potential is *Macrobrachium americanum* (Ponce-Palafox *et al.*, 2002; Anger, 2013) among others. These species have the conditions of the key species in the lentic ecosystems, their size, and adaptation to aquaculture production systems, price, and their market demand. Many studies on biology, reproduction, physiology, biochemistry and growth aspects for caridean shrimp *M. americanum* have been conducted (Ponce-Palafox *et al.*, 2014). However, there are few studies on biotechnological aspects of growth in early juveniles (Mendez-Martinez *et al.*, 2017) and late juvenile-adults (Ponce-Palafox *et al.*, 2014). The generation of technology aspect for the culture of *M. americanum* juveniles, sub-adults and adults is essential for its growth and survival in commercial semiintensive and intensive farming systems.

The size and survival at stocking density have had a significant impact on production for the culture of crustaceans (Jones & Ruscoe, 2000), so it is necessary to study their behavior in aquaculture systems for different species. Numerous studies on freshwater prawns have shown the effect of density on growth, survival and production in ponds, tanks, cages and pens (Marques et al., 2000, 2012; Lopez-Uriostegui et al., 2014). Density in these species have been compared in mono, polyculture and mixed cultured conditions (Wohlfarth et al., 1985; Alam et al., 2001; Hossain & Islam, 2006; Uddin et al., 2008), suggesting that in the grow-out phase the best densities are between 2 and 6 org/m² depending on the species and the culture system. In *M. americanum*, the density effect in juveniles has been studied, obtaining 80% survival (Garcia-Guerrero & Apun-Molina, 2008; Ponce-Palafox et al., 2014). However, little is known as to the effect of density on *M. americanum* in grow-out phase in monoculture systems. The aim of this study was to evaluate the effect of stocking density on growth and survival in M. americanum juvenile-adult, cage-cultured and to determine the optimal density for grow-out production.

Material and methods

Site and cage system

The experiment was carried out in a 1,422 m² earthen pond located at the Centro de Acuicultura San Cayetano, Tepic, Nayarit Mexico $(21^{\circ}27'24.23''N; 104^{\circ}49'29.67''W)$, where 16 bottom cages $(3 \times 1 \times 1 m)$ were introduced in a semi-rustic rectangular pond. The cage system was used and managed according to what was described by Lopez-Uriostegui *et al.* (2014). The cages were constructed from polyethylene mesh (0.7 mm diameter) and iron frames. Circular feeding trays 50 cm in diameter, made of polyethylene screening were placed in the cages and synthetic mesh bags (raffia bags) were provided as refuges.

Water quality

Dissolved oxygen (DO), temperature (Yellow Springs Instruments YSI-85), pH (Bernauer F-1002) and water transparency (Secchi disk) were measured daily inside the cage at a depth of 30 cm. Samples of water were collected monthly inside the cages, to determine total ammonia nitrogen (TAN) and alkalinity (YSI 9100 photometer, Yellow Springs Instruments).

Experimental procedures

Shrimps of *M. americanum* (n=228) of 12.8 ± 1.63 g were placed in 16 cages with a stocking density of 1,

3, 6 and 9 org/m³, respectively, using a completely randomized design with four replicates by treatment. The prawns were fed twice daily with Camaronina 35% (Agribrands, Purina, Mexico, Inc., 35% protein, 8% lipid and 12% moisture), in the same proportion in each cage. The daily feed ratio was adjusted from 5% to 2% at the end of the experiment, based on feed demand and after periodic monitoring of the feed trays. All prawns were collected and weighed monthly to evaluate their growth and adjust the amount of feed supplied. After 152 days, the prawns growth was determined: mean initial weight (W_i) and mean final weight (W_f) in g, growth rate (GR) in g/week, production (P) in g/m³, harvest biomass (HB) in g/cage, specific growth rate (SGR) in %/day, production-size index (PSI), feeding conversion ratio (FCR) and survival (%). Biological parameters were determined as follows: GR = Final mean weight - Initial mean weight / Time; FCR = Feed fed / Weight gained; SGR = $100 \times [(\ln W_{e} - \ln W_{i}) /$ $(t_r - t_i)$; and PSI = (Production × Average weight) / 1000 (Tidwell et al., 1999).

Statistical analysis and model

The growth data were analyzed by Shapiro-Wilks and Bartlett tests, to assess normality and homoscedasticity, respectively. A paired t-test was used to compare the morning and afternoon data of water quality. The growth data were subjected to a one-way ANOVA (Montgomery, 2013). Differences among treatments were measured with Tukey's multiple comparison test of the means. The results were evaluated at 5% significance level. Survival data were square root arcsine transformed prior to analysis. The analyses were conducted using Statistica package v10 (StatSoft, Tulsa, OK, USA). The coefficient of variation was calculated as $CV = (Standard deviation / Mean) \times 100$.

A regression was applied for each stocking density *vs* production relationship of all animals (Bhujel, 2008). The relationship was a quadratic equation of the form:

$$Production = \beta_0 + \beta_1 d + \beta_2 d^2 \tag{1}$$

where β_2 is a quadratic coefficient (other than 0), β_1 is the linear coefficient, β_0 is the intercept and d is density (Bhujel, 2008).

The maximum density was obtained as follows:

$$d_{Max} = \frac{\beta_1}{2\beta_2} \tag{2}$$

where maximum production (*Production* $_{Max}$) is:

Production
$$_{\text{Max}} = \beta_0 + \frac{\beta_1^2}{2\beta_2} - \frac{\beta_1}{2}$$
 (3)

Final optimum production for stocking density was calculated from the first order derivative of the quadratic regressions (*i.e.* when $dP/dS_{density}=0$) (Buhjel, 2008).

Results

The water temperature, DO, pH and TAN concentrations did not show any difference (p>0.05) among different stocking densities (Table 1). Alkalinity concentration in the afternoon was significantly higher than in the morning (p<0.05).

M. americanum grew from a mean of 12.83 ± 0.64 g to 53.08 ± 29.28 g in 152 days at different stocking densities (Fig. 1; Table 2). After 60 days, differences were found between the densities of 1 to 3 org/m³ and 6 to 9 org/m³. Differences were also found between 1 and 3 org/m³ at 120 days (Fig. 1).

The final weight of the organisms decreased as the density increased from 82.4 to 16.2 g. The SD and CV were the highest in the density of 6 org/m³ with values of 20.7 and 47.0%, respectively. The organisms with the lowest growth had the lowest weight dispersion in the density of 9 org/m³ (Table 3).

The mean weight and survival of prawn at harvest $(82.4 \pm 9.1 \text{ g}; 100\%$, respectively) were significantly greater (p<0.05) at the lowest stocking density (1 org/m³), at 152 days (Table 2). Final weight, GR, HB, production, yield, K condition, SGR, and survival were the lowest in the highest density (9 org/m³). The prawn production and yield in 6 org/m³ were significantly higher than in other treatments (p<0.05) and reached 190.3 g/m³ and 1,903.0 kg/ha (152 days), respectively. FCR was significantly lower in stocking densities 1 and 3 org/m³, ranged from 1.4 to 1.6. In general, it was found

Table 1. Variations of water quality variables (mean±SD) in ponds used for rearing of *Macrobrachium americanum* at different stocking densities during a 152 days period.

Variables	Morning	Afternoon
Temperature (°C)	$27.0\pm2.4^{\rm a}$	$28.8\pm2.2^{\rm a}$
Secchi disk (cm)	35.0 ± 0.5	-
Dissolved oxygen (mg/L)	$6.2\pm1.6^{\rm a}$	$6.6\pm0.6^{\rm a}$
pH	$8.4\pm0.2^{\rm a}$	$8.5\pm0.1^{\rm a}$
Alkalinity (mg CaCO ₃ /L)	$110.0\pm0.90^{\text{b}}$	$135\pm1.3^{\rm a}$
Total ammonia N (mg/L)	$0.19\pm0.05^{\mathrm{a}}$	$0.15\pm0.08^{\mathrm{a}}$

Values with the same superscript within the same row are not significantly different at p<0.05.

that there was tendency to present a higher uniformity in the mean weight of the prawn in the treatment with the highest density (9 org/m³), the highest dispersion of weights of the population in stocking density 6 org/m³ and the highest weight in the treatment of 1 org/m³, with ranges from 74.3 to 96.8 g.

The quadratic regressions = $-8.0222x^2 + 78.629x + 17.54$ (R^2 = 0.9751; Fig. 2) indicated that the optimum for maximum production increases with stocking density up 3 org/m³ and was estimated to be 192.1 g/m³. Since that space does not increase in the cage, prawn production ceases when their standing crop reaches about 6 org/m³, and survival decreases from 100 to 92%.

Discussion

Water parameters were within range for *Macrobrachium* species (Boyd & Zimmerman, 2000; New, 2002) and those recorded in the culture of




Figure 1. Growth curves for *Macrobrachium americanum* at different stocking densities during a 152 days period. Errors bars: \pm SE.

$T3 = 6 \text{ org/m}^3$, $T4 = 9 \text{ org/m}^3$.	<i>c</i> ,	1	<i>, c</i>	, ,
Variables	T1	T2	Т3	T4
Initial weight (g)	$13.2\pm2.0^{\rm a}$	$12.5\pm1.9^{\rm a}$	$13.5\pm2.3^{\rm a}$	$12.1\pm1.7^{\rm a}$
Final weight (g)	$82.4\pm9.1^{\mathtt{a}}$	$69.6\pm7.7^{\rm b}$	$44.1\pm20.4^{\rm c}$	$16.2\pm1.9^{\rm d}$
Growth rate (g/week)	$3.2\pm1.2^{\rm a}$	$2.7\pm1.1^{\text{b}}$	$1.5\pm1.0^{\circ}$	$0.2\pm0.1^{\rm d}$
Harvest biomass (g/cage)	$247.2\pm9.7^{\text{b}}$	$576.3\pm11.7^{\rm a}$	$571.5\pm11.2^{\rm a}$	$236.2\pm8.9^{\circ}$
Production (g/m ³)	$82.4\pm12.7^{\rm b}$	$192.1\pm21.9^{\rm a}$	$190.5\pm18.5^{\rm a}$	$78.7\pm10.6^{\text{b}}$
Yield (kg/ha (152 days))	$824.3\pm92.4^{\rm b}$	$1{,}921.0\pm122.2^{\mathtt{a}}$	$1{,}905.1 \pm 115.7^{\rm a}$	$787.3\pm63.9^{\rm c}$
Specific growth rate (%/day)	$1.2\pm0.1^{\rm a}$	$1.1\pm0.2^{\rm a}$	$0.8\pm0.1^{\rm b}$	$0.2\pm0.1^{\circ}$
Production-size index	$67.9\pm6.9^{\circ}$	$133.7\pm12.1^{\mathtt{a}}$	$84.0\pm8.6^{\rm b}$	$12.8\pm1.6^{\rm d}$
Feeding conversion ratio	$1.4\pm0.1^{\text{c,b}}$	$1.6\pm0.3^{\rm b}$	$1.9\pm0.6^{\rm b}$	$2.6\pm0.7^{\rm a}$
K condition	$1.5\pm0.1^{\rm a}$	$1.4\pm0.1^{\rm a}$	$1.2\pm0.2^{\rm a,b}$	$0.9\pm0.1^{\rm b}$
Survival (%)	$100\pm2.1^{\mathtt{a}}$	$92\pm3.8^{\text{b}}$	$72\pm2.5^{\circ}$	$54\pm1.2^{\rm d}$

Table 2. Growth performance, production and survival of *Macrobrachium americanum* at different stocking densities during a 152 days period (mean \pm SD). T1= 1 org/m³, T2= 3 org/m³, T3= 6 org/m³, T4= 9 org/m³.

Values with the same superscript within the same row are not significantly different at p < 0.05).

stocking sizes in semi-natural ponds for *M. americanum* (Ponce-Palafox *et al.*, 2014). This study confirms that *M. americanum* can be grown in a grow-out phase in cage-cultured in a pond system at low densities, finding an inverse relationship between size and density, as it has been reported for this species (Ponce-Palafox *et al.*, 2014) and other native species of Latin America (Marques *et al.*, 2010; Lopez-Uriostegui *et al.*, 2014). The high growth rates of the prawn *M. americanum* in the cage-cultured in a pond system was due to the high availability of live food associated with periphyton, zooplankton and benthos (Uddin *et al.*, 2008; Marques *et al.*, 2010; Mohanty, 2010; Marques & Lombardi, 2011), and commercial feed with 35% protein in diet (Ponce-Palafox *et al.*, 2014).

When the density increased from 1 to 6 org/m³, the weight of the prawn decreased 50%, similar to what happens to *M. rosenbergii* in earthen ponds (Wohlfarth *et al.*, 1985). The positive correlation of FCR and density found showed that as the density increases, the efficiency of the feed decreases (Martin *et al.*, 1998). Stocking size affects the population structure, juvenile grading and selectively harvesting in prawns grown

Table 3. Characterization of the distribution of weights in *Macrobrachium americanum* at different stocking densities to the final weight (W_f) in a 152 days period. T1= 1 org/m³, T2= 3 org/m³, T3= 6 org/m³, T4= 9 org/m³.

8,	- 0) = -	8,	8
Parameters	T1	T2	Т3	T4
$W_{f-mean}(g)$	82.4	69.6	44.1	16.2
SD	9.1	7.7	20.7	1.9
CV	11.0	11.0	47.0	12.0
$W_{f-min}(g)$	74.3	63.0	15.5	13.0
$W_{f-max}(g)$	96.7	82.0	61.8	18.0

in cages (Karplus et al., 1986a; Lopez-Uriostegi et al., 2014). In this study it was determined that at low density, juveniles play smaller roles in the subsequent morphological development. This showed a structure similar to the ungraded population. This differs from that found in *M. rosenbergii* where juvenile jumpers are transformed into large prawns and laggards into small males (Karplus, 2005). However, this is similar to that found in M. amazonicus, where prawn development can show a directional process which produces a stable regulation through environmental factors (Maciel & Valenti, 2009). In 9 org/m³ there was a partial compensatory growth in this species, which has been reported for *M. rosenbergii* and other species in high densities (Daniels et al., 1995; Preto et al., 2010; Marques & Lombardi, 2011).

The coefficient of variation of weight was higher in 6 org/m³ (47.0%) because the larger prawns negatively affect the growth of the smallest ones and the structure of the population was more homogeneous in low densities. The CV (11%) of the densities of 1 to 3 org/m³ showed that in this species, the population tended to a "steady stable" of the morphotype distribution (D'Abramo *et al.*, 1989; Sampaio & Valenti, 1996). The CV showed that increasing stocking density up to 6 org/m³ increased production, but decreased commercial production by decrease mean weights.

At the density of 6 org/m³, strong density dependence was found at a high mortality and low growth rate. Strong density-dependence was established when growth in mean weight was high. The effect of "heterogeneous individual growth" (HIG) was greater at a density of 6 org/m³. The population was more complex at a density of 9 org/m³, and smaller sized transformation occurs without much increase to their body size, which




Figure 2. Relationship between production and survival of *Macrobrachium americanum* at different densities in the cage-culture in a pond system.

has been observed in other species (Ranjeet & Kurup, 2002). Space and natural food can be limiting in high densities, but the agonistic and social behavior increases disproportionately (Moraes-Valenti *et al.*, 2010).

In freshwater prawns stocked up to 6 org/m³, yields were significantly greater (p<0.05) than in 1 org/m³ (1,905.1 and 824.3 kg/ha (152 days), respectively). Also, weight and SGR were decreased (p<0.05) in prawns stocked at 6 and 9 org/m³. From 1 to 3 org/m³, higher values of SGR (1.1 to 1.2) were obtained than those recorded by Ponce-Palafox *et al.* (2014) for this species. However, the SGR was below those recorded for *M. rosenbergii* (3.86 to 4.96) in ponds (Tidwell *et al.*, 2004; Hossain & Islam, 2006).

The results obtained with *M. americanum* were in agreement with those found for *M. rosenbergii*, where the increase in production reduced the sizes at high density (Karplus *et al.*, 1986b). The increase in PSI with increased density up to 3 org/m³ indicated a positive balance in the production/size ratio. In this density, the production was more important than the low weight (Tidwell *et al.*, 2000). The values obtained in the density of 3 org/m³ (133.7 ± 12.1) were similar to those reported for *M. rosenbergii* at 133 days (Tidwell *et al.*, 2003), obtaining a high growth.

The quadratic regressions (Fig. 2) showed that the optimum stocking density considering the effect of production and survival (maximum standing crop) in semi-intensive systems was estimated to be 3 org/m³ or 192.1 g/m³. This density is within the range reported for high growth in grow-out phase in this species (Ponce-Palafox *et al.*, 2014).

Survival of *M. americanum* was inversely proportional to density, which is in agreement with that found in most of the monocultures of freshwater prawn (Valenti & New, 2000; Uddin *et al.*, 2008). The survival obtained at densities of 1 to 6 org/m³ was found within the range of 72 to 100%, suitable for commercial freshwater prawn culture (Karplus *et al.*, 1986b; Hosain & Islam, 2006; Ponce-Palafox *et al.*, 2014).

Our results indicate that *M. americanum* shows a density-dependent effect when the density was ≥ 6 org/m³ in grow-out, and mortality was high from 9 org/m³. The results of this research utilized bioengineering information on handling *M. americanum* in grow-out cages system. However, further research is required on the uniformity of size, feeding rates, and shelter, among others.

References

- Alam MJ, Jahan DA, Pramanikand WA, Hoq ME, 2001. Polyculture of freshwater prawn, *Macrobrachium rosenbergii*, de Man with carps: effects of prawn stocking density. Bangladesh J Fish Res 5 (2): 135-144.
- Anger K, 2013. Neotropical *Macrobrachium* (Caridea: Palaemonidae): on the biology, origin, and radiation of freshwater-invading shrimp. J Crustacean Biol 33 (2): 151-183. https://doi.org/10.1163/1937240X-00002124
- Bhujel RC, 2008. Statistics for aquaculture. Wiley-Blackwell Sci Publ, Ames, IA, USA.
- Boyd C, Zimmerman S, 2000. Grow-out systems- water quality and soil management. In: Freshwater prawn culture:

the farming of *Macrobrachium rosenbergii*; New MB & Valenti WC, eds., pp: 221-236. Blackwell Sci Inc., Malden, MA, USA. https://doi.org/10.1002/9780470999554.ch14

- D'Abramo LR, Heinen JM, Robinette HR, Collins JS, 1989. Production of the freshwater prawn *Macrobrachium rosenbergii* stocked as juveniles at different densities in temperate zone ponds. J World Aquacult Soc 20 (2): 81-89. https://doi.org/10.1111/j.1749-7345.1989.tb00527.x
- Daniels WH, D'Abramo LR, Fondren MW, Durant MD, 1995. Effects of stocking density and feed on pond production characteristics and revenue of harvested freshwater prawns *Macrobrachium rosenbergii* stocked as size-graded juveniles. J World Aquacult Soc 26: 38-47. https://doi.org/10.1111/j.1749-7345.1995.tb00207.x
- García-Guerrero MU, Apun-Molina JP, 2008. Density and shelter influence the adaptation of wild juvenile cauque prawns *Macrobrachium americanum* to culture conditions. N Am J Aquac 70 (3): 343-346. https://doi.org/10.1577/ A07-041.1
- Hossain MA, Islam MS, 2006. Optimization of stocking density of freshwater prawn *Macrobrachium rosenbergii* (de Man) in carp polyculture in Bangladesh. Aquacult Res 37: 994-1000. https://doi.org/10.1111/j.1365-2109.2006.01518.x
- Jones CM, Ruscoe IM, 2000. Assessment of stocking size and density in the production of redclaw crayfish, *Cherax quadricarinatus* (von Martens) (Decapoda: Parastacidae), cultured under earthen pond conditions. Aquaculture 189: 63-71. https://doi.org/10.1016/S0044-8486(00)00359-8
- Karplus I, 2005. Social control of growth in *Macrobrachium rosenbergii* (De Man): a review and prospects for future research. Aquacult Res 36 (3): 238-254. https://doi.org/10.1111/j.1365-2109.2005.01239.x
- Karplus I, Hulata G, Wolfarth GW. Halevy A, 1986a. The effect of size grading juvenile *Macrobrachium rosenbergii* prior to stocking on their population structure and production in polyculture. 1. Dividing the population structure into two fractions. Aquaculture 56: 257-270. https://doi.org/10.1016/0044-8486(86)90341-8
- Karplus I, Hulata G, Wohlfarth GW, Halevy A, 1986b. The effect of density of *Macrobrachium rosenbergii* raised in earthen ponds on their population structure and weight distribution. Aquaculture 52: 307-320. https://doi. org/10.1016/0044-8486(86)90373-X
- López-Uriostegui F, Ponce-Palafox JT, Arredondo-Figueroa JL, Benítez-Mandujano MA, García-Ulloa M, Castillo-Vargasmachuca SG, Esparza-Leal HM, 2014. Effect of stocking density on growth and survival of the prawn *Macrobrachium tenellum* cultured in a cage-pond system. N Am J Aquac 76: 164-169. https://doi.org/10.1080/1522 2055.2014.886646
- Maciel CR, Valenti WC, 2009. Biology, fisheries and aquaculture of the Amazon River prawn *Macrobrachium amazonicum*: a review. Nauplius 17 (2): 61-79.

- Marques HLA, Lombardi JV, 2011. Compensatory growth of Malaysian prawns reared at high densities during the nursery phase. R Bras Zootec 40 (4): 701-707. https://doi. org/10.1590/S1516-35982011000400001
- Marques HLA, Lombardi JV, Boock MV, 2000. Stocking densities for nursery phase culture of the freshwater prawn *Macrobrachium rosenbergii* in cages. Aquaculture 187: 127-132. https://doi.org/10.1016/S0044-8486(00)00304-5
- Marques HLA, Lombardi JV, Mallasen M, De Barros HP, Boock MV, 2010. Stocking densities in cage rearing of Amazon river prawn (*Macrobrachium amazonicum*) during nursery phases. Aquaculture 307: 201-205. https:// doi.org/10.1016/j.aquaculture.2010.07.035
- Marques HLA, Barros HP, Mallasen M, Boock MV, Valenti PMCM, 2012. Influence of stocking densities in the nursery phase on the growth of *Macrobrachium amazonicum* reared in net pens. Aquaculture 358-359: 240-245. https:// doi.org/10.1016/j.aquaculture.2012.06.011
- Martin JLM, Veran Y, Guelorget O, Pham D, 1998. Shrimp rearing: stocking density, growth, impact on sediment, waste output and their relationships studied through the nitrogen budget in rearing ponds. Aquaculture 164: 135-149. https://doi.org/10.1016/S0044-8486(98)00182-3
- Mendez-Martinez Y, Yamasaki-Granados S, García-Guerrero MU, Martinez-Cordova LR, Rivas-Vega ME, Arcos-Ortega FG, Cortes-Jacinto E, 2017. Effect of dietary protein content on growth rate, survival and body composition of juvenile cauque river prawn, *Macrobrachium americanum* (Bate 1868). Aquacult Res 48: 741-751. https://doi. org/10.1111/are.13193
- Mohanty RK, 2010. Impact of phased harvesting on population structure, feed intake pattern and growth performance of *Macrobrachium rosenbergii* De Man (giant freshwater prawn) in polyculture with carps in concurrent rice-fish culture. Aquacult Int 18: 523-537. https://doi.org/10.1007/ s10499-009-9263-5
- Montgomery DC, 2013. Design and analysis of experiments, 8th ed. Wiley, NY.
- Moraes-Valenti P, De Morais PA, Preto BL, Valenti WC, 2010. Effect of density on population development in the Amazon River prawn *Macrobrachium amazonicum*. Aquat Biol 9: 291-301. https://doi.org/10.3354/ab00261
- New MB, 2002. Farming freshwater prawns. A manual for the culture of the giant river prawn (*Macrobrachium rosenbergii*). FAO Fisheries Technical Paper No. 428. Food and Agriculture Organization of the United Nations, Rome, 212 pp.
- Ponce-Palafox JT, Arana-Magallón FC, Cabanillas-Beltrán H, Esparza-Leal H, 2002. Bases biológicas y técnicas para el cultivo de los camarones de agua dulce nativos del Pacífico Americano *Macrobrachium tenellum* (Smith, 1871) y *M. americanum* (Bate, 1968). I Congreso IberoamericanoVirtual de Acuicultura, CIVA 2002 (España), pp: 534-546.

- Ponce-Palafox JT, López-Uriostegui F, Arredondo-Figueroa JL, Benítez-Mandujano MA, García-Ulloa M, Esparza-Leal HM, Spanopoulos MH, 2014. Effect of stocking size on growth performance, biomass, production, yield, and survival of caridean shrimp cage-cultured in a pond system. N Am J Aquac 76: 430-435. https://doi.org/10.108 0/15222055.2014.936538
- Preto BL, Kimpara JM, Moraes-Valenti P, Valenti WC, 2010. Population structure of pond-raised *Macrobrachium amazonicum* with different stocking and harvesting strategies. Aquaculture 307: 206-211. https://doi.org/10.1016/j.aquaculture.2010.07.023
- Ranjeet K, Kurup BM, 2002. Heterogeneous individual growth of *Macrobrachium rosenbergii* male morphotypes. NAGA 25 (2): 13-18. https://www.worldfishcenter. org/content/heterogeneous-individual-growthmacrobrachium-rosenbergii-male-morphotypes-1
- Sampaio CMS, Valenti W, 1996. Growth curves for Macrobrachium rosenbergii in semi-intensive culture in Brazil. J World Aquacult Soc 27 (3): 353-358. https://doi. org/10.1111/j.1749-7345.1996.tb00619.x
- Tidwell JH, Coyle S, Webel C, Evans J, 1999. Effects and interactions of stocking density and added substrate on production and population structure of freshwater prawns *Macrobrachium rosenbergii*. J World Aquacult Soc 30 (2): 174-179. https://doi.org/10.1111/j.1749-7345.1999. tb00864.x
- Tidwell JH, Coyle SD, Van Arnum A, Weibel C, 2000. Production response of freshwater prawns *Macrobrachium rosenbergii* to increasing amounts of artificial substrate.

J World Aquacult Soc 31: 452-457. https://doi. org/10.1111/j.1749-7345.2000.tb00895.x

- Tidwell JH, Coyle SD, Bright LA, Arnum AV, Weibel C, 2003. The effects of size grading and length of nursery period on growth and population structure of freshwater prawns stocked in temperature zone ponds with added substrates. Aquaculture 218: 209-218. https://doi.org/10.1016/S0044-8486(02)00341-1
- Tidwell JH, Coyle SD, Dasgupta S, 2004. Effects of stocking different fractions of size graded juvenile prawns on production and population structure during a temperaturelimited growout period. Aquaculture 231: 123-134. https://doi.org/10.1016/S0044-8486(03)00544-1
- Uddin MS, Milsten A, Azim ME, Wahab MA, Verdegem M, Verreth J, 2008. Effects of stocking density, periphyton substrate and supplemental feed on biological processes affecting water quality in earthen tilapia–prawn polyculture ponds. Aquacult Res 39 (12): 1243-1257. https://doi.org/10.1111/j.1365-2109.2008.01987.x
- Valenti WC, New MB, 2000. Grow-out systems-monoculture. In: The farming of *Macrobrachium rosenbergii*; New MB, Valenti WC (eds.). pp: 177-186. Blackwell Sci, Oxford, UK. https://doi.org/10.1002/9780470999554. ch10
- Wohlfarth GW, Hulata G, Karflus I, Halevy A, 1985. Polyculture of the freshwater prawn *Macrobrachium rosenbergii* in intensively manured ponds, and the effect of stocking rate of prawns and fish on their production characteristics. Aquaculture 46: 143-156. https://doi. org/10.1016/0044-8486(85)90198-X