

SNP-SNP interactions in the *BDNF*, *COMT*, *CBR1* and *CCK* genes, associated with post-traumatic stress disorder in urban residents of Itagüí, Colombia

Interacciones SNP-SNP en los genes *BDNF*, *COMT*, *CBR1* y *CCK* asociadas al trastorno de estrés postraumático en la población urbana de Itagüí, Colombia

Mariana Duque-Quintero¹,  Juliana Martínez-Garro¹,  Pablo Andrés Guzmán-González¹,  Gloria María Sierra-Hincapié²,  Yolanda Torres-de Galvis² 

¹ Universidad CES - Department of Sciences and Biotechnology - Biology Academic Program - Medellín - Colombia.

² Universidad CES - Department of Medicine- CESIM Mental Health Study Center - Medellín - Colombia.

Corresponding author: Juliana Martínez-Garro. Department of Sciences and Biotechnology, Universidad CES. Calle 10A No. 22-04, block: B, floor: 6. Phone: +57 4 4548230, ext.: 1546. Medellín. Colombia. E-mail: jmartinezg@ces.edu.co.

Abstract

Introduction: Single nucleotide polymorphisms (SNPs) in the *BDNF*, *COMT*, *CBR1* and *CCK* genes have been associated with the process of fear extinction in humans. Since fear extinction plays a key role in recovering from psychological trauma, there is a possibility that these genes modulate the risk of developing post-traumatic stress disorder (PTSD).

Objective: To assess unilocus and multilocus associations between SNPs in the *BDNF*, *COMT*, *CBR1* and *CCK* genes and the risk of developing PTSD.

Materials and methods: 129 inhabitants of the municipality of Itagüí, Colombia, who had experienced psychological trauma at least once, were genotyped for these polymorphisms (38 cases of PTSD and 91 controls). Logistic regression was used to perform unilocus and multilocus association tests for single SNPs and existing SNP-SNP genotypic combinations.

Results: No unilocus associations were found, but interactions between the *BDNF* and *CBR1* genes and between the *COMT* and *CCK* genes were observed. Of these interactions, the genotypic combinations that behaved as risk factors were AG-AA (OR=13.52, $p<0.05$) in the *BDNF-CBR1* interaction, and TC-AA (OR=13.70, $p<0.05$) in the *CCK-COMT* interaction.

Conclusions: The two pairs of interacting polymorphisms found in this study could act additively and generate a greater risk of developing PTSD after suffering psychological trauma. People who have a single allele have a lower risk of developing PTSD than those who have two alleles in the interacting genes.

Keywords: Stress Disorders, Post-Traumatic; Extinction, Psychological; Polymorphism, Single Nucleotide; Risk Ratio (MeSH).

Resumen

Introducción. Los polimorfismos de un solo nucleótido (SNP, por su sigla en inglés) en los genes *BDNF*, *COMT*, *CBR1* y *CCK* han sido asociados con el proceso de extinción del miedo en humanos. Dado que la extinción del miedo es clave para la recuperación del trauma psicológico, es posible que estos genes modulen el riesgo de desarrollar trastorno de estrés postraumático (TEPT).

Objetivo. Evaluar las asociaciones unilocus y multilocus entre los SNP en los genes *BDNF*, *COMT*, *CBR1* y *CCK* y el riesgo de desarrollar TEPT.

Materiales y métodos. 129 habitantes del municipio de Itagüí, Colombia, que habían experimentado trauma psicológico al menos una vez, fueron genotipificados para estos polimorfismos (38 casos de TEPT y 91 controles). Se realizaron pruebas de asociación unilocus y multilocus por regresión logística para SNP únicos y las combinaciones genotípicas SNP-SNP existentes.

Resultados. No se encontraron asociaciones unilocus, pero se observaron interacciones entre *BDNF* y *CBR1*, y *CCK* y *COMT*. De estas interacciones, las combinaciones genotípicas que se comportaron como factores de riesgo fueron AG-AA (OR=13.52, $p<0.05$) de *BDNF-CBR1* y TC-AA (OR=13.70, $p<0.05$) de *CCK-COMT*.

Conclusiones: Los dos pares de polimorfismos en interacción encontrados en el presente estudio podrían actuar de forma aditiva y generar un mayor riesgo de desarrollar TEPT después de sufrir trauma psicológico. Quienes portan un solo alelo tienen un menor riesgo de desarrollar el trastorno que quienes portan dos alelos en genes que interactúan entre sí.

Palabras clave: Trastornos por estrés postraumático; Extinción psicológica; Polimorfismo de nucleótido simple; Factores de riesgo (DeCS).

Duque-Quintero M, Martínez-Garro J, Guzmán-González PA, Sierra-Hincapié GM, Torres-de Galvis Y. SNP-SNP interactions between *BDNF*, *COMT*, *CBR1* and *CCK* genes associated with post-traumatic stress disorder in urban residents of Itagüí, Colombia. Rev. Fac. Med. 2020;68(4):550-5. English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n4.77723>.

Duque-Quintero M, Martínez-Garro J, Guzmán-González PA, Sierra-Hincapié GM, Torres-de Galvis Y. [Interacciones SNP-SNP en los genes *BDNF*, *COMT*, *CBR1* y *CCK* asociadas al trastorno de estrés postraumático en la población urbana de Itagüí, Colombia]. Rev. Fac. Med. 2020;68(4):550-5. English. doi: <http://dx.doi.org/10.15446/revfacmed.v68n4.77723>.

Introduction

Post-traumatic stress disorder (PTSD) is a pathological response to stressful life events that can cause psychological trauma.¹ The prevalence of such events in conflicted societies facilitates the appearance of this mental health condition.² That is the case of Itagüí (Colombia), a city historically affected by violence, where an 11% PTSD rate among the adult population living in its urban area was reported in 2012.³ Besides environmental factors, genetic studies have shown that genetic inheritance plays a vital role in whether this condition is developed after being exposed to a trauma or not. In this sense, a twin study conducted on war veterans showed that genetic factors accounted for 30% of the risk of developing PTSD.⁴

According to the fear conditioning behavioral models that have been used to explain this pathology,⁵ PTSD symptoms can be triggered by exposing individuals who have experienced psychological trauma to stimuli associated with said traumatic event. However, in some cases, those associations remain active in time due to the inability of the individuals to inhibit the fear response and, therefore, PTSD symptoms may appear any time over a long period of time, even if the subject is in a safe environment.⁶ Persistent fear response is thought to be the result of the individual's failure to learn that said emotional response is no longer useful in a safe environment.⁷

It has been proposed that an individual's genotype can increase the risk of developing PTSD since it can influence how fear is processed in the underlying neural circuit after the subject has undergone a fear conditioning process. Also, this genotype could be associated with prefrontal cortex hypoactivity and hyperactivity in the amygdala;⁵ thus, it could be involved in how fear memory is consolidated in the corticolimbic pathway.⁸

SNP polymorphisms in genes that exert a certain degree of influence on different brain pathways have been associated with a fear extinction deficit, including *COMT*, which codes for the catechol-O-methyltransferase enzyme; *BDNF*, which codes for the brain-derived neurotrophic factor; *CBR1*, which codes for the cannabinoid receptor 1; and *CCK*, which codes for the cholecystokinin neuropeptide.

In the *COMT* gene, the *rs4680* single nucleotide polymorphism (SNP) yields a guanine for adenine substitution, which in turn leads to replacement of valine by methionine in the *COMT* enzyme, catalyzing the breakdown of catecholamines in the brain. In a fear conditioning experiment conducted on humans, those who had the Met/Met genotype showed a fear inhibition deficit.⁸ In this regard, Valente *et al.*⁹ in a study carried out with Brazilian victims of urban violence, reported that this polymorphism was associated with PTSD. In addition, the Val/Met mutation is also present in the *BDNF* protein due to a G/A substitution in SNP *rs6265* of the *BDNF* gene. Both mice and humans that carry the Met allele of the *BDNF* gene have exhibited deficit in fear extinction.¹⁰

On the other hand, in the *CBR1* gene, *rs2180619* SNP can have the A or the G allele. In this regard, Heitland *et al.*¹¹ reported that the participants with A allele extinguish fear less effectively than those with the G allele. Likewise, the *CBR1* gene has also been associated with PTSD.¹² Finally, an association between the T allele found in the SNP *rs1799923* in the *CCK* gene and

PTSD was reported in war veterans.¹³ It has been suggested that cholecystokinin (CCK) neuropeptide acts within the cannabinoid system to modulate fear inhibition; therefore, it is involved in fear processing.¹⁴

These findings allow suggesting that the combined effect of polymorphisms in the *BDNF*, *COMT*, *CBR1* and *CCK* genes may modulate the risk of developing PTSD after exposure to psychological trauma. In the present case-control study, carried out in the city of Itagüí, an association between PTSD and the aforementioned polymorphisms was tested to determine if these genes may modulate said risk. Both the unilocus and the multilocus effect over the phenotype were considered.

Materials and methods

Previous research

Patient data and genetic samples used in the present study were obtained from a previous mental health survey carried out in 2012 in Itagüí, Colombia.¹⁵ For that study, 896 people, living in six urban areas of this municipality, were administered the World Health Organization World Mental Health Composite International Diagnostic Interview (WHO WMH-CIDI). Their saliva samples were collected using the DNA Genotek's saliva-based collection kits. Chelex resins were used for DNA extraction. Finally, the obtained nucleic acids were stored in a Tris-EDTA buffer solution at -20°C.

Ethical considerations

All participants signed and submitted a written informed consent form in which they authorized the use of their DNA samples for future genetic studies on mental health. Data and sample collection procedures complied with the ethical principles for medical research established in the Declaration of Helsinki (2013)¹⁶ and the ethical principles for human research stated in Resolution 8430 issued by the Colombian Ministry of Health and Social Protection.¹⁷ This study was approved by the Bioethics Committee of Universidad CES on its session 83, held on July 13, 2015.

Participants

Out of the 896 subjects that participated in the study mentioned above, a sample of 129 respondents aged from 13 to 65 years was selected to conduct the present study. All participants from the original sample that had experienced psychological trauma at least once were included in the study. Cases (referred to as PTSD+) consisted of 38 people that were diagnosed with the disorder using the WHO WMH-CIDI. The criteria for diagnosis, according to the interview were: A) exposure to a traumatic event, B) persistently re-experiencing the event, C) persistent avoidance of stimuli associated with the trauma and numbing of general responsiveness, D) hyperarousal, E) experiencing disturbances B, C and D for more than a month, and F) clinically significant distress that impairs normal functioning.¹⁸ Controls (referred to as PTSD-) were made up of 91 subjects who only met the diagnostic criterion A.

Sex distribution was similar in both PTSD+ and PTSD- groups (42.1% and 40.6% men, respectively, $p > 0.5$, Fisher's exact test). On the other hand, distribution

of comorbidity with major depressive disorder (MDD) was significantly different between cases and controls (52.6% individuals diagnosed with MDD in the PTSD+ group vs. 13.2% in the control group, $p < 0.001$, Fisher's exact test). Finally, a higher prevalence of alcohol dependence was also observed in the cases (15.8% in the PTSD+ group compared to 1.1% in the control group, $p < 0.005$ Fisher's exact test).

Genotyping

SNP *rs6265* in the *BDNF* gene, SNP *rs4680* in the *COMT* gene, SNP *rs2180619* in the *CBR1* gene, and SNP *rs1799923* in the *CCK* gene were genotyped using competitive allele specific PCR (KASP™) (LGC Genomics, USA). Since it was not possible to find out the genotype of some participants, the following results were obtained: 113 genotypes for SNP *rs6265* (33 cases and 80 controls), 107 for SNP *rs4680* (32 cases and 75 controls), 109 for SNP *rs2180619* (32 cases and 77 controls), and 112 for SNP *rs1799923* (33 cases and 79 controls). Selection for double genotyping was blind and random, and only 12.16% of the samples genotyped were selected. The genetic results obtained after this process showed a 100% match between each sample and its replica. Negative controls were also processed blindly and, as expected, it was not possible to genotype them.

Statistical analysis

Statistical data processing was done using the R software version 3.2.2.¹⁹ The SNPAssoc package²⁰ was used to perform the Hardy-Weinberg test and the unilocus and multilocus analyses. The Epitools package²¹ was used to carry out a post-hoc analysis to study the allele combinations that could explain gene-gene

interactions. All alleles were found in the study population with a frequency greater than 1%. In addition, the Hardy-Weinberg Exact Test²² was used to evaluate the adjustment of the allele frequencies to the Hardy-Weinberg Model, which allowed determining that the allele frequencies of the four genes were in equilibrium.

The unilocus association with PTSD was assessed for each of the candidate genes. The Odds Ratio (OR) was calculated as a measure of risk, while significance was assessed with a maximum likelihood test. The Akaike information criterion (AIC) was used to judge which inheritance model (codominant, dominant, recessive, and over-dominant) fit the data better. The multilocus genetic association between every possible combination of SNP pairs was evaluated as an interaction in the logistic regression model. Age and sex were controlled in the unilocus and multilocus tests. Afterwards, Fisher's exact tests were performed to determine which genotypic combinations of the previously found interactions were significant as risk factors. This was made by comparing each genotypic combination to the rest of the existent genotypic combinations in the sample. Significance tests are reported with a 95% confidence interval.

Results

Unilocus analysis

Only the inheritance models that showed the best fit for the data (smallest AIC) were considered for the unilocus analysis. These inheritance models allowed determining the possible risk genotype for each independent gene. No significant association with post-traumatic stress disorder was reported in any SNP. However, OR intervals tended to be displaced to the right for the following genotypes: AG in the *BDNF* gene, AA in the *COMT* gene, and the TC in *CCK* gene (Table 1).

Table 1. Candidate genes as unilocus risk factors in PTSD cases.

Gene inheritance model	Genotypes compared	Allelic combinations	PTSD + *	PTSD - †	Odds Ratio	p-value‡
<i>BDNF</i> : Over-dominant model of inheritance	Risk genotype	AG	13	18	2.3 [0.9-5.4]	>0.05
	Reference	GG, AA	20	62		
<i>COMT</i> : Recessive model of inheritance	Risk genotype	AA	9	11	2.3 [0.8-6.4]	>0.1
	Reference	GG, AG	23	64		
<i>CBR1</i> : Dominant model of inheritance	Risk genotype	GA, GG	20	57	0.6 [0.2-1.4]	>0.1
	Reference	AA	12	20		
<i>CCK</i> : Over-dominant model of inheritance	Risk genotype	TC	12	17	2.1 [0.9-5.2]	>0.1
	Reference	TT, CC	21	62		

BDNF: brain-derived neurotrophic factor gene; *COMT*: catechol-O-methyltransferase; *CBR1*: cannabinoid receptor 1; *CCK*: cholecystokinin.

* Number of cases diagnosed with the disease.

† Number of controls.

‡ Probability.

Source: Own elaboration.

Multilocus analysis

When considering all possible combinations of SNP pairs in the regression model, interactions between *BDNF* and *CBR1* ($p < 0.05$) and between *CCK* and *COMT* ($p < 0.005$) were significant in the PTSD+ group. Then, existent ge-

notype combinations for these interactions were studied individually as risk factors and the following outcomes were obtained: when AG genotype of *BDNF* gene is combined with AA of *CBR1*, and when TC of *CCK* is combined with AA of *COMT*, the risk for developing PTSD increases significantly in comparison to other existent combinations (Table 2).

Table 2. Genotype pairs of SNP-SNP interactions identified as risk factors in PTSD cases.

SNP-SNP interaction	BDNF - CBR1	Genotype combinations	PTSD + *	PTSD - †	Odds Ratio	p-value ‡
	Risk Haplotype	AG-AA		5	1	13.5 [1.5-121.0]
Reference	GG-AA + GG-GA + GG-GG + AG-GA + AG-GG + AA-AA		27	72		
SNP-SNP interaction	CCK7 - COMT8	genotype combinations	PTSD + *	PTSD - †	Odds Ratio	p-value ‡
	Risk Haplotype	TC-AA	5	1	13.7 [1.5-122.7]	<0.05
Reference	TC-GA + TC-GG + CC-AA + CC-GA + CC-GG + TT-AA + TT-GA		27	74		

SNP: Single nucleotide polymorphism; *BDNF*: brain-derived neurotrophic factor gene; *CBR1*: cannabinoid receptor 1; *COMT*: catechol-O-methyltransferase; *CCK*: cholecystokinin.

* Number of cases diagnosed with PTSD.

† Number of controls.

‡ Probability.

Source: Own elaboration based on the data obtained in the study.

Discussion

Unilocus vs. multilocus results

According to the reports, risk allele A of the *BDNF*, *CBR1* and *COMT* genes and risk allele T of the *CCK* gene have been associated with increased vulnerability for PTSD^{9,13} or reduced capacity to inhibit the fear response.^{8,10,11} However, in the present study, no association between carrying these alleles, either in a homozygous or a heterozygous form, and developing PTSD as a result of being exposed to a traumatic event was found.

Nevertheless, when these alleles were portrayed in two different genes, they behaved as risk factors for developing the disease. This was the case for A allele as a heterozygous in genotype AG of *BDNF* combined with A allele as a homozygous in genotype AA of *CBR1*, and for T allele as a heterozygous in genotype TC of *CCK* combined with A allele as a homozygous in genotype AA of *COMT*. The low frequencies of genotype AA of *BDNF* (0.7%) and genotype TT of *CCK* (2.8%) that were observed in this study might explain why this effect is only present when these alleles appear as heterozygous. Thus, portraying two risk alleles in two different interacting genes increases the risk of developing PTSD after exposure to trauma.

SNP-SNP interactions associated with posttraumatic stress disorder

BDNF interacts with the endocannabinoid system inducing the synthesis and release of neuromodulators (endocannabinoids and neurotrophins). Once they are released, they activate their *CBR1* receptor, starting a long term depression of inhibitory synapses.²³ It is known that the substitution of valine for methionine in the *BDNF* precursor protein affects the depolarization dependent-secretion of this growing factor in hippocampal neurons.²⁴ If *BDNF* is secreted in a less effective way, its activity on the endocannabinoid system may be affected; this system plays important role in the expression and retention of fear extinction.²⁵ A minor activity of *BDNF*, together with portraying AA genotype in the SNP *rs2180619* of the *CBR1* gene might explain why an

association between these two interacting genes and PTSD was found in the present study.

Regarding the interaction between the gene products of *CCK* and *COMT*, it is known that the *CCK* neuropeptide exerts antagonistic effects on dopaminergic function through its action on the *CCK-B* receptor.²⁶ The *CCK* polymorphism under study appears to reduce the transcription of this gene.²⁷ Meanwhile, when the *COMT* enzyme is mutated with *Met* allele, its catalytic activity drops by 40%, increasing dopamine availability in the brain.²⁸ These two effects combined may impact the antagonistic effects of *CCK* over dopaminergic signaling, where the antagonistic effect of *CCK* is important to modulate behavior motivated by stress.²⁶ Yet, determining how a reduction of this modulatory action could be associated with PTSD requires further research.

Limitations

One of the limitations of this study is that the frequency of comorbidity with MDD was higher in the cases than in the controls. As a result, the genetic associations found here may be a consequence of an MDD diagnosis instead of a PTSD diagnosis. However, it is well known that comorbidity between MDD and PTSD is usually the standard. Depressive symptoms appear along with PTSD symptoms, and some authors consider that separating MDD into a different disorder when it co-occurs with PTSD is arbitrary.²⁹ Consequently, both diseases could share the genetic factors that increase the risk of developing them after an individual is exposed to a traumatic event.

Another limitation of this study is that the sample of PTSD patients retrieved from the previous study was low since only 38 people were diagnosed with the disorder out of 896 people that were originally interviewed. Thus, the presented results may need additional support by further studies.

Conclusions

This study shows that the probability of developing PTSD is higher when two risk alleles are present in two interacting genes than when the allele is only found in one

of the two genes. This means that to understand the genetic etiology of this disease better, the effect of multiple polymorphisms acting together on the phenotype should be considered. It is worth noting that the interactions found here should be analyzed at a biochemical level to understand how they influence the neural networks in which fear processing occurs and establish a causal relation between said influence and PTSD. Although the way how these genes interact at the neuronal level is not clearly known, it was found that certain genetic combinations, considered as risk factors, could be used to help achieving a genetic diagnosis of the disease.

Conflicts of interest

None stated by the authors.

Funding

This study was funded by a grant awarded by Universidad CES.

Acknowledgments

None stated by the authors.

References

- Segman RH, Cooper-Kazaz R, Macciardi F, Goltser T, Halfon Y, Dobroborski T, *et al.* Association between the dopamine transporter gene and posttraumatic stress disorder. *Mol Psychiatry*. 2002;7(8):903-7. <https://doi.org/bbwx8p>.
- Echeburúa E, de Corral P, Amor PJ. Evaluación del daño psicológico en las víctimas de delitos violentos. *Psicothema*. 2002;14(1):139-46.
- Torres-de Galvis Y, Agudelo-Martinez A, Sierra-Hincapié GM, Salas-Zapata C. Prevalencia de trastornos mentales en población general del municipio de Itagüí (Colombia), 2012. *Rev CES Med*. 2014;28(1):49-60.
- True WR, Rice J, Eisen SA, Heath AC, Goldberg J, Lyons MJ, *et al.* A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Arch Gen Psychiatry*. 1993;50(4):257-64. <https://doi.org/fv4vbb>.
- Johnson LR, McGuire J, Lazarus R, Palmer AA. Pavlovian fear memory circuits and phenotype models of PTSD. *Neuropharmacology*. 2012;62(2):638-46. <https://doi.org/cxbpsk>.
- Zoladz PR, Diamond DM. Current status on behavioral and biological markers of PTSD: a search for clarity in a conflicting literature. *Neurosci Biobehav Rev*. 2013;37(5):860-95. <https://doi.org/ffd2>.
- Gluck MA, Mercado E, Myers CE. Learning and memory: from brain to behavior. New York: Worth Publishers; 2008.
- Wendt J, Neubert J, Lindner K, Ernst FD, Homuth G, Weike AI, *et al.* Genetic influences on the acquisition and inhibition of fear. *Int J Psychophysiol*. 2014;98(3 Pt 2):499-505. <https://doi.org/f76ps6>.
- Valente NL, Vallada H, Cordeiro Q, Bressan RA, Andreoli SB, Mari JJ, *et al.* Catechol-O-methyltransferase (COMT) val158met polymorphism as a risk factor for PTSD after urban violence. *J Mol Neurosci*. 2011;43(3):516-23. <https://doi.org/crjhcc>.
- Soliman F, Glatt CE, Bath KG, Levita L, Jones RM, Pattwell SS, *et al.* A genetic variant BDNF polymorphism alters extinction learning in both mouse and human. *Science*. 2010;327(5967):863-6. <https://doi.org/d6pd4v>.
- Heitland I, Klumpers F, Oosting RS, Evers DJJ, Leon-Kenemans J, Baas JMP. Failure to extinguish fear and genetic variability in the human cannabinoid receptor 1. *Transl Psychiatry*. 2012;2(9):e162. <https://doi.org/f4d2mf>.
- Lu AT, Ogdie MN, Järvelin MR, Moilanen IK, Loo SK, McCracken JT, *et al.* Association of the Cannabinoid Receptor Gene (CNR1) With ADHD and Post-Traumatic Stress Disorder. *Am J Med Genet Part B Neuropsychiatr Genet*. 2008;147B(8):1488-94. <https://doi.org/bb38f7>.
- Badour CL, Hirsch RL, Zhang J, Mandel H, Hamner M, Wang Z. Exploring the association between a cholecystokinin promoter polymorphism (rs1799923) and posttraumatic stress disorder in combat veterans. *J Anxiety Disord*. 2015;36:78-83. <https://doi.org/ffd3>.
- Chhatwal JP, Gutman AR, Maguschak KA, Bowser ME, Yang Y, Davis M, *et al.* Functional interactions between endocannabinoid and CCK neurotransmitter systems may be critical for extinction learning. *Neuropsychopharmacology*. 2009;34(2):509-21. <https://doi.org/bsxvvg>.
- Torres de Galvis Y, editor. Primer estudio poblacional de Salud Mental, Itagüí 2012. Itagüí: Facultad de Medicina, Universidad CES; 2012.
- World Medical Association (WMA). WMA Declaration of Helsinki – Ethical principles for medical research involving human subjects. Fortaleza: 64th WMA General Assembly; 2013.
- Colombia. Ministerio de Salud. Resolución 8430 de 1993 (octubre 4): Por la cual se establecen las normas científicas, técnicas y administrativas para la investigación en salud. Bogotá D.C.; octubre 4 de 1993.
- U.S. Department of Health and human services. Appendix E: DSM-IV-TR Criteria for Posttraumatic Stress Disorder. In: Substance Abuse Treatment: Addressing the Specific Needs of Women. Rockville (MD): Substance Abuse and Mental Health Services Administration (US); 2009 [cited 2015 Apr 26]. Available from: <https://bit.ly/34iTCgk>.
- R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2015. Available from: <https://bit.ly/31sacb4>.
- González JR, Armengol L, Guinó E, Solé X, Moreno V. SNPassoc: SNPs-based whole genome association studies. 2014 [cited 2015 Apr 26]. Available from: <https://bit.ly/2Hpf8qc>.
- Aragon TJ, Fay MP, Wollschlaeger D, Omidpanah A. Package "epitools". 2012 [cited 2015 Apr 26]. Available from: <https://bit.ly/31vfnXR>.
- Wigginton JE, Cutler DJ, Abecasis GR. A Note on Exact Tests of Hardy-Weinberg Equilibrium. *Am J Hum Genet*. 2005;76(5):887-93. <https://doi.org/fs98gz>.
- Zhao L, Yeh MLW, Levine ES. Role for Endogenous BDNF in Endocannabinoid-Mediated Long-Term Depression at Neocortical Inhibitory Synapses. *eNeuro*. 2015;2(2). <https://doi.org/ffd4>.
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, *et al.* The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell*. 2003;112(2):257-69. <https://doi.org/bzxmdn>.
- Lafenêtre P, Chaouloff F, Marsicano G. The endocannabinoid system in the processing of anxiety and fear and how CB1 receptors may modulate fear extinction. *Pharmacol Res*. 2007;56(5):367-81. <https://doi.org/fpfw8k>.
- Rotzinger S, Bush DEA, Vaccarino FJ. Cholecystokinin Modulation of Mesolimbic Dopamine Function: Regulation of Motivated Behaviour. *Pharmacol Toxicol*. 2002;91(6):404-13. <https://doi.org/fqpfft>.

27. Nielsen FC, Pedersen K, Hansen TV, Rourke IJ, Rehfeld JF. Transcriptional regulation of the human cholecystokinin gene: composite action of upstream stimulatory factor, Sp1, and members of the CREB/ATF-AP-1 family of transcription factors. *DNA Cell Biol.* 1996;15(1):53-63. <https://doi.org/csf945>.
28. Chen J, Lipska BK, Halim N, Ma QD, Matsumoto M, Melhem S, *et al.* Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain. *Am J Hum Genet.* 2004;75(5):807-21. <https://doi.org/fs9sbc>.
29. O'Donnell ML, Creamer M, Pattison P. Posttraumatic stress disorder and depression following trauma: understanding comorbidity. *Am J Psychiatry.* 2004;161(8):1390-6. <https://doi.org/c3z7bm>.