



Mastitis, somatic cell count, and its impact on the quality of dairy-products ... An omission in Colombia?: A review

Mastitis, conteo de células somáticas y su impacto en la calidad de los productos lácteos ... ¿Una omisión en Colombia?: Revisión de literatura

Mastite e seu impacto na qualidade dos laticínios ... ¿Uma omissão na Colômbia?: Revisão de literatura

David S Riveros-Galán^{id}; Mónica Obando-Chaves*^{id}.

Facultad de Ingeniería Agronómica, Departamento de Producción y Sanidad Vegetal, Programa de Ingeniería Agroindustrial, Universidad del Tolima, Ibagué, Tolima (Colombia).

To cite this article:

Riveros-Galán DS, Obando-Chaves M. Mastitis and its impact on dairy-product quality ... An omission in Colombia?: A review. *Rev Colomb Cienc Pecu* 2021; 34(4): 241–253. DOI: <https://doi.org/10.17533/udea.rccp.v34n4a01>

Abstract

Mastitis is the most common disease in dairy herds and the main cause of economic losses in milk production worldwide. This inflammatory reaction of the mammary gland affects the quantity, composition and quality of milk produced and its suitability for the dairy industry. Despite of its importance, Colombia has no regulations on somatic cell count (SCC); that is, no official upper limits have been established for the dairy industry. The current quality-based payment system for raw milk does not encourage local producers to reduce the level of somatic cells. Consequently, Colombia is at a disadvantage compared to countries that include this parameter in their payment schemes and subscribe to international free trade agreements, affecting the competitiveness of the Colombian dairy sector. This article reviews the types of somatic cells, the microbiology of mastitis, its etiology and diagnosis, the changes that generate the composition of milk, and the impact of high SCCs on the quality of dairy products, such as yogurt, cheese, and milk powder. The final section offers a reflection on the problem of high SCCs in Colombia and the lack of regulations in this regard.

Keywords: *dairy; dairy industry; dairy cattle; dairy products; mammary gland; mastitis etiology; milk derivatives; milk products; milk quality; raw milk; somatic cell count; udder.*

Received: April 7, 2020; accepted: July 6, 2020

*Corresponding author. Calle 42 # 1-02, Barrio Santa Helena (Parte Alta), 564, Ibagué, Tolima, Colombia.

E-mail: mobando@ut.edu.co



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Resumen

La mastitis es la enfermedad más frecuente en los hatos lecheros y es la principal causa de pérdidas económicas en la producción de leche alrededor del mundo. Esta reacción inflamatoria de la glándula mamaria afecta la cantidad, composición, calidad y aptitud de la leche para procesamiento por la industria. Sin embargo, en Colombia no existe normatividad sobre el recuento de células somáticas, principal indicador de mastitis; es decir, no existen límites oficiales que sirvan de referencia para la industria lechera. El sistema de pago basado en calidad no incentiva al productor a disminuir el nivel de células somáticas. Esto coloca al país en desventaja frente a otros países que exigen este parámetro en su esquema de pago y con los cuales Colombia tiene tratados de libre comercio, afectando negativamente la competitividad del sector lechero colombiano. En este artículo se hace una revisión sobre los tipos de células somáticas, la microbiología de la mastitis, su etiología, el diagnóstico, los cambios que generan la composición de la leche y el impacto que tienen los altos recuentos de células somáticas sobre la calidad de algunos derivados lácteos como yogures, quesos y leche en polvo. En la parte final se presenta una reflexión sobre la problemática de los altos recuentos de células somáticas en Colombia y la falta de normatividad al respecto.

Palabras clave: *calidad de la leche; conteo de células somáticas; derivados lácteos; etiología de la mastitis; ganado de leche; glándula mamaria; industria láctea; leche cruda; lechería; productos lácteos; ubre.*

Resumo

A mastite é a doença mais frequente nos rebanhos leiteiros e é a principal causa de perdas econômicas na produção de leite em todo o mundo. Essa reação inflamatória da glândula mamária causa efeitos na quantidade, composição, qualidade e aptidão de leite processado pela indústria. No entanto, na Colômbia, não há regulamentação sobre a contagem de células somáticas até o momento, o principal indicador de mastite, ou seja, não há limites oficiais que sejam referência para a indústria leiteira. O sistema de pagamento baseado na qualidade não incentiva o produtor a diminuir o nível de células somáticas. Isso coloca o país em desvantagem em comparação com outros países que exigem esse parâmetro em seu esquema de pagamento e com quem a Colômbia possui acordos de livre comércio, afetando negativamente a competitividade do setor. Este documento analisa os tipos de células somáticas, a microbiologia da mastite, sua etiologia, diagnóstico, alterações na composição do leite e o impacto da alta contagem de células somáticas na qualidade do leite. Alguns derivados de leite, como iogurtes, queijos e leite em pó. Na parte final, é feita uma reflexão sobre o problema da alta contagem de células somáticas na Colômbia e a falta de regulamentação nesse sentido.

Palavras-chave: *contagem de células somáticas; derivados de leite; etiologia da mastite; gado leiteiro; glândula mamária; indústria de laticínios; laticínio; leite cru; produtos lácteos; qualidade do leite; úbere.*

Introduction

High somatic cell count (SCC) is an indicator of mastitis in dairy cattle and an indirect measure of the quality of raw milk for the dairy industry. Mastitis is the most common disease in dairy herds and is the leading cause of economic losses in dairy production worldwide (Hogeveen *et al.*, 2011). The composition of milk may be affected by multiple factors, such as breed, age, lactation stage, climate, and animal diet, among others. Mastitis, an inflammatory reaction of the mammary gland, is also known to affect the quantity, quality, and suitability of milk for the dairy industry. Beyond its implications for animal health and food safety, mastitis is a problem for the dairy industry because milk changes result in lower yields, decreased shelf life, poor quality of dairy products, and economic losses for the industry (Le Marechal *et al.*, 2011).

In Colombia, quality-based milk price was established in Decree #616 (Social Protection Ministry, 2006). However, SCC -an udder-health indicator and an indirect measure of raw milk quality- is not considered in this decree, meaning that no official reference limits are considered in the payment system. Consequently, producers do not feel encouraged to lower somatic-cell levels. This results in a disadvantage compared to countries that include this parameter in their payment schemes and have free trade agreements with Colombia, negatively affecting its competitiveness.

This article reviews the literature on the microbiology and etiology of mastitis, and how milk quality and safety are affected by high SCCs. This study seeks to draw attention to the vital importance of developing the dairy sector in Colombia, its effect on farmers, the dairy industry, the country's economy, and -of course- the consumers.

Somatic cells and mastitis

Somatic cells count in milk is an indicator of udder health as these cells protect the mammary gland from infections. The SCC is influenced by

many factors, such as animal species, lactation status, individual and environmental factors, and animal management (Pyörälä, 2003).

The four main types of somatic cells (leukocytes) are macrophages, polymorphonuclear neutrophil cells, lymphocytes, and epithelial cells. Mastitis can affect a single udder quarter or several of them. In addition to the udder immune function, somatic cells provide numerous endogenous enzymes (*e.g.*, lipases, oxygenases, proteases, glycosidases). Proteases and lipases have specific activity and, when released into milk, can influence dairy-industry processes related to the transformation of raw milk, negatively affecting the quality of dairy products, producing rancidity (Li *et al.*, 2014).

Normal values of somatic cells per udder quarter are between 68.000 and 187.000 cells/mL in bovine milk (Djabri *et al.*, 2002). In contrast, the corresponding values in sick animals are >200.000 cells/mL, varying according to the pathogen causing the infection (Ruegg and Pantoja, 2013). The main microorganisms responsible for mastitis are *Corynebacterium bovis*, *Escherichia coli*, *Staphylococcus aureus*, *Streptococcus uberis*, and *Streptococcus agalactiae* (Le Maréchal *et al.*, 2011). Different pathogens induce different symptoms associated with different types of mastitis. Mastitis can be subclinical or clinical, in different ranges, as shown in Table 1.

Table 1. Characteristics and symptoms of mastitis.

Mastitis type	Symptoms
<i>Subclinical</i>	Not apparent swelling of the mammary gland. Requires diagnostics. It's the most prevalent disease on herds.
<i>Mild clinical</i>	Minimal changes in udders, presence of small clots in milk.
<i>Moderate clinical</i>	Visible swelling on udder quarters, presence of milk clots or blood in milk. Absence of systemic disease.
<i>Severe</i>	General swelling of the udder with serious local and systemic symptoms.

Le Maréchal *et al.* (2011).

Legal limits for acceptable SCC levels in the

dairy industry vary by country. For example, in the European Union, if cow's milk has SCCs >200.000 cells/mL, the udder is considered infected. When the SCC is over 400.000 cells/mL, milk is deemed unsuitable for human consumption. SCC on the tank is 100.000 cells/mL in Germany, 500.000 cells/mL in Canada, and 750.000 cells/mL in the United States (Ruegg and Pantoja, 2013).

Milk with high SCCs also represents a significant risk to public health. This has been insufficiently studied in developed countries because most milk consumed in such countries undergoes some heat treatment. In Colombia, 43% of the milk produced goes to informal vendors (DNP, 2010) and is mostly used to make artisanal peasant cheese without any heat treatment or pasteurization, thus, presence of pathogens and toxins is highly probable, thereby posing a risk to human health. *Listeria monocytogenes* in “campesino” cheese — a popular fresh product made in Colombia with unpasteurized milk- has been reported (D_xl_{ue} *et al.*, 2018).

An increase in SCC is normally associated with a decrease in milk production in ruminant animals. After clinical mastitis, milk production decreases in the cow and will never return to its normal levels (before mastitis). The percentage of milk yield loss depends on the infecting pathogen (Le Marechal *et al.*, 2011).

Definition of mastitis

Mastitis can be defined as an inflammation of the mammary gland, caused by the growth of pathogenic microorganisms introduced either naturally by the breast canal or by physical means during milking or cleaning of the animal. As microorganisms multiply, they release toxic substances into the mammary gland causing vascular permeability of the tissue, leading to a compositional change in milk due to the entry of blood components such as blood serum, proteins, salts, and enzymes. In turn, casein and lactose production substantially decreases, as is the quality of fatty acids present in dairy fat. These

changes affect the taste and quality of milk and dairy products. From a clinical point of view, mastitis is a multifactorial disease, a combination of the glandular defense mechanisms, exposure to pathogens, and environmental factors present at the site (Welleberg *et al.*, 2002).

Mastitis etiology

The process of infection of the mammary gland can be due to several factors. The presence of dirt or feces in the udder, poor hygiene among the milking staff, poor washing of the nipples in the context of automatic milking, and poor milking practices, among others. Although more than 150 species of microorganisms are capable of infecting the mammary gland (Kuang *et al.*, 2009; Shome *et al.*, 2011; El-sayed *et al.*, 2017), including viruses, yeasts, mycoplasma, fungi, and bacteria, more than 95% of mastitis cases worldwide are caused by about 10 microbial groups (Table 2). *S. aureus*, *S. agalactiae*, and *S. dysgalactiae* are the microorganisms present in most mastitis cases produced by contagion. Among the microorganisms present in the environment, four groups can be identified: Gram-negative microorganisms (*E. coli*), *Corynebacterium*, *Mycoplasma*, and others, such as *Nocardia*, *Prototheca*, and some yeasts (Ranjan *et al.*, 2006; Vakkamäki *et al.*, 2017).

S. agalactiae and *S. dysgalactiae* are obligatory inhabitants of the mammary gland and very easily spread among healthy cows causing subclinical or intermittent mastitis (Mc Dougall, 2002). *S. aureus* is not an obligatory inhabitant of the mammary gland but can be easily transmitted by sick cows by cross-contamination of milkers or poor cleaning. Once established, *S. aureus* is difficult to eliminate (Divers and Peek, 2008).

Microorganisms present in the environment, such as *E. coli* or *Mycoplasma*, are transmitted horizontally between asymptomatic animals, either orally in lactating calves, by contaminated feces, or by nasal secretions of infected but asymptomatic animals (Punyapornwithaya *et al.*, 2010). Proliferation of such microorganisms

is linked to high temperatures, wet surfaces, contaminated bed materials (straw), and poor milking practices, among other factors. Coliforms can enter the breast canal and multiply rapidly in short periods (16 hours), causing an inflammatory response (Divers and Peek, 2008).

Prevalence of mastitis in herds has declined dramatically in recent years due to improved milking practices and health control of farms and animals. It should be noted that mastitis prevalence varies greatly from region to region depending on climate, temperature, livestock breed, and handling at milking sites (Ruegg, 2012; Reyes *et al.*, 2017).

Table 2. Pathogens causing mastitis.

Type of microorganisms	Relevant species	Transmission
Gram-positive cocci	<i>S. aureus</i>	Natural inhabitants of the mammary gland and epithelial tissue. Calves could be infected after drinking mastitic milk.
	<i>S. agalactiae</i>	
	<i>S. dysgalactiae</i>	
	<i>S. uberis</i>	
	Other negative coagulase staphylococci like:	
	<i>S. xylois</i>	
Gram-negative rods	<i>S. warneri</i>	Produced due to lack of proper hygiene of animals and their surroundings. As a rule of thumb any situation that increases coliforms population will increase the probability of an infection
	<i>S. simulans</i>	
	<i>Escherichia coli</i>	
	<i>Klebsiella</i> sp.	
	<i>Enterobacter</i> sp.	
Mycoplasma	<i>Pseudomonas</i> sp.	<i>Mycoplasma</i> is found naturally on the respiratory tract of animals. It is transmitted between animals by means of direct contact or by the farmworkers through cross contamination
	<i>Serratia</i> sp.	
	<i>M. bovis</i>	
	<i>M. californicum</i>	
	<i>M. dispar</i>	
	<i>M. bovirhinis</i>	
	<i>M. bovis genitalum</i>	
	<i>M. alkalescens</i>	
	<i>M. canadense</i>	
	<i>C. bovis</i>	
<i>Arcanobacterium pyogenes</i> (formerly known as <i>C. pyogenes</i>)		
Other	Yeast:	Diseases produced by these microorganisms are generated after a bacterial infection where an antibiotic therapy had no effect on these microorganisms causing opportunistic infections that can be fatal
	<i>Candida</i> spp.	
	Fungi:	
	<i>Aspergillus</i> sp.	
	Algae:	
<i>Prototheca zopfii</i>		
<i>Prototheca trispora</i>		

Adapted from Ruegg (2017), Carrillo-Casas and Miranda-Morales (2012), and Hillerton and Berry (2005).

The countries with the lowest incidence of mastitis in Latin America are Argentina and Chile, due in part to the implementation of education for animal handlers along with good agricultural practices and programs aimed at decreasing mastitis prevalence (Calvinho, 2019). In other Latin American countries, studies have examined dairy areas, such as the Arequipa or Huaira Basin in Perú (Velázquez and Vega, 2012), and the province of Córdoba in Argentina (Vissio, 2015). However, it is not easy to obtain up-to-date data on the incidence of mastitis in specific countries, making it difficult to design control policies at the governmental level (Table 3). In Colombia, mastitis incidence has been investigated in the traditional dairy basins, such as Boyacá (Calderón and Rodríguez, 2008; Andrade *et al.*, 2014) and Cundinamarca provinces (Martínez, 2006).

Diagnosis

Mastitis can be diagnosed as clinical mastitis -noticeable due to milk clots, inflammation, discoloration, and fever in the mammary gland- or subclinical mastitis, which is the most prevalent in herds. Subclinical mastitis is challenging to diagnose due to the absence of obvious symptoms, making it a risk factor for the animal and the quality of milk. It should be noted that mastitis can be acute or chronic, producing inflammation episodes that can last for months, leading to fibrotic udder (Nielsen, 2009).

The most effective way to diagnose mastitis is direct somatic cell counting; however, this methodology is not always feasible, because it is labor-intensive and must be done routinely. The gold standard test is somatic cell counting in the industry and the California Mastitis Test (CMT) in farms (Mollenhorst *et al.*, 2010). The latter relates to the amount of DNA present in a milk sample, which correlates with an approximate number of somatic cells; however, this number is not always comparable to that derived from other methods, such as cell counting (Blowey and Edmonson, 2010). Compared to the CMT, direct somatic cell counting is a better indicator of the likelihood of infection of an animal. Counts of somatic cells above 50.000 cells/mL result in economic losses for the producer, and counts above 500.000 cells/mL indicate a greater than 50% chance of infection in the animal. It is also known that an increase in the number of somatic cells in milk directly affects milk quality (Emanuelson *et al.*, 2009; Viguier *et al.*, 2009), as well as the quality of products derived from it. Alternative methods for early detection of mastitis are electrical conductivity (Addis *et al.*, 2016; 2017), lactose percentage (Frey *et al.*, 2013), lactate dehydrogenase (Chagunda *et al.*, 2006), and acute phase proteins (Shaheem *et al.*, 2016), which indicate the presence of somatic cells and could replace the direct count in the near future.

Table 3. Mastitis prevalence in different countries.

Country	Staphylococcus aureus (%)	Streptococcus agalactiae (%)	Environment Streptococcus spp	Negative coagulase Staphylococci	Environment pathogens (%)
Argentina	2.0	0.3	-	25.3	-
Hungary	32.5	-	12.8	41.0	6.8
Holland	-	-	-	10.8	-
United States (Wisconsin)	-	-	-	12.8	-
Canada	74.0	4.4	-	-	-
Germany	5.0	-	8.7	17.17	-

Adapted from Carrillo-Casas and Miranda-Morales (2012).

The use of molecular techniques such as real-time polymerase chain reaction (PCR) has also been used to determine pathogenic bacteria in milk (Shome *et al.*, 2011; Oikonomou *et al.*, 2012; Griffioen *et al.*, 2016; Vakkamäki *et al.*, 2017).

Changes in milk associated with high SCC

The main changes in milk caused by high SCCs are summarized in Table 4.

Table 4. Main changes in milk composition caused by mastitis.

Parameter	Change
Lactose	Decreased
Milk fat	Increased free fatty acids
Casein	Decreased
Serum proteins	Increased, particularly serum albumin and immunoglobulins
Chloride	Increased
Sodium	Increased
Calcium	Decreased
pH	Increased

Adapted from Bobbo *et al.* (2017).

It can be observed that blood serum proteins (mainly serum-albumins and immunoglobulins) increase, and α and β caseins decrease. There is no such trend for fatty acids; however, some studies report a decreased percentage and quality of fatty acids, with increased free fatty acids (Murphy *et al.*, 2016). Milk pH increases, and sodium, chlorine, and calcium levels decrease. Enzyme activity increases (i.e., plasmin, which is a native protease of milk with an optimal activity of 37 °C present in the udder even before the cow is milked), can hydrolyze α and β caseins. Lipases also are increased, as a result, there are increasing levels of free fatty acids and short-chain fatty acids, leading to defects in dairy products, such as rancidity (Jadhav *et al.*, 2016).

Mastitis affects the quality of pasteurized milk and reduces its shelf life. Raw milk with high SCCs (>700.000 cells/mL) generates taste and odor defects such as rancidity, bitter taste, and astringency in pasteurized milk (Ma *et al.*, 2000). These conditions are attributed to

increased lipolysis and proteolysis occurring in milk, which are noticeable between 14 and 21 days of cold storage, decreasing the shelf life of these products. Ma *et al.* (2000) observed an increase in the amount of total protein, fat, and casein in milk with high SCCs (comparing pre- and post-infection milk), but the increase was explained by a decrease of approximately 30% of the volume of milk produced by the animal. During mastitis, the synthesis of casein and fat in the mammary gland decreases. In Ultra High Temperature (UHT) milk, produced with high SCCs, there were also defects in taste and rancidity due to the growth of bacteria and the proteolysis and lipolysis of milk components (Murphy *et al.*, 2016).

Impact of high levels of somatic cells in yogurt. There are not many studies of high SCCs in yogurt; however, decreased shelf life of this product is due to the appearance of foreign flavors and texture changes during storage. Yogurts produced from milk with counts >800.000 cells/mL had sensory defects and a decrease in consistency after 20 days of storage at 5 °C, attributed to proteolysis generated by the enzymes present due to high SCC (Oliveira *et al.*, 2002).

Impact of high level of somatic cells in cheeses. In cheese products, the impact of high SCCs appears to vary depending on the type of cheese. In short, changes in coagulation properties, increased moisture content in cheese, and development of foreign flavors are the main defects (Le Márechal *et al.*, 2011).

Delay in coagulation times, along with poor curd firmness, low cheese yields, and poor final cheese quality, can be attributed to increased milk pH, decreased lactose, and degradation of casein fractions (Bobbo *et al.*, 2017). In acid coagulation cheeses, the fermentative process is affected by the antimicrobial components produced in mastitic milk, which alter the development of normal aromatic compounds in cheese. Low cheese yields due to high SCCs in milk have been found in Prato, Mozzarella, and Zamorano cheese. High levels of proteolysis in

maturation were detected in Swiss and Cheddar cheeses (Le Marechal *et al.*, 2011).

A recent study in Colombia investigated the impact of SCCs on the technological aptitude of milk and the physicochemical and sensory quality of peasant cheese. Milk samples were taken with a range of 150.000 to 1.200.000 cells/mL, and 30 peasant cheeses produced with this milk. The study found a decrease in the technological aptitude of milk with >200.000 cells/mL, increased coagulation time, and decreased yield. The sensory quality was affected by counts greater than 800.000 cells/mL (Vásquez *et al.*, 2014).

Impact of high levels of somatic cells in powdered milk. The impact of high SCCs is weaker in powdered products. Recently, high SCCs were found to increase particle size in skim milk powder and to decrease its uniformity, affecting its functional properties, such as solubility, dispersibility, and wettability (Sert *et al.*, 2016).

The problem of high somatic-cell levels in Colombia

Money losses due to mastitis have been calculated in studies carried out in the United States. Unfortunately, no studies have been conducted in Colombia to quantify the economic losses due to this disease. Annual livestock guild losses are even harder to calculate. On the other hand, the quality-based milk payment system in Colombia does not include SCCs; therefore, only some farmers are receiving money as bonuses upon the quality payment system. Colanta, a Colombian dairy company, recognizes bonuses for milk with <400.000 cel/mL and pays a higher price for milk with <200.000 cel/mL (Agrocolanta, 2020). For the farm, economic losses include decreasing milk yield, animal losses, increases in veterinary costs (assistance and antibiotics), management costs, etc. Additionally, the dairy industry loses money by receiving milk with high SCCs due to decreased shelf life, product yields (especially cheeses), and loss of sensory attributes. The dairy industry

cannot produce high-quality products unless it receives high-quality (microbiologically and compositionally) raw milk. Heat treatments do not improve milk received on the dairy platform; a more internationally competitive dairy sector can come about only with clear governmental support in terms of sectorial policies and commitment of dairy farmers.

The vast majority of milk produced in Colombia is processed by few companies: Colanta, Alpina, Alquería-Freskaleche, Parmalat, and Gloria Colombia (SIC, 2013). These companies have livestock development departments that strive to improve the quality of milk they receive and manage their quality standards based on Colombian regulations. Being close to the dairy basins, which have specialized livestock (for example, in Antioquia), they receive milk with good production standards (approx. 16.46 L of milk/cow/day) and good quality (<100.000 CFU/mL and 250.000 cells/mL on average; Barrios and Oliveira, 2013). However, other provinces, such as Sucre, do not have the same standards. Romero *et al.* (2018) reported between 530.000 and 624.000 cel/mL. Only 20.6% of the total raw milk samples tested had <400.000 cel/mL. Other researchers analyzed raw milk samples from a Colombian dairy company in 11 collection centers (*i.e.*, around Bogotá, the Middle Magdalena Valley, the Viejo Caldas region, the Antioquia province, and the Atlantic coast), reporting that 41.8% of the tanks had <400.000 cel/mL and only 11.03% had <200.000 cel/mL (Vásquez *et al.*, 2012). More recently, Aldana *et al.* (2019) found SCCs between 230.000 and 338.000 cel/mL in 27 farms in Anaime-Cajamarca (Tolima province).

Other countries that have included SCC in their quality-based milk payment system have markedly improved raw milk quality. Since 1995, Chile has applied a scheme of payment where SCC was incorporated to improve raw milk quality aiming to reach high quality standards. Statistics of some Southern Chile dairy industries show that average SCC diminished from 460,000 to 330,000 cell/mL from 1997 to 2000, and the same trend was observed from 2001 to 2011: average SCC

was 161,131 cell/mL of milk, while in 2017 raw SCC was 151,518 cell/mL (Sebastino *et al.*, 2020). In Brazil, Botaro *et al.* (2013) found that financial rewards induced changes in milk quality at the central dairy cooperative in south Brazil decreasing counts in bulk tank SCCs and total bacteria. Furthermore, SCC has gained international relevance for the dairy industry worldwide due to the inclusion of this item as a requisite for any export of dairy products and derivatives (<400.000 cel/mL). Therefore, many countries have established quality-based milk payment systems that include this feature (More, 2009).

In Colombia, 43% of milk is marketed informally. Provinces with high levels of informality (over 60%), such as Cauca, Huila, Tolima, Valle del Cauca, Sucre, Santander, Norte de Santander, Meta, and Casanare, where productivity per animal is low (4.5 L/cow/day) also have low raw milk quality (DNP, 2010). Artisanal cheese is the main product of milk informally marketed, and 60% of this cheese is used in bakery products (*e.g.*, *achiras*, *arepas boyacenses*, and *almojábanas*). The small factories processing this milk do not have good quality standards, and the National Agency for the Control of Food and Medicines (INVIMA) usually gives them unfavorable or pending concepts (FEDEGAN, 2013).

As long as the government, dairy farmers, industry, and other stakeholders do not agree to establish a fair payment based on quality, the quality of raw milk in Colombia will not improve. On one hand, The National Dairy Council, who advises the Colombian government, is questioned due to the prevalence of dairy industry members over farmers, generating unfair decisions related to the price of raw milk. On the other hand, livestock owners and farmer associations argue that they want an international market-tied fare (Contexto Ganadero, 2017; 2018). Thus, more effort is required from the government, research centers, universities, and stakeholders to improve the quality of raw milk produced in Colombia. It is important to emphasize that provinces with high informality must improve

their standards, productivity and quality of raw milk. This could lead to better economic returns, generating greater development in these regions and the sector as a whole. Studying and disseminating the impact of high SCCs on cheese yield in Colombia and on shelf life of dairy products could elicit greater attention from the government and the dairy industry to establish official limits for SCCs and its legislation as part of the quality payment systems.

Declarations

Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

Author contributions

The authors declare they have no conflicts of interest with regard to the work presented in this report.

Author contributions

Mónica Obando Chaves: Construction of the article and writing on the influence of mastitis on technological aptitude of milk. David Santiago Riveros Galán: Construction of the article and writing about mastitis microbiology and etiology.

References

- Addis M, Bronzo V, Puggioni G, Cacciotto C, Tedde V, Pagnozzi B, Locatelli C, Casula A, Curone G, Uzzao S, Moroni P. Relationship between milk cathelicidin abundance and microbiologic culture in clinical mastitis. *J Dairy Sci* 2017; 100(4): 2944–2953. DOI: <https://doi.org/10.3168/jds.2016-12110>
- Addis MF, Tedde V, Puggioni G, Pisanu S, Casula A, Locatelli C, Rota N, Bronzo V, Moroni P, Uzzau S. Evaluation of milk cathelicidin for detection of bovine mastitis. *J Dairy Sci* 2016; 99(10): 8250–8258. DOI: <https://doi.org/10.3168/jds.2016-11407>

Agrocolanta. Prevención y control de la mastitis y de las células somáticas –RCS, 2020. Available at: <http://www.agrocolanta.com/mejoramiento-de-la-calidad/prevencion-y-control-de-la-mastitis-y-las-celulas-somaticas-r-c-s/>

Aldana L, Fandiño C, García I, Mora J. Perfil composicional, higiénico y sanitario de la leche cruda en sistemas de producción ganadera en el Cañon de Anaime-Cajamarca. *Rev Colomb Cienc Pecu* 2019; 32(Supl). <https://revistas.udea.edu.co/index.php/rccp/article/view/340343>

Andrade R, Caro-Carvajal Z, Pulido M, Porras J, Vargas-Abella J. Prevalencia de bacterias causantes de mastitis en fincas lecheras de Toca (Boyacá, Colombia). *Rev Cienc Agron* 2014; 11(1): 47–53. DOI: <https://doi.org/10.19053/01228420.3487>

Barrios D, Oliveira M. Análisis de competitividad del sector lechero: Caso aplicado al norte de Antioquia Colombia. *Innovar-Rev Cienc Ad* 2013; 48:33–41. ISSN 2248-6968.

Blowey R, Edmonson P. Mastitis control on dairy herds. CABI. 2nd edition; 2010; 274.p. 155–157.

Bobbo T, Ruegg L, Stocco G, Fiore G, Giancesella M, Morgante M, et al. Association between pathogen specific cases of subclinical mastitis and milk yield, quality, protein composition, and cheese making traits in dairy cows. *J Dairy Sci* 2017; 100(6): 4868–4883. DOI: <https://doi.org/10.3168/jds.2016-12353>

Botaro B, Gameiro A, Santos M. Quality based payment program and milk quality in dairy cooperatives of Southern Brazil: an econometric analysis. *Scientia Agricola* 2013; 70(1): 21–26. DOI: <https://doi.org/10.1590/S0103-90162013000100004>

Calderón A, Rodríguez V. Prevalencia de mastitis bovina y su etiología infecciosa en sistemas especializados en producción de leche en el altiplano cundiboyacense (Colombia). *Rev Colomb Cienc Pec* 2008; 21: 582–589.

Calvinho LF. Mastitis bovina: evolución del control en Argentina y nuevos horizontes de investigación. *Anales de la Academia Nacional de Agronomía*

y Veterinaria 2017 (tomo LXX); 151–165. ISSN: 0327-8093. Available at: http://sedici.unlp.edu.ar/bitstream/handle/10915/87779/Documento_completo.pdf-PDFA.pdf?sequence=1&isAllowed=y

Carrillo-Casas M, Miranda-Morales E. Bovine mastitis pathogens: Prevalence and effects on somatic cell count. *Milk production – An up-to-date overview of animal nutrition, management and health*. 2012. 360.374. ISBN: 978-953-51-5322-1.

Chagunda G, Larsen T, Bjerring M, Ingvarsen K. L-lactate dehydrogenase and N-acetyl- β -D-glucosaminidase activities in bovine milk as indicators of non-specific mastitis. *J Dairy Res* 2006; 73(4): 431–440. DOI: <https://doi.org/10.1017/S0022029906001956>

Contexto Ganadero. Balance de los 5 años de la resolución 017 de 2012 para pago de leche. Available at: <https://www.contextoganadero.com/economia/balance-de-los-5-anos-de-la-resolucion-017-de-2012-para-pago-de-leche>

Contexto Ganadero. CNL insistirá en la revisión del pago de leche al productor. 2018. Available at: <https://www.contextoganadero.com/politica/cnl-insistira-en-la-revision-del-pago-de-leche-al-productor>

DNP (Departamento Nacional de Planeación). Política Nacional para mejorar la competitividad del sector lácteo colombiano. Documento Conpes 3675. 2010. Available at: <https://www.minagricultura.gov.co/ministerio/direcciones/Documents/d.angie/conpes%203675.pdf>

Divers T, Peek S. *Rebhun's diseases of dairy cattle*. St. Louis: Saunders Elsevier. 2008; 360–364. ISBN: 9780323396592.

Djabri B, Bareille N, Beaudeau F, Seegers H. Quarter milk somatic cell count in infected dairy cows: a meta-analysis. *Vet Res* 2002; 33(4): 335–357. DOI: <https://doi.org/10.1051/vetres:2002021>

Duque M, Duque S, Duque A. The popular dairy chain: A look from foodborne diseases (ETA) in Antioquia, Colombia (2008–2015). *Rev electron Vet* 2018; ISSN 1695-7504.

- El-sayed A, Awad W, Abdou N, Castañeda H. Molecular biological tools applied for identification of mastitis causing pathogens. *I J Vet Sci Med* 2017(5): 89–97. DOI: <https://doi.org/10.1016/j.ijvsm.2017.08.002>
- Emanuelson B, Berglund E, Strandberg. Relationship between somatic cell count and milk yield in different stages of lactation. *J Dairy Sci* 2009; 92: 3124–3133. DOI: <https://doi.org/10.3168/jds.2008-1719>
- FEDEGAN. Informalidad de la Cadena Láctea: Retos para su modernización. 2013. Available at: <http://www.contextoganadero.com/blog/informalidad-de-la-cadena-lactea-retos-para-su-modernizacion>
- Frey I, Peña J, Thomann A, Schwendener S, Perreten V. Genetic characterization of antimicrobial resistance in coagulase-negative staphylococci from bovine mastitis milk. *J Dairy Sci* 2013; 96(4): 2247–2257. DOI: <https://doi.org/10.3168/jds.2012-6091>
- Griffioen K, Hop G, Holstege M, Velthuis A, Lam, T. Dutch dairy farmers need for microbiological mastitis diagnostics. *J Dairy Sci* 2016; 99(7): 5551–5561. DOI: <https://doi.org/10.3168/jds.2015-10816>
- Hillerton J, Berry E. Treating mastitis in the cow – a tradition or an archaism. *J Appl Microbiol* 2005; 98(6): 1250–1255. DOI: <https://doi.org/10.1111/j.1365-2672.2005.02649.x>
- Hogeveen H, Huijps K, Lam TJGM. Economic aspects of mastitis: New developments. *New Zeal Vet J* 2011; 59:16–23. DOI: <https://doi.org/10.1080/00480169.2011.547165>
- Jadhav P, Tarate S, Bhuvana M, Das D, Shome B. Somatic cell count as a monitoring system for hygienic milk production in India: A review. *Asian J Dairy Food Res* 2016; 35(4): 270–277. DOI: <https://doi.org/10.18805/ajdfr.v35i4.6624>
- Kuang Y, Tani K, Synnott AJ, Ohshima K, Higuchi H, Nagahata H, Tanji Y. Characterization of bacterial population of raw milk from bovine mastitis by culture-independent PCR – DGGE method. *Biochem Eng J* 2009; 45: 76–81 DOI: <https://doi.org/10.1016/j.bej.2009.02.010>
- Le Marechal C, Thiéry R, Vautor E, Le Loir Y. Mastitis Impact on technological properties of milk and quality milk products: a review. *Dairy Sci Technol* 2011; 91: 247–282. DOI: <https://doi.org/10.1007/s13594-011-0009-6>
- Li N, Richoux N, Boutinaud M, Martin P, Gagnaire V. Role of Somatic Cells on Dairy Processes and Products: a review. *Dairy Sci Technol* 2014; 94: 517–538. DOI: <http://dx.doi.org/10.1007/s13594-015-0221-x>
- Ma Y, Ryan C, Barbano M, Galton M, Rudan A, Boor J. Effect of somatic cell count on quality and shelf life of pasteurized fluid milk. *J Dairy Sci* 2000; 83:264–274. DOI: [https://doi.org/10.3168/jds.S0022-0302\(00\)74873-9](https://doi.org/10.3168/jds.S0022-0302(00)74873-9)
- Martínez G. Comportamiento de la mastitis bovina y su impacto económico en algunos hatos de la Sabana de Bogotá, Colombia. *Rev Med Vet* 2006; 12: 35–55. DOI: <https://doi.org/10.19052/mv.2052>
- Mc Dougall S. Bovine mastitis: epidemiology, treatment and control. *New Zealand Vet J* 2002; 50(3): 81–84. DOI: <https://doi.org/10.1080/00480169.2002.36274>
- Mollenhorst H, Van der Tol P, Hogeveen H. Somatic cell count assessment at the quarter level or cow milking level. *J. Dairy Sci.* 2010; 93 (7): 3358–3364. DOI: <https://doi.org/10.3168/jds.2009-2842>
- More S. Global trends in milk quality: Implications for the Irish dairy industry. *Ir Vet J* 2009; 62(Supl):5–14. DOI: <https://doi.org/10.1186/2046-0481-62-S4-S5>
- Murphy S, Martin N, Barbano M, Wiedman M. Influence of raw milk quality on processed dairy products: How do raw milk quality test results relate to product quality and yield? *J Dairy Sci* 2016; 99(12): 10128–10149. DOI: <https://doi.org/10.3168/jds.2016-11172>

- Nielsen C. Economic impact of mastitis in dairy cows. *A U Agr-Suecia*, 2009. ISBN 978-91-86195-76-2. Available at: https://pub.epsilon.slu.se/1968/1/Christel_Nielsen_kappa.pdf
- Oikonomou G, Machado V, Santisteban C, Schukken Y, Bicalho C. Microbial diversity of bovinemastiticmilkasdescribedbypyrosequencing of metagenomic 16s rDNA. *Plos One* 2012; 7: 10. DOI: <https://doi.org/10.1371/journal.pone.0047671>
- Oliveira CAF, Fernandes A, Cunha OC, Laranja L, Telles E, Carvalho S. Composition and sensory evaluation of whole yogurt produced from milk with different somatic cell counts. *Aust J Dairy Technol* 2002; 57:192–196. ISSN: 0004-9433
- Pyörälä S. Indicators of inflammation in the diagnosis of mastitis. *Vet. Res.* 34; 2003; 565–578. DOI: <https://doi.org/10.1051/vetres:2003026>
- Punyapornwithaya V, Fox L, Hancock D, Gay J, Alldredge J. Association between an outbreak strain causing *Mycoplasma bovis* mastitis and its asymptomatic carriage in the herd: A case study from Idaho, USA. *Prev Vet Med* 2010; 93: 66–70. DOI: <https://doi.org/10.1016/j.prevetmed.2009.08.008>
- Ranjan R, Swarup D, Patra C, Nandi D. Bovine protothecal mastitis: a review. *CAB Reviews*. 2006; 17: 1–8. DOI: <https://doi.org/10.1079/PAVSNR20061017>
- Reyes J, Chaffer J, Rodriguez-Lecompte C, Sánchez J, Zadoks N, Robinson N, Cardona X, Ramírez N, Keefe G. Short communication: Molecular epidemiology of *Streptococcus agalactiae* differs between countries. *J Dairy Sci* 2017; 100(11): 9294–9297. DOI: <https://doi.org/10.3168/jds.2017-13363>
- Romero A, Calderón A, Rodríguez V. Evaluation of raw milk quality in three subregions of the department of Sucre, Colombia. *Recia* 2018; 10(1): 43–50. DOI: <https://doi.org/10.24188/recia.v10.n1.2018.630>
- Ruegg P. A 100-Year Review: Mastitis detection, management, and prevention. *J Dairy Sci* 2017; 100 (12): 10381–10397. DOI: <https://doi.org/10.3168/jds.2017-13023>
- Ruegg P. Mastitis in dairy cows. *Vet Clin N Am-Food* 2012; 28(2): 227–229. DOI: <https://doi.org/10.1016/j.cvfa.2012.04.003>
- Ruegg P, Pantoja J. Understanding and using somatic cells count to improve milk quality. *Irish J Agr Food Res* 2013; 52: 101–117. DOI: <http://repositorio.unifesp.br/11600/44140>
- Sebastino K, Uribe H, González H. Effect of test year, parity number and days in milk on somatic cell count in dairy cows of Los Ríos region in Chile. *Austral journal of veterinary sciences* 2020; 52(1):1–7. DOI: <https://dx.doi.org/10.4067/S0719-81322020000100102>
- Sert D, Mercan E, Aydemir S, Civelek M. Effect of milk somatic cell counts on some physicochemical and functional characteristics of skim and whole milk powders. *J Dairy Sci* 2016; 99: 5254–5264. DOI: <https://doi.org/10.3168/jds.2016-10860>
- Shaheem M, Tantary H, Nabi SU. A Treatise on bovine mastitis: disease and disease economics, etiological basis, risk factors, impact on human health, therapeutic management, prevention and control strategy. *Adv Dairy Res* 2016; 4:1–10. Available at: <https://www.longdom.org/open-access/a-treatise-on-bovine-mastitis-disease-and-disease-economics-etiologicalbasis-risk-factors-impact-on-human-health-therapeutic-manag-23-29-888X-1000150.pdf>
- Shome BR, Das Mitra S, Bhuvana M, Krithiga N, Velu D, Isloor S. Multiplex PCR assay for species identification of bovine mastitis pathogens. *J Appl Microbiol* 2011; 111(6): 1349–1356. DOI: <https://doi.org/10.1111/j.1365-2672.2011.05169.x>
- SIC (Superintendencia de Industria y Comercio). Análisis del mercado de la leche y derivados lácteos en Colombia (2008–2012). Delegatura de Protección de la Competencia. Colombia. 2013. Available at: https://www.sic.gov.co/recursos_user/documentos/promocion_competencia/Estudios_Economicos/Estudios_Economicos/Estudio_Sectorial_Leche1.pdf

- Social Protection Ministry. Decree #616, 2006. Reglamento técnico sobre los requisitos que debe cumplir la leche para consumo humano que se obtenga, procese, envase, transporte, comercializa, expendia, importe o exporte en el país. Available at: <https://www.ica.gov.co/getattachment/15425e0f-81fb-4111-b215-63e61e9e9130/2006D616.aspx>
- Vakkamäki J, Taponen S, Heikkilä A, Pyörälä S. Bacteriological etiology and treatment of mastitis in Finnish dairy herds. *Acta Vet Scand* 2017; 59: 1–9. DOI: <https://doi.org/10.1186/s13028-017-0301-4>
- Vásquez J, Loaiza E, Olivera M. The hygienic and sanitary quality of raw milk collected from different regions in Colombia. *Orinoquia* 2012; 16(2):13–23. ISSN electrónico 2011-2629.
- Vásquez J, Novoa C, Carulla J. Efecto del recuento de células somáticas sobre la aptitud quesera de la leche y la calidad fisicoquímica y sensorial del queso campesino. *Rev Fac Med Vet Zootec (Universidad Nacional de Colombia)* 2014; 61: 171–185. DOI: <https://doi.org/10.15446/rfmvz.v61n2.44680>
- Velázquez C, Vega J. Calidad de leche y mastitis subclínica en establos de la provincia de Huaura, Lima. *Rev Invest Vet-Perú* 2012; 23(1):65–71. Available at: DOI: http://www.scielo.org.pe/scielo.php?script=sci_arttext&pid=S1609-91172012000100008
- Viguiet C, Arora S, Gilmartin N, Welbeck K, O’Kennedy R. Mastitis detection: current trends and future perspectives. *Trends Biotechnol* 2009; 27: 486–492. DOI: <https://dx.doi.org/10.1016/j.tibtech.2009.05.004>
- Vissio C. Pérdidas productivas y económicas diarias ocasionadas por la mastitis y erogaciones derivadas de su control en establecimientos lecheros de Córdoba, Argentina. *Arch Med Vet* 2015; 47: 7–14. DOI: http://www.scielo.org.pe/scielo.php?script=sci_arttext&pid=S1609-91172012000100008
- Wellenberg J, Poel W, Van Der M, Oirschot J, Van T. Viral infections and bovine mastitis: a review. *Vet Microbiol* 2002; 88: 27–45. DOI: [https://doi.org/10.1016/S0378-1135\(02\)00098-6](https://doi.org/10.1016/S0378-1135(02)00098-6)