



On Pointwise Smooth Dendroids

SERGIO MACÍAS ✉

Universidad Nacional Autónoma de México, Instituto de Matemáticas, Ciudad
Universitaria, México.

Abstract. This is a survey paper on dendroids, smooth dendroids and mainly on pointwise smooth dendroids based on the work of J. J. Charatonik and C. Eberhart (dendroids and smooth dendroids) and S. T. Czuba (pointwise smooth dendroids). We present several characterizations of pointwise smooth dendroids, including one using the strict point \mathcal{T} -asymmetry property, defined by D. P. Bellamy.

Keywords: Aposyndetic continuum, continuum, dendrite, dendroid, pointwise smooth dendroid, Jones' set functions \mathcal{T} and \mathcal{K} , semi-aposyndetic continuum, smooth dendroid, strictly point \mathcal{T} -asymmetric continuum.

MSC2010: 54F15, 54F50, 54F55.

Sobre Dendroides Puntualmente Suaves

Resumen. Éste es un artículo expositivo sobre dendroides, dendroides suaves y, principalmente, sobre dendroides puntualmente suaves basado en el trabajo de J. J. Charatonik y C. Eberhart (dendroides y dendroides suaves) y S. T. Czuba (dendroides puntualmente suaves). Presentamos varias caracterizaciones de dendroides puntualmente suaves, incluyendo una utilizando la propiedad de \mathcal{T} -asimetría puntual estricta, definida por D. P. Bellamy.

Palabras clave: Continuo aposindético, continuo, dendrita, dendroide, dendroide puntualmente suave, funciones \mathcal{T} y \mathcal{K} de Jones, continuo semiaposindético, dendroide suave, continuo estrictamente puntualmente \mathcal{T} -asimétrico.

1. Introduction

This is a survey paper on dendroids, smooth dendroids and mainly on pointwise smooth dendroids based on the work of Janusz J. Charatonik and Carl Eberhart [2], [4] and [12] (dendroids and smooth dendroids) and of Stanisław T. Czuba [7], [9], [10], and [11] (pointwise smooth dendroids). We present several characterizations of pointwise smooth

E-mail: sergiom@matem.unam.mx ✉

Received: 25 October 2021, Accepted: 1 June 2022.

To cite this article: S. Macías, On Pointwise Smooth Dendroids, *Rev. Integr. Temas Mat.*, 40 (2022), No. 2, 137-158. doi: 10.18273/revint.v40n2-2022001

dendroids. In particular, a new one using the strict point \mathcal{T} -asymmetry property (defined by David P. Bellamy [19, p. 392]) and the nonexistence of R^3 -continua in the dendroid. We give the appropriate references for the results whose proofs we do not include.

The paper is divided in six sections. After this Introduction, in section 2, we have the preliminaries, here we include the definitions and some needed results for the rest of the paper. In section 3, we introduce dendroids and present some of their properties, for example that each subcontinuum of a dendroid is a dendroid (Theorem 3.1); that dendroids are hereditarily decomposable continua (Theorem 3.2); that being a dendroid is preserved by monotone maps (Theorem 3.4); we prove a characterization of monotone maps between dendroids (Theorem 3.5), and characterize dendrites as aposyndetic dendroids (Corollary 3.9). In section 4, we talk about smooth dendroids and prove that each subcontinuum of a smooth dendroid is a smooth dendroid (Theorem 4.2); that smoothness allows us to construct certain types of subcontinua (Theorem 4.6); and that smooth dendroids are contractible (Theorem 4.7). In section 5, we introduce the main topic of the paper, pointwise smooth dendroids, we show that a subcontinuum of a pointwise smooth dendroid is a pointwise smooth dendroid (Theorem 5.1); that a fan is pointwise smooth if and only if it is smooth (Theorem 5.3); we present a characterization of pointwise smooth dendroids using Professor F. Burton Jones' set function \mathcal{T} and irreducible subcontinua (Theorem 5.11); we give another characterization of pointwise smooth dendroids using the images of singletons using Professor Jones' set function \mathcal{K} (Theorem 5.19); we present some interrelationships of the set functions \mathcal{T} and \mathcal{K} on pointwise smooth dendroids (Theorem 5.22); we show a characterization of pointwise smooth dendroids using semi-aposyndesis and the nonexistence of R^3 -continua in the dendroid (Theorem 5.24). In section 6, we introduce strict point \mathcal{T} -asymmetry for continua, defined by David P. Bellamy [19, p. 392]. We present new results about strict point \mathcal{T} -asymmetry. We show that strict point \mathcal{T} -asymmetry is equivalent to semi-aposyndesis (Theorem 6.2); we prove that each pointwise smooth dendroid is strictly point \mathcal{T} -asymmetric (Corollary 6.3); we demonstrate that, for fans, smoothness, pointwise smoothness and strict point \mathcal{T} -asymmetry are equivalent properties (Corollary 6.6) and we give a new characterization of pointwise smooth dendroids using the strict point \mathcal{T} -asymmetry property and the nonexistence of R^3 -continua in the dendroid (Theorem 6.7).

2. Preliminaries

Let Z be a metric space. If A is a subset of Z , then $Int(A)$, $Bd(A)$ and $Cl(A)$ denote the interior, the boundary and the closure of A in Z , respectively. A *map* is a continuous function. The symbol \mathbb{N} denotes the set of positive integers.

A *continuum* is a nonempty compact, connected, metric space. A *subcontinuum* is a continuum contained in a metric space. A continuum X is *connected im kleinen* at a point p of X provided that for each open subset U of X containing p , there exists a subcontinuum W of X such that $p \in Int(W) \subset W \subset U$. A continuum X is *locally connected* at a point p if for every open subset U of X containing p , there exists a connected open subset V of X such that $p \in V \subset U$. A continuum X is *locally connected* provided that it is locally connected at each of its points. We do not define connectedness im kleinen globally because a continuum is connected im kleinen at each of its points if and only if it is locally connected [22, Theorem 1.4.18]. An *arc* is a continuum Z that

is homeomorphic to $[0, 1]$. If $h: [0, 1] \rightarrow Z$ is a homeomorphism, then $h(0)$ and $h(1)$ are the *endpoints* of Z .

Let X be a continuum and let p be a point of X . Then p is an *endpoint* of X provided that it is an endpoint of every arc in X that contains it. A *simple triod* is a continuum Z such that Z is the union of three arcs α_1, α_2 and α_3 such that $\alpha_1 \cap \alpha_2 = \alpha_1 \cap \alpha_3 = \alpha_2 \cap \alpha_3 = \{v\}$, where v is called the *vertex* of the simple triod.

A continuum X is *unicoherent* if each time A and B are subcontinua of X such that $X = A \cup B$, we have that $A \cap B$ is connected. A continuum X is *hereditarily unicoherent* provided that each of its subcontinua is unicoherent. Note that this is equivalent to say that a continuum is hereditarily unicoherent if the intersection of each pair of its subcontinua is connected.

Let X and Y be continua and let $f: X \rightarrow Y$ be a surjective map. We say that f is *monotone* provided that for each connected subset C of Y , we have that $f^{-1}(C)$ is a connected subset of X .

Theorem 2.1. *Let X and Y be continua and let $f: X \rightarrow Y$ be a map. If X is hereditarily unicoherent and f is monotone, then Y is hereditarily unicoherent.*

Proof. Suppose Y has a subcontinuum Z that is not unicoherent. Then there exist two subcontinua K and L of Z such that $Z = K \cup L$ and $K \cap L$ is not connected. Since f is monotone, $f^{-1}(Z), f^{-1}(K)$ and $f^{-1}(L)$ are subcontinua of X , and $f^{-1}(Z) = f^{-1}(K) \cup f^{-1}(L)$. Since $f^{-1}(K) \cap f^{-1}(L) = f^{-1}(K \cap L)$, we obtain that $f^{-1}(K) \cap f^{-1}(L)$ is not connected. Hence, X is not hereditarily unicoherent. \square

A continuum is *decomposable* if it can be written as the union of two of its proper subcontinua; it is *indecomposable* if it is not decomposable. A continuum is *hereditarily decomposable* provided that every nondegenerate subcontinuum of it is decomposable. A continuum is *hereditarily indecomposable* if each of its subcontinua is indecomposable.

Let X be a continuum and let \mathcal{G} be a family of subcontinua of X . Then \mathcal{G} is a *clump* if $\bigcup \mathcal{G}$ is a subcontinuum of X and there exists a subcontinuum C of X , called the *centre of the clump*, such that C is a proper subcontinuum of every element of \mathcal{G} , and for each pair G and G' of elements of \mathcal{G} , we have that $G \cap G' = C$.

A continuum X is *contractible* provided that there exist a point p and a map $F: X \times [0, 1] \rightarrow X$ such that, for each element x of X , $F((x, 0)) = x$ and $F((x, 1)) = p$.

Given a continuum X , we define its *n-fold hyperspace* as:

$$\mathcal{C}_n(X) = \{A \subset X \mid A \text{ is a nonempty closed subset of } X \text{ with at most } n \text{ components}\}.$$

We topologize $\mathcal{C}_n(X)$ with the topology given by the *Hausdorff metric* [21, Theorem 1.8.3]. With this topology, $\mathcal{C}_n(X)$ is an arcwise connected continuum [21, Theorem 1.8.12]. A *Whitney map* for $\mathcal{C}_n(X)$ is map $\mu: \mathcal{C}_n(X) \rightarrow [0, 1]$ such that $\mu(\{x\}) = 0$, for all points $x \in X$, $\mu(X) = 1$, and if A and B are two elements of $\mathcal{C}_n(X)$ and $A \subsetneq B$, then $\mu(A) < \mu(B)$ [24, (0.50)].

A continuum X is *semi-aposyndetic* provided that for each pair of points p and q of X , there exists a subcontinuum W of X such that $\{p, q\} \cap \text{Int}(W) \neq \emptyset$ and $\{p, q\} \cap (X \setminus W) \neq \emptyset$. A continuum X is *aposyndetic* if for each pair of points p and q of X , there exists a subcontinuum W of X such that $p \in \text{Int}(W) \subset W \subset X \setminus \{q\}$.

Given a continuum X and a point p of X , the *composant of p* in X , κ_p , consists of the union of all proper subcontinua of X that contain p .

Let X be a continuum and let $\{A_n\}_{n=1}^{\infty}$ be a sequence of nonempty closed subsets of X . The *limit inferior* of the sequence, denoted by $\liminf A_n$, is the set:

$$\{x \in X \mid \text{for each open subset } U \text{ of } X \\ \text{such that } x \in U, \text{ there exists } N \in \mathbb{N} \text{ such that} \\ U \cap A_n \neq \emptyset \text{ for all } n \geq N\}.$$

The *limit superior* of the sequence, denoted by $\limsup A_n$, is the set:

$$\{x \in X \mid \text{for every open subset } U \text{ of } X \\ \text{such that } x \in U, \text{ we have that } U \cap A_n \neq \emptyset \\ \text{for infinitely many indexes } n \in \mathbb{N}\}.$$

If there exists a subset A of X such that $\liminf A_n = \limsup A_n = A$, we say that the sequence $\{A_n\}_{n=1}^{\infty}$ *converges* to A .

Theorem 2.2. [2, Lemma 1] *Let X be a hereditarily unicoherent continuum. If $\{W_n\}_{n=1}^{\infty}$ is a sequence of subcontinua of X , then $\liminf W_n$ is a subcontinuum of X .*

The following result is useful for studying smooth dendroids.

Theorem 2.3. *Let X be a continuum and let p and q be two elements of X and let α be an arc contained in X whose endpoints are p to q . If U is an open subset of X such that $p \in U$ and $q \in X \setminus U$, then there exists an open subset V of X such that $Bd(V) \cap \alpha$ consists of just one point.*

Proof. Let U be an open subset of X such that $p \in U$ and $q \in X \setminus U$. Since X is a regular space, there exists an open subset W of X such that $p \in W \subset Cl(W) \subset U$. Since α is a connected set, $p \in W$ and $q \in X \setminus W$, we have that $\alpha \cap Bd(W) \neq \emptyset$ [18, Theorem 1, p. 127]. Let t be the first element of α in $Bd(W)$, from p to q . Since X is a completely normal space, there exist two disjoint open subsets Y and Z of X such that $pt \setminus \{t\} \subset Y$ and $tq \setminus \{t\} \subset Z$, where pt and tq are the subarcs of α , from p to t and from t to q , respectively. Let $V = Y \cap U$. Hence, $t \in Bd(V)$. Suppose that there exists a point s in $(Bd(V) \setminus \{t\}) \cap \alpha$. Then, we obtain that $s \in tq \setminus \{t\}$, because $pt \setminus \{t\} \subset V \subset Y$. This implies that $Y \cap Z \neq \emptyset$, a contradiction. Therefore, $\alpha \cap Bd(V) = \{t\}$. Also, $p \in V \subset Cl(V) \subset Cl(W) \subset U$. \square

Let X be a continuum. Define Professor F. Burton Jones set functions \mathcal{T} and \mathcal{K} as follows: if A is a subset of X , then

$$\mathcal{T}(A) = X \setminus \{x \in X \mid \text{there exists a subcontinuum} \\ W \text{ of } X \text{ such that } x \in \text{Int}(W) \subset W \subset X \setminus A\}.$$

Observe that for any subset A of X , $\mathcal{T}(A)$ is a closed subset of X and $A \subset \mathcal{T}(A)$. A continuum X is \mathcal{T} -additive provided that for each pair A and B of closed subsets of X , we have that $\mathcal{T}(A \cup B) = \mathcal{T}(A) \cup \mathcal{T}(B)$.

Next, if A is a subset of X , then

$$\mathcal{K}(A) = \bigcap \{W \mid W \text{ is a subcontinuum of } X \text{ and } A \subset \text{Int}(W)\}.$$

Note that for any subset A of X , $\mathcal{K}(A)$ is a closed subset of X and $A \subset \mathcal{K}(A)$.

The next theorem follows from the definition of the set function \mathcal{K} :

Theorem 2.4. *If X is a hereditarily unicoherent continuum and A is a nonempty closed subset of X , then $\mathcal{K}(A)$ is a subcontinuum of X .*

Theorem 2.5. [22, Theorem 7.7.2] *Let X be a continuum and let W be a subcontinuum of X . Then $\mathcal{K}(W) = \{x \in X \mid \mathcal{T}(\{x\}) \cap W \neq \emptyset\}$.*

Let X be a continuum and let K be a proper subcontinuum of X . Then K is an R^1 -continuum if there exist an open subset U of X , containing K , and two sequences $\{C_n^1\}_{n=1}^\infty$ and $\{C_n^2\}_{n=1}^\infty$ of components of U such that $\limsup C_n^1 \cap \limsup C_n^2 = K$.

Remark 2.6. In [22] an R^1 -continuum is called an R -continuum.

Let X be a continuum and let K be a proper subcontinuum of X . Then K is an R^3 -continuum if there exist an open subset U of X , containing K , and a sequence $\{C_n\}_{n=1}^\infty$ of components of U such that $\liminf C_n = K$.

From [8, Corollary 11], we have the following:

Lemma 2.7. *Let X be a dendroid. Then each R^1 -continuum contains an R^3 -continuum. Also, if an R^1 -continuum is a single point, then it is also both an R^2 and an R^3 -continuum.*

3. Dendroids

The class of dendroids was introduced by Bronisław Knaster in his seminar at the Institute of Mathematics of the University of Wrocław. We present properties of them.

A *dendroid* is an arcwise connected continuum that is hereditarily unicoherent; i.e., a dendroid is an arcwise connected continuum in which the intersection of every pair of its subcontinua is connected.

As an example of a dendroid, we have the *comb space*:




Figure 1: Comb Space

Theorem 3.1. *If X is a dendroid and Z is a subcontinuum of X , then Z is a dendroid.*

Proof. Let X be a dendroid and let Z be a subcontinuum of X . We only need to show that Z is arcwise connected. Let z_0 and z_1 be two points of Z . Since X is arcwise connected, there exists an arc α in X whose endpoints are z_0 and z_1 . Consider $\alpha \cap Z$. Since X is hereditarily unicoherent, $\alpha \cap Z$ is connected. Hence, since $\{z_0, z_1\} \subset (\alpha \cap Z)$, we obtain that $\alpha \cap Z = \alpha$. Therefore, $\alpha \subset Z$ and Z is a dendroid. \square

Theorem 3.2. *If X is a dendroid, then X is hereditarily decomposable.*

Proof. Let X be a dendroid. By Theorem 3.1, we only need to show that X is decomposable. Suppose that X is an indecomposable continuum. Let x_1 and x_2 be two points of X that belong to two distinct composants of X (indecomposable continua have uncountably many composants [16, Theorem 3-46] and they are pairwise disjoint [16, Theorem 3-47]). Since X is arcwise connected, there exists an arc α in X having x_1 and x_2 as its endpoints. Since x_1 and x_2 are in distinct composants of X and X is indecomposable, the only subcontinuum of X containing $\{x_1, x_2\}$ is X . Hence, $\alpha = X$. This is a contradiction, since an arc is a decomposable continuum. Therefore, X is hereditarily decomposable. \square

Notation. Let X be a dendroid. Since X is hereditarily unicoherent and arcwise connected, given to elements x_0 and x_1 of X , there exists a unique arc whose endpoints are x_0 and x_1 , this arc is denoted by $\overline{x_0x_1}$.

Theorem 3.3. [2, Corollary 1] *Let X be a dendroid, let $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ be two convergent sequences of elements of X . If $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ converge to a and b , respectively, then $\liminf \overline{a_nb_n}$ is a subcontinuum of X , and $ab \subset \liminf \overline{a_nb_n}$.*

Theorem 3.4. *Let X and Y be continua and let $f: X \rightarrow Y$ be a surjective map. If X is a dendroid and f is monotone, then Y is a dendroid.*

Proof. Suppose X is a dendroid and $f: X \rightarrow Y$ is a surjective monotone map. By Theorem 2.1, Y is hereditarily unicoherent. We need to show that Y is arcwise connected. Let y_0 and y_1 be two elements of Y . Since f is surjective, there exist two points x_0 and x_1 in X such that $f(x_0) = y_0$ and $f(x_1) = y_1$. Consider the arc $\overline{x_0x_1}$ in X . By [25, Theorem 8.14], we have that $f(\overline{x_0x_1})$ is a locally connected subcontinuum of Y containing $f(x_0)$ and $f(x_1)$. Since locally connected continua are arcwise connected [25, Theorem 8.23], the arc $\overline{y_0y_1}$ exists and is contained in Y . Therefore, Y is arcwise connected and it is a dendroid. \square

The following result gives a characterization of monotone maps between dendroids.

Theorem 3.5. *Let X and Y be dendroids and let $f: X \rightarrow Y$ be a surjective map. Then f is monotone if and only if $f(\overline{x_0x_1}) = \overline{f(x_0)f(x_1)}$ for each pair of points x_0 and x_1 of X .*

Proof. Suppose f is monotone and let x_0 and x_1 be two elements of X . Note that $\{f(x_0), f(x_1)\} \subset \overline{f(x_0)f(x_1)}$. Hence, $\overline{f(x_0)f(x_1)}$ is an arc contained in the subdendroid $f(\overline{x_0x_1})$ (Theorem 3.1). Since f is monotone, $f^{-1}(\overline{f(x_0)f(x_1)})$ is a subdendroid of X that contains $\{x_0, x_1\}$. Thus, $\overline{x_0x_1} \subset f^{-1}(\overline{f(x_0)f(x_1)})$. This implies that $f(\overline{x_0x_1}) \subset \overline{f(f^{-1}(\overline{f(x_0)f(x_1)}))} = \overline{f(x_0)f(x_1)}$. Thus, $f(\overline{x_0x_1}) = \overline{f(x_0)f(x_1)}$.

Now, assume that $f(\overline{x_0x_1}) = \overline{f(x_0)f(x_1)}$ for each pair of points x_0 and x_1 of X . Let y be an element of Y and let x_0 and x_1 be two elements of $f^{-1}(y)$. By our assumption, $f(\overline{x_0x_1}) = \overline{f(x_0)f(x_1)} = \{y\}$. Note that $\overline{x_0x_1} \subset f^{-1}(\overline{f(x_0)f(x_1)}) = f^{-1}(\{y\}) = \overline{f^{-1}(y)}$. Hence, $f^{-1}(y)$ is a connected subset of X . Thus, by [22, Theorem 1.4.46], f is a monotone map. \square

Important properties of dendroids are the following:

Theorem 3.6. [1, p. 18] *Let X be a dendroid. If $\xi: [0, \infty) \rightarrow X$ is a one-to-one map, then the closure of $\xi([0, \infty))$ is an arc ($\xi([0, \infty))$ is called a Borsuk ray).*

Theorem 3.7. [1, Theorem 2] *If X is a dendroid and $f: X \rightarrow X$ is a map, then there exists a point x in X such that $f(x) = x$.*

A *dendrite* is a locally connected dendroid. Dendrites constitute an important class of continua, the main properties of these spaces are given in [3].

Theorem 3.8. *A dendroid is a dendrite if and only if $\mathcal{T}(\{x\}) = \{x\}$, for all x in X .*

Proof. If X is a dendrite, by [22, Theorem 2.1.37], $\mathcal{T}(\{x\}) = \{x\}$, for all x in X .

Suppose X is a dendroid that satisfies that $\mathcal{T}(\{x\}) = \{x\}$, for all x in X . Let p be an element of X and let A be a nonempty closed subset of X such that $p \in \mathcal{T}(A)$. Since X is hereditarily unicoherent, X is \mathcal{T} -additive [22, Theorem 2.2.12]. Hence, by [22, Corollary 2.2.13], we have that $\mathcal{T}(A) = \bigcup\{\mathcal{T}(\{a\}) \mid a \in A\}$. Thus, there exists a point a in A such that $p \in \mathcal{T}(\{a\})$. Since $\mathcal{T}(\{a\}) = \{a\}$, we obtain that $p = a$, and $p \in A$. Hence, by [22, Corollary 2.1.31], X is connected im kleinen at p . Since p is an arbitrary point of X , we have that X is connected im kleinen at each of its points. Therefore, X is locally connected [22, Theorem 1.4.18], and X is a dendrite. \square

The next result says that for the class of dendroids aposyndesis is equivalent to local connectedness.

Corollary 3.9. *A dendroid is a dendrite if and only if X is aposyndetic.*

Proof. Let X be a dendroid. If X is a dendrite, by Theorem 3.8, we have that $\mathcal{T}(\{x\}) = \{x\}$, for all x in X . Hence, by [22, Theorem 2.1.34], X is aposyndetic. If X is an aposyndetic dendroid, we obtain that, by [22, Theorem 2.1.34], that $\mathcal{T}(\{x\}) = \{x\}$, for all x in X . Thus, by Theorem 3.8, X is a dendrite. \square

A *fan* is a dendroid with exactly one ramification point; i.e., a point that is the only endpoint in common of at least three otherwise disjoint arcs. This point is called the *top* of the fan. The cone over the harmonic sequence $\{\frac{1}{n}\}_{n=1}^{\infty} \cup \{0\}$ is an example of a fan, the so-called *harmonic fan*:




Figure 2: Harmonic Fan

4. Smooth Dendroids

The class of smooth dendroids was introduced by Janusz J. Charatonik and Carl Eberhart [4] and [12].

A dendroid X is *smooth at a point* p of X provided that for each sequence $\{x_n\}_{n=1}^{\infty}$ of points of X converging to a point x in X , the sequence of arcs $\{\overline{px_n}\}_{n=1}^{\infty}$ converges to the arc \overline{px} . The point p is called a *point of smoothness* of X . A dendroid X is *smooth* if it is smooth at some of its points. A fan is *smooth* provided that it is smooth at its top.

As an example of a smooth dendroid we have the comb space and the harmonic fan (Figures 1 and 2, respectively). Also, the cone over the Cantor set, called the *Cantor fan*, is a smooth dendroid:




Figure 3: Cantor Fan

As a consequence of [2, Theorem 9, p. 27] and [12, Corollary 4], we have:

Theorem 4.1. *A fan is smooth if and only if it can be embedded in the Cantor fan.*

Theorem 4.2. *If X is a smooth dendroid and Z is a subcontinuum of X , then Z is a smooth dendroid.*

Proof. Let X be a smooth dendroid, let Z be a subcontinuum of X , and let p be the point of smoothness of X . By Theorem 3.1, Z is a dendroid. If p belongs to Z , we are done. Assume that $p \in X \setminus Z$. Let z' be an element of Z and consider the arc $\overline{pz'}$. Let t be the first element of the arc $\overline{pz'}$ in Z . We show that Z is smooth at t . Let $\{z_n\}_{n=1}^\infty$ be a sequence of elements of Z converging to a point z in Z . Note that, for each $n \in \mathbb{N}$,

$$\overline{pz_n} = \overline{pt} \cup \overline{tz_n}.$$

Hence,

$$\liminf \overline{pz_n} = \overline{pt} \cup \liminf \overline{tz_n}$$

[17, p. 336], and

$$\limsup \overline{pz_n} = \overline{pt} \cup \limsup \overline{tz_n}$$

[17, p. 337]. Since X is smooth at p ,

$$\liminf \overline{pz_n} = \limsup \overline{pz_n}.$$

Thus,

$$\overline{pt} \cup \liminf \overline{tz_n} = \overline{pt} \cup \limsup \overline{tz_n}.$$

Since $\overline{pt} \cap \liminf \overline{tz_n} = \{t\}$ and $\overline{pt} \cap \limsup \overline{tz_n} = \{t\}$, we obtain that:

$$\liminf \overline{tz_n} = \limsup \overline{tz_n}.$$

Therefore, Z is smooth at t . ☑

Remark 4.3. An important characterization of smooth dendroids in terms of prohibited subdendroids is given in [15].

Theorem 4.4. [4, Corollary 10] *Let X and Y be dendroids and let $f: X \rightarrow Y$ be a surjective map. If X is smooth and f is monotone, then Y is smooth.*

Note that if X is a dendroid that is not smooth at the point p , then there exists a sequence $\{z_n\}_{n=1}^{\infty}$ of elements of X converging to a point z of X such that the sequence of arcs $\{\overline{pz_n}\}_{n=1}^{\infty}$ either does not converge or $\lim \overline{pz_n} \neq \overline{pz}$. The next theorem shows that we may always assume the second possibility.

Theorem 4.5. *Let X be a dendroid and let p be a point of X . If X is not smooth at p , then there exists a sequence $\{a_n\}_{n=1}^{\infty}$ of points of X converging to an element a of X such that the sequence of arcs $\{\overline{pa_n}\}_{n=1}^{\infty}$ converges to a subdendroid L of X and $\overline{pa} \neq L$.*

Proof. Suppose that X is not smooth at p . Let $\{a_n\}_{n=1}^{\infty}$ be a sequence of points of X that converges to an element a of X and the sequence $\{\overline{pa_n}\}_{n=1}^{\infty}$ does not converge. Since $\overline{pa} \subset \liminf \overline{pa_n}$, we have that there exists a subsequence $\{a_{n_k}\}_{k=1}^{\infty}$ of $\{a_n\}_{n=1}^{\infty}$ such that $\lim \overline{pa_{n_k}} \neq \overline{pa}$. To see this, note that, since $\{\overline{pa_n}\}_{n=1}^{\infty}$ does not converge, we have that $\liminf \overline{pa_n} \neq \limsup \overline{pa_n}$. Let $x \in \limsup \overline{pa_n} \setminus \liminf \overline{pa_n}$ and let V be an open subset of X containing x . Since $x \in \limsup \overline{pa_n}$, we obtain that V intersects infinitely many arcs $\overline{pa_n}$. Hence, since $\mathcal{C}_1(X)$ is compact [21, Theorem 1.8.5], we have that the sequence of arcs $\{\overline{pa_n}\}_{n=1}^{\infty}$ has a convergent subsequence $\{\overline{pa_{n_k}}\}_{k=1}^{\infty}$. Thus, $\{a_{n_k}\}_{k=1}^{\infty}$ is a subsequence of $\{a_n\}_{n=1}^{\infty}$ such that the sequence of arcs $\{\overline{pa_{n_k}}\}_{k=1}^{\infty}$ converges. Since $x \in \lim \overline{pa_{n_k}}$, $\overline{pa} \subset \liminf \overline{pa_n} \subset \lim \overline{pa_{n_k}}$ and $x \in X \setminus \liminf \overline{pa_n}$, we obtain that $\lim \overline{pa_{n_k}} \neq \overline{pa}$. \square

Theorem 4.6. *Let X be a dendroid that is smooth at the point p . If K is a closed subset of X , then $L = \bigcup_{z \in K} \overline{pz}$ is a subcontinuum of X .*

Proof. Note that L is a connected subset of X since L is the union of arcs all containing the point p . Let $x \in Cl(L)$. Then there exists a sequence $\{x_n\}_{n=1}^{\infty}$ of elements of L that converges to x . For each $n \in \mathbb{N}$, there exists an element z_n in K such that $x_n \in \overline{pz_n}$. Since K is compact, without loss of generality, we assume that there exists a point z in K such that the sequence $\{z_n\}_{n=1}^{\infty}$ converges to z . Since X is smooth at p , we have that the sequence of arcs $\{\overline{pz_n}\}_{n=1}^{\infty}$ converges to the arc \overline{pz} . Observe that we have that $x \in \overline{pz}$. Hence, $x \in L$ and L is a subcontinuum of X . \square

The following result is originally proved in [23, Theorem 1.16]. The proof given here is a modification of [20, p. 41].

Theorem 4.7. *If X is a smooth dendroid, then X is contractible.*

Proof. Suppose X is a dendroid that is smooth at the point p . Let $\mu: \mathcal{C}_1(X) \rightarrow [0, 1]$ be a Whitney map. For each element x of X and every number $t \in [0, 1]$, let x_t be the unique point in the arc \overline{px} such that $\mu(\overline{px_t}) = (1-t)\mu(\overline{px})$. Let $F: X \times [0, 1] \rightarrow X$ be given by $F((x, t)) = x_t$. Note that, for every $x \in X$, $F((x, 0)) = x$ and $F((x, 1)) = p$.

We need to show that F is continuous. Let $\{(x_n, t_n)\}_{n=1}^{\infty}$ be a sequence of points of $X \times [0, 1]$ converging to a point (x, t) in $X \times [0, 1]$. For each $n \in \mathbb{N}$, we have that $F((x_n, t_n)) = x_{nt_n}$. Since X is a continuum, we may assume that there exists an element w of X such that the sequence $\{x_{nt_n}\}_{n=1}^{\infty}$ converges to w . We prove that $F((x, t)) = w$.

From the definition of F , we obtain that, for every $n \in \mathbb{N}$, $\mu(px_{n_{t_n}}) = (1 - t_n)\mu(px_n)$. Since X is smooth at p , $\{x_n\}_{n=1}^\infty$ converges to x and $\{x_{n_{t_n}}\}_{n=1}^\infty$ converges to w , we have that $\lim \overline{px_n} = \overline{px}$, $\lim \overline{px_{n_{t_n}}} = \overline{pw}$ and $\overline{pw} \subset \overline{px}$. Using the continuity of μ and the fact that $\lim t_n = t$, these equalities imply that $\mu(\overline{pw}) = (1 - t)\mu(\overline{px})$. Hence, $F((x, t)) = w$. Therefore, F is continuous. \square

We end this section with a couple of characterizations of smooth dendroids.

Theorem 4.8. [4, Theorem 12] *A dendroid X is smooth if and only if there exists a point p in X such that for each pair of convergent sequences $\{a_n\}_{n=1}^\infty$ and $\{b_n\}_{n=1}^\infty$, the conditions:*

$$\lim a_n = a, \quad \lim b_n = b,$$

and

$$a_n \in \overline{pb_n}$$

imply that:

$$\lim \overline{a_n b_n} = ab.$$

Theorem 4.9. [4, Theorem 6] *A dendroid X is smooth if and only if for each pair of points x_1 and x_2 of X , we have that $\overline{x_1 x_2} \cap \mathcal{T}(\{x_1\}) = \{x_1\}$ or $\overline{x_1 x_2} \cap \mathcal{T}(\{x_2\}) = \{x_2\}$.*

5. Pointwise Smooth Dendroids

The class of pointwise smooth dendroids was introduced by Stanisław T. Czuba [7] as a generalization of the class of smooth dendroids. He continued the study of this class of dendroids in [9], [10], and [11].

A dendroid X is said to be *pointwise smooth* if, for each point x of X , there exists a element $p(x)$ of X such that for every sequence $\{x_n\}_{n=1}^\infty$ of points of X that converges to x , the sequence of arcs $\{\overline{p(x)x_n}\}_{n=1}^\infty$ converges to the arc $\overline{p(x)x}$. The point $p(x)$ is called an *initial point* for x in X . Note that in the case of a smooth dendroid the point of smoothness is the initial point of every element of the dendroid.

Theorem 5.1. *If X is a pointwise smooth dendroid and Z is a subcontinuum of X , then Z is a pointwise smooth dendroid.*

Proof. Let X be a pointwise smooth dendroid and let Z be a subcontinuum of X . By Theorem 3.1, Z is a dendroid. Let z be a point of Z . Since X is pointwise smooth, there exists a point $p(z)$ in X that is an initial point for z in X . We assume that $p(z) \in X \setminus Z$. Let $p'(z)$ be the first point of the arc $\overline{p(z)z}$ that is in Z . We show that $p'(z)$ is an initial point for z in Z . To this end, let $\{z_n\}_{n=1}^\infty$ be a sequence of elements of Z converging to z . Note that $\overline{zp(z)} = \overline{zp'(z)} \cup \overline{p'(z)p(z)}$ and, for each $n \in \mathbb{N}$, $\overline{z_n p(z)} = \overline{z_n p'(z)} \cup \overline{p'(z)p(z)}$. A similar argument to the one given in Theorem 4.2 shows that the sequence of arcs $\{\overline{z_n p'(z)}\}_{n=1}^\infty$ converges to the arc $\overline{zp'(z)}$. Therefore, Z is pointwise smooth. \square

Remark 5.2. Note that the following dendroid:




Figure 4.

is pointwise smooth but not smooth.

The next theorem shows that for the class of fans, pointwise smoothness and smoothness are equivalent [7, Theorem 1].

Theorem 5.3. *A fan X is pointwise smooth if and only if X is smooth.*

Proof. Suppose X is a pointwise smooth fan, let τ be the top of X , and let x be a point of X . We prove that τ is the initial point for x . Let $\{x_n\}_{n=1}^\infty$ be a sequence of elements of X converging to x . If there exists $N \in \mathbb{N}$ such that for $n \geq N$, we have that both x_n and x belong to the same component of $X \setminus \{\tau\}$ (this is a semi-open arc), we obtain that $\overline{\lim \tau x_n} = \overline{\tau x}$. Without loss of generality, we assume that for each $n \in \mathbb{N}$, we have that $\overline{x_n p(x)} = \overline{x_n \tau} \cup \overline{\tau p(x)}$ (if there exists an infinite set \mathbb{K} of \mathbb{N} and there exists a component Z of $X \setminus \{\tau\}$ such that if $n \in \mathbb{K}$, then $x_n \in Z$, we obtain that $\lim \overline{x_n \tau} = \overline{\tau x}$ ($n \in \mathbb{K}$) and we only need to take care of the indexes that belong to $\mathbb{N} \setminus \mathbb{K}$, this set is infinite and if n_1 and n_2 are two distinct elements of $\mathbb{N} \setminus \mathbb{K}$, then x_{n_1} and x_{n_2} belong to two distinct components of $X \setminus \{\tau\}$). Thus, for each $n \in \mathbb{N}$, we have that

$$\overline{x_n p(x)} = \overline{x_n \tau} \cup \overline{\tau p(x)}.$$

Hence,

$$\liminf \overline{x_n p(x)} = \liminf \overline{x_n \tau} \cup \overline{\tau p(x)}$$

[17, p. 336] and that

$$\limsup \overline{x_n p(x)} = \limsup \overline{x_n \tau} \cup \overline{\tau p(x)}$$

[17, p. 337]. Since X is pointwise smooth fan and $p(x)$ is the initial point for x in X , we obtain that

$$\liminf \overline{x_n p(x)} = \limsup \overline{x_n p(x)},$$

from this, we have

$$\overline{\tau p(x)} \cup \liminf \overline{x_n \tau} = \overline{\tau p(x)} \cup \limsup \overline{x_n \tau}.$$

Observe that $\overline{\tau x} \subset \liminf \overline{x_n \tau} \subset \limsup \overline{x_n \tau}$. We prove that $\limsup \overline{x_n \tau} \subset \overline{\tau x}$. Suppose that $(\limsup \overline{x_n \tau}) \setminus \overline{\tau x} \neq \emptyset$ and that $(\limsup \overline{x_n \tau}) \setminus \overline{\tau x} = \overline{z \tau} \setminus \{\tau\}$ [22, Theorem 1.1.30]. Let $w \in \overline{z \tau} \setminus \{z, \tau\}$. Let $\{x_{n_\ell}\}_{\ell=1}^\infty$ be a subsequence of $\{x_n\}_{n=1}^\infty$ such that $\overline{z \tau} \subset \lim \overline{x_{n_\ell} \tau}$. For every $\ell \in \mathbb{N}$, let w_ℓ be the first point of the arc $\overline{x_{n_\ell} \tau}$, from x_{n_ℓ} to τ , such that

$d(w, w_\ell) \leq \frac{1}{\ell}$. The sequence $\{w_\ell\}_{\ell=1}^\infty$ converges to w . Note that $\overline{z\tau} \subset \liminf \overline{w_\ell\tau}$. Hence, there does not exist an initial point $p(w)$ for w in X , a contradiction. Thus, $\limsup \overline{x_n\tau} \subset \overline{x\tau}$, and $\lim \overline{x_n\tau} = \overline{x\tau}$. Therefore, X is a smooth fan. The reverse implication is clear. \square

The following two theorems give characterizations of pointwise smooth dendroids.

Theorem 5.4. [22, Theorem 7.3.15] *Let X be a dendroid. Then X is pointwise smooth if and only if for each pair of different points x_1 and x_2 of X , we have that either $\overline{x_1x_2} \cap \mathcal{T}(\{x_1\}) = \{x_1\}$ or $\overline{x_1x_2} \cap \mathcal{T}(\{x_2\}) = \{x_2\}$ or $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) = \emptyset$.*

Theorem 5.5. [22, Corollary 7.3.16] *A dendroid X is pointwise smooth if and only if for each point x of X , there exists a point $p(x)$ in X such that for each sequence $\{x_n\}_{n=1}^\infty$ of points of X converging to x , the sequence of arcs $\{\overline{p(x)x_n}\}_{n=1}^\infty$ converges to the arc $\overline{p(x)x}$ and $x \in \mathcal{T}(\{p(x)\})$.*

Lemma 5.6. [22, Lemma 7.3.10] *If X is a dendroid and A and B are closed subsets of X such that $A \cap \mathcal{T}(B) = \emptyset$, $\mathcal{T}(A) \cap B = \emptyset$ and $\mathcal{T}(A) \cap \mathcal{T}(B) \neq \emptyset$, then there exist two points $p \in A$ and $q \in B$ such that $p \in X \setminus \mathcal{T}(\{q\})$, $q \in X \setminus \mathcal{T}(\{p\})$ and $\mathcal{T}(\{p\}) \cap \mathcal{T}(\{q\}) \neq \emptyset$.*

Remark 5.7. Lemma 5.6 is important since it gives a simple way to obtain the noncontractibility of a dendroid [22, Theorem 7.2.2].

The next theorem gives a sufficient condition for a continuum to contain an R^1 -continuum (R^3 -continuum (Lemma 2.7)).

Theorem 5.8. [22, Theorem 7.3.11] *If X is a dendroid and A and B are subcontinua of X such that $A \cap \mathcal{T}(B) = \emptyset$, $\mathcal{T}(A) \cap B = \emptyset$, and $\mathcal{T}(A) \cap \mathcal{T}(B) \neq \emptyset$, then $\mathcal{T}(A) \cap \mathcal{T}(B)$ is an R^1 -continuum. Therefore, it contains an R^3 -continuum.*

Corollary 5.9. [22, Corollary 7.3.12] *If X is a dendroid and A and B are closed subsets of X such that $A \cap \mathcal{T}(B) = \emptyset$, $\mathcal{T}(A) \cap B = \emptyset$ and $\mathcal{T}(A) \cap \mathcal{T}(B) \neq \emptyset$, then X contains an R^1 -continuum. Therefore, it contains an R^3 -continuum.*

If X is a dendroid and K and L are two disjoint and nonempty closed subsets of X , then

$$Irr(K, L) = \bigcap \{W \mid W \text{ is a subcontinuum of } X \text{ such that } K \cup L \subset W\}.$$

The set $Irr(K, L)$ is called the *irreducible subcontinuum of X about $K \cup L$* .

Remark 5.10. If X is a dendroid and K and L are two disjoint subcontinua of X , then by the arcwise connectedness and the hereditary unicoherence of X , we have that there exist two points k_0 in K and l_0 in L such that the arc $\overline{k_0l_0}$ satisfies that $K \cap \overline{k_0l_0} = \{k_0\}$, $L \cap \overline{k_0l_0} = \{l_0\}$ and $Irr(K, L) = K \cup \overline{k_0l_0} \cup L$.

The following result is [9, Corollary (3.3)].

Theorem 5.11. *A dendroid is pointwise smooth if and only if for any two disjoint subcontinua K and L of X , we have that $\mathcal{T}(K) \cap Irr(K, L) = K$ or $\mathcal{T}(L) \cap Irr(K, L) = L$ or $\mathcal{T}(K) \cap \mathcal{T}(L) = \emptyset$.*

Proof. Assume that X is a pointwise smooth dendroid and that K and L are two disjoint subcontuna of X such that

$$(\mathcal{T}(K) \cap Irr(K, L)) \setminus K \neq \emptyset \text{ and } (\mathcal{T}(L) \cap Irr(K, L)) \setminus L \neq \emptyset.$$

Observe that (Remark 5.10),

$$\mathcal{T}(K) \cap \overline{k_0 l_0} = \mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0} \text{ and } \mathcal{T}(L) \cap \overline{k_0 l_0} = \mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0}.$$

We show these two equalities are true. Note that $\mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0} \subset \mathcal{T}(K) \cap \overline{k_0 l_0}$ [22, Proposition 2.1.7]. Let z be an element of $\mathcal{T}(K) \cap \overline{k_0 l_0}$. Hence, if W is a subcontinuum of X such that $z \in Int(W)$, then $W \cap K \neq \emptyset$. Since X is hereditarily unicoherent, we obtain that $k_0 \in W$. Thus, $z \in \mathcal{T}(\{k_0\})$, and $z \in \mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0}$. Therefore, $\mathcal{T}(K) \cap \overline{k_0 l_0} = \mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0}$. Similarly, we have that $\mathcal{T}(L) \cap \overline{k_0 l_0} = \mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0}$. Next, we claim that

$$\mathcal{T}(K) \cap L = \emptyset \text{ and } K \cap \mathcal{T}(L) = \emptyset.$$

Assume that $\mathcal{T}(K) \cap L \neq \emptyset$. Since $\mathcal{T}(K)$ is a subcontinuum of X [22, Theorem 2.1.27] and X is a dendroid, we have that $\overline{k_0 l_0} \subset \mathcal{T}(K)$. Hence, $\mathcal{T}(K) \cap \overline{k_0 l_0} = \overline{k_0 l_0}$. Since $\mathcal{T}(K) \cap \overline{k_0 l_0} = \mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0}$ and $\mathcal{T}(L) \cap \overline{k_0 l_0} = \mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0}$, we obtain that $l_0 \in \mathcal{T}(\{k_0\}) \cap \mathcal{T}(\{l_0\})$. Also, since $(\mathcal{T}(K) \cap Irr(K, L)) \setminus K \neq \emptyset$ and $(\mathcal{T}(L) \cap Irr(K, L)) \setminus L \neq \emptyset$, we have that $\mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0} \neq \{k_0\}$ and $\mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0} \neq \{l_0\}$. Similarly, if $K \cap \mathcal{T}(L) \neq \emptyset$, then $k_0 \in \mathcal{T}(\{k_0\}) \cap \mathcal{T}(\{l_0\})$, $\mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0} \neq \{k_0\}$ and $\mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0} \neq \{l_0\}$. Thus, in any case, we obtain a contradiction to Theorem 5.4. Therefore, $\mathcal{T}(K) \cap L = \emptyset$ and $K \cap \mathcal{T}(L) = \emptyset$.

Since $\mathcal{T}(K) \cap \overline{k_0 l_0} = \mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0}$, we have that:

$$\begin{aligned} \mathcal{T}(K) \cap Irr(K, L) &= \mathcal{T}(K) \cap (\overline{k_0 l_0} \cup K \cup L) = \\ &= (\mathcal{T}(K) \cap \overline{k_0 l_0}) \cup K = (\mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0}) \cup K. \end{aligned}$$

Similarly, we obtain:

$$\mathcal{T}(L) \cap Irr(K, L) = (\mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0}) \cup L.$$

Hence,

$$\mathcal{T}(K) \cap \mathcal{T}(L) \cap Irr(K, L) = \mathcal{T}(\{k_0\}) \cap \mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0}.$$

Recall that $(\mathcal{T}(K) \cap Irr(K, L)) \setminus K \neq \emptyset$ implies that $\mathcal{T}(\{k_0\}) \cap \overline{k_0 l_0} \neq \{k_0\}$ and that $(\mathcal{T}(L) \cap Irr(K, L)) \setminus L \neq \emptyset$ implies that $\mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0} \neq \{l_0\}$. Thus, by Theorem 5.4, we have that

$$\mathcal{T}(\{k_0\}) \cap \mathcal{T}(\{l_0\}) \cap \overline{k_0 l_0} = \mathcal{T}(K) \cap \mathcal{T}(L) \cap Irr(K, L) = \emptyset.$$

Since X is hereditarily unicoherent, we obtain that $\mathcal{T}(K) \cap \mathcal{T}(L) = \emptyset$.

The reverse implication is obtained by Theorem 5.4, taking $K = \{x_1\}$ and $L = \{x_2\}$. Note that, in this case, $Irr(\{x_1\}, \{x_2\}) = \overline{x_1 x_2}$. \square

Let X be a dendroid and let p and x two be points of X . Define the sets:

$$P(x) = \{z \in X \mid z \text{ is an initial point for } x \text{ in } X\}$$

and

$$Q(p) = \{z \in X \mid p \text{ is an initial point for } z \text{ in } X\}.$$

Some consequences of the definition of these sets are:

Theorem 5.12. *Let X be a dendroid. Then the following hold:*

- (1) X is pointwise smooth if and only if $\bigcup\{Q(p) \mid p \in X\} = X$.
- (2) X is smooth if and only if $\bigcap\{P(x) \mid x \in X\} \neq \emptyset$.
- (3) X is smooth if and only if there exists an element p of X such that $Q(p) = X$.

Theorem 5.13. *Let X be a dendroid and let x_0 be a point of X . Then the following are equivalent:*

- (1) $P(x_0) = X$;
- (2) $x_0 \in Q(x_0)$;
- (3) $x_0 \in P(x_0)$;
- (4) $x_0 \in X \setminus \mathcal{T}(\{x\})$, for all $x \in X \setminus \{x_0\}$;
- (5) X is connected im kleinen at x_0 .

Proof. Assume that $P(x_0) = X$. This means that each element x of X is an initial point for x_0 in X . In particular, x_0 is an initial point for itself in X . Thus, $x_0 \in Q(x_0)$. Next, if $x_0 \in Q(x_0)$, then x_0 is an initial point for itself in X . Hence, $x_0 \in P(x_0)$. Now, assume that $x_0 \in P(x_0)$. This implies that for each sequence $\{x_n\}_{n=1}^\infty$ of elements of X converging to x_0 , we have that the sequence of arcs $\{\overline{x_0x_n}\}_{n=1}^\infty$ converges to $\{x_0\}$. Thus, $\lim \text{diam}(\overline{x_0x_n}) = 0$. From this, we obtain that if p is a point of X and $\{x_n\}_{n=1}^\infty$ is a sequence of elements of X converging to x_0 , we have that the sequence of arcs $\{\overline{px_n}\}_{n=1}^\infty$ converges to the arc $\overline{px_0}$. Therefore, $P(x_0) = X$. Hence, (1), (2) and (3) are equivalent.

Suppose that $x_0 \in X \setminus \mathcal{T}(\{x\})$, for every point x in $X \setminus \{x_0\}$. We show that X is connected im kleinen at x_0 . To this end, let A be a closed subset of X such that $x_0 \in \mathcal{T}(A)$. Since X is hereditarily unicoherent, by [22, Theorem 2.2.12], X is \mathcal{T} -additive. Thus, by [22, Corollary 2.2.13], we have that $\mathcal{T}(A) = \bigcup\{\mathcal{T}(\{a\}) \mid a \in A\}$. Hence, there exists a point a in A such that $x_0 \in \mathcal{T}(\{a\})$. By our assumption, we obtain that $x_0 = a$ and $x_0 \in A$. By [22, Corollary 2.1.31], we conclude that X is connected im kleinen at x_0 . Now, assume that X is connected im kleinen at x_0 , let x be an element of $X \setminus \{x_0\}$ and let U be an open subset of $X \setminus \{x\}$ that contains x_0 . Since X is connected im kleinen at x_0 , there exists a subcontinuum W of X such that $x_0 \in \text{Int}(W) \subset W \subset U \subset X \setminus \{x\}$. This implies that $x_0 \in X \setminus \mathcal{T}(\{x\})$. Thus, (4) and (5) are equivalent.

Suppose that X is connected im kleinen at x_0 . Then x_0 is in the interior of subcontinua of X of arbitrary small diameters. By Lemma 3.1 each of these subcontinua is a dendroid. This implies that if $\{x_n\}_{n=1}^\infty$ is a sequence of points of X converging to x_0 ,

then $\lim \text{diam}(\overline{x_0x_n}) = 0$. Hence, if p is an element of X and $\{x_n\}_{n=1}^\infty$ is a sequence of points of X converging to x_0 , then the sequence of arcs $\{\overline{px_n}\}_{n=1}^\infty$ converges to the arc $\overline{px_0}$. Therefore, $P(x_0) = X$. Now, assume that X is not connected in the kleinian sense at x_0 . Thus, there exists an open subset U of X containing x_0 such that no subcontinuum of X contained in U has x_0 in its interior. Since X is a regular space, there exists an open subset V of X such that $x_0 \in V \subset Cl_X(V) \subset U$. Let W be the component of $Cl(V)$ that contains x_0 . By [22, Theorem 1.4.36], we have that W is a nondegenerate subcontinuum of X . By our assumption, $x_0 \in W \setminus \text{Int}(W)$. Hence, there exists a sequence $\{x_n\}_{n=1}^\infty$ of points of $X \setminus W$ that converges to x_0 . Consider the sequence of arcs $\{\overline{x_0x_n}\}_{n=1}^\infty$. Observe that, for each $n \in \mathbb{N}$, we obtain that $\overline{x_0x_n} \cap (X \setminus W) \neq \emptyset$. We have two possibilities, either the sequence $\{\overline{x_0x_n}\}_{n=1}^\infty$ converges and in this case, $(X \setminus W) \cap \lim \overline{x_0x_n} \neq \emptyset$ or such sequence does not converge. In either case, we obtain that $x_0 \in X \setminus P(x_0)$. Therefore, $P(x_0) \neq X$. Hence, (5) and (1) are equivalent. \square

By Theorem 4.7, we know that a smooth dendroid is contractible. It was asked if the same was true for pointwise smooth dendroids. Note the following result.

Theorem 5.14. *Let X be a dendroid. Then X is pointwise smooth if and only if $C_n(X)$ is contractible, for each $n \in \mathbb{N}$.*

Proof. By [5, Corollary, p. 411], we have that X is a pointwise smooth dendroid if and only if $C_1(X)$ is contractible. Hence, by [21, Theorem 6.1.17], we obtain the result for $n \geq 2$. \square

A dendroid X has *property (CS)* if there exists a clump \mathcal{G} of smooth subdendroids of X whose centre is C such that:

- (1) $\bigcup \mathcal{G} = X$;
- (2) there exists an element $p \in C$ such that p is the point of smoothness of each of the elements of \mathcal{G} ;
- (3) the set $Cl(\bigcup \mathcal{G} \setminus C) \cap C$ is totally disconnected.

Theorem 5.15. [11, Theorem 3] *If X is a pointwise smooth dendroid having property (CS), then X is a smooth dendroid.*

Regarding the contractibility of a pointwise smooth dendroid, we have:

Theorem 5.16. [11, Corollary 4] *Let X be a dendroid having property (CS). Then the following are equivalent:*

- (1) X is hereditarily contractible;
- (2) X is pointwise smooth;
- (3) X is smooth.

Let X be a continuum. If L is a subcontinuum of X that is contained in a subset A of X , then $C(A, K)$ denotes the component of A containing L .

Theorem 5.17. *Let X be a dendroid. Then the following are equivalent:*

- (1) X is pointwise smooth;
- (2) for each point x of X , there exists a point z of X such that for every subcontinuum K of X which contains an arc \overline{xz} and for each open subset U of X that contains K , we have that $x \in \text{Int}(C(U, K))$;
- (3) for each element x of X , there exists a point z of X such that for every open subset U of X containing \overline{xz} , we have $x \in \text{Int}(C(U, K))$.

Proof. To show that (1) implies (2), let x be a point of the pointwise smooth dendroid X and let K be a subcontinuum of X containing the arc $\overline{xp(x)}$. Let U be an open subset of X containing K . If $x \in X \setminus \text{Int}(C(U, K))$, then there exists a sequence $\{x_n\}_{n=1}^\infty$ of points of $X \setminus C(U, K)$ converging to x . Observe that $\overline{x_np(x)} \setminus U \neq \emptyset$. Thus, since X is pointwise smooth, $\lim \overline{x_np(x)} \cap (X \setminus U) \neq \emptyset$. Since $K \subset U$, we have that $K \cap (X \setminus U) = \emptyset$. By Theorem 3.1, $\lim \overline{x_np(x)} = \overline{xp(x)} \subset K$, a contradiction. Hence, $x \in \text{Int}(C(U, K))$.

It is clear that (2) implies (3). Suppose (3), we prove (1). Let z be a point of X and let $\{x_n\}_{n=1}^\infty$ be a sequence of elements of X converging to an element x of X . If $\limsup \overline{x_nz} \neq \overline{xz}$, then let $w \in \limsup \overline{x_nz} \setminus \overline{xz}$. Let U be an open subset of X such that $w \in \text{Cl}(U)$ and $\overline{xz} \subset U$. Note that for each $n \in \mathbb{N}$, $x_n \in X \setminus C(U, \{x\})$. Thus, $x \in X \setminus \text{Int}(C(U, \{x\}))$, a contradiction. Since $\liminf \overline{x_nz}$ is a continuum (Theorem 2.2) containing x and z , we have that $\liminf \overline{x_nz} = \limsup \overline{x_nz} = \overline{xz}$. Hence, we may take z to be $p(x)$, the initial point of x in X . □

Theorem 5.18. [10, Proposition 4] *A dendroid X is pointwise smooth if and only if for each point x of X and every sequence $\{x_n\}_{n=1}^\infty$ of elements of X converging to x , there exists a point q of X such that $\limsup \overline{x_nx} = \overline{qx}$ and for each initial point $p(x)$ for x in X , we have that $\overline{qx} \subset \overline{xp(x)}$.*

Theorem 5.19. *A dendroid X is pointwise smooth if and only if for each point x of X , $\mathcal{K}(\{x\})$ is an arc \overline{xz} , where z may be taken as an initial point for x in X ($z = p(x)$), in the sense of Theorem 5.5).*

Proof. Let X be a pointwise smooth dendroid, let x be an element of X and let $p(x)$ be an initial point for x in X (Theorem 5.5). Since $x \in \mathcal{T}(\{p(x)\})$, by [22, Lemma 7.7.3], $p(x) \in \mathcal{K}(\{x\})$. Thus, $\{x, p(x)\} \subset \mathcal{K}(\{x\})$ and, by Theorem 2.4, we have that $\overline{xp(x)} \subset \mathcal{K}(\{x\})$. If $\mathcal{K}(\{x\}) \setminus \overline{xp(x)} \neq \emptyset$, let $w \in \mathcal{K}(\{x\}) \setminus \overline{xp(x)}$. We consider three cases.

Case (1). $x \in \overline{wp(x)}$.

In this case, $\{w\} \neq \overline{xw}$. Since $w \in \mathcal{K}(\{x\})$, by [22, Lemma 7.7.3], $x \in \mathcal{T}(\{w\})$. Hence, $\overline{xw} \subset \overline{wp(x)} \subset \mathcal{T}(\{w\})$ [22, Theorem 2.1.27]. Also, $\{p(x)\} \neq \overline{xp(x)}$. Since $x \in \mathcal{T}(\{p(x)\})$, we obtain that $\overline{xp(x)} \subset \overline{wp(x)} \cap \mathcal{T}(\{p(x)\})$. Form all of this, we have that $\{w\} \neq \overline{wp(x)} \cap \mathcal{T}(\{w\})$, $\{p(x)\} \neq \overline{wp(x)} \cap \mathcal{T}(\{p(x)\})$ and $x \in \mathcal{T}(\{w\}) \cap \mathcal{T}(\{p(x)\})$. Thus, by Theorem 5.4, we have that X is not a pointwise smooth dendroid.

Case(2). $p(x) \in \overline{xw}$.

It follows from Theorem 5.18 that $p(x)$ is not an initial point for x in X .

Case (3). $\overline{wx} \cup \overline{xp(x)}$ is a simple triod.

In this case, there exists an element t of X such that $\overline{wx} \cup \overline{xp(x)} = \overline{wt} \cup \overline{tp(x)} \cup \overline{tx}$, where t is the vertex of the simple triod. Hence, $\{w\} \neq \overline{wt} \subset \overline{wx}$. Since $w \in \mathcal{K}(\{x\})$, by [22, Lemma 7.7.3], $x \in \mathcal{T}(\{w\})$. Thus, $\overline{wx} \subset \mathcal{T}(\{w\})$ [22, Theorem 2.1.27]. Also, $\{x\} \neq \overline{tx} \subset \overline{wx}$. Since $w \in \mathcal{K}(\{x\})$, we obtain that $\overline{wx} \subset \mathcal{T}(\{x\})$ [22, Theorem 2.1.27]. From all this, we have that $\{w\} \neq \overline{wx} \cap \mathcal{T}(\{w\})$, $\{x\} \neq \overline{wx} \cap \mathcal{T}(\{x\})$ and $t \in \mathcal{T}(\{w\}) \cap \mathcal{T}(\{x\})$. Thus, by Theorem 5.4, we have that X is not a pointwise smooth dendroid.

From these cases, we obtain that $\mathcal{K}(\{x\}) = \overline{xp(x)}$.

Now, suppose X is not a pointwise smooth dendroid. Then, by Theorem 5.4, there exist two elements a and b of X such that $\{a\} \neq \overline{ab} \cap \mathcal{T}(\{a\})$, $\{b\} \neq \overline{ab} \cap \mathcal{T}(\{b\})$, and $\mathcal{T}(\{a\}) \cap \mathcal{T}(\{b\}) \neq \emptyset$. Since X is hereditarily unicoherent, there exists a point $x \in \mathcal{T}(\{a\}) \cap \mathcal{T}(\{b\}) \cap \overline{ab} \setminus \{a, b\}$. By [22, Lemma 7.7.3], we have that $\{a, b\} \subset \mathcal{K}(\{x\})$. This implies that $\mathcal{K}(\{x\})$ is not an arc with x as one of its endpoints. \square

As a consequence of Theorems 5.5 and 5.19, we have:

Corollary 5.20. *If X is a pointwise smooth dendroid, then for each element x of X , there exists a unique point $p(x)$ (namely, the endpoint of the arc $\mathcal{K}(\{x\})$ different from x) such that for each sequence $\{x_n\}_{n=1}^\infty$ of points of X converging to x , $\lim p(x)x_n = p(x)x$, and $x \in \mathcal{T}(\{p(x)\})$.*

Corollary 5.21. *If X is a pointwise smooth dendroid, then no point from the open arc $\mathcal{K}(\{x\}) \setminus \{x, p(x)\}$ is an initial point for x in X , where $\mathcal{K}(\{x\}) = \overline{p(x)x}$.*

Theorem 5.22. *If X is a pointwise smooth dendroid, then for each pair of points x_1 and x_2 of X , the following are satisfied:*

- (1) $\mathcal{K}(\{x_1\}) = \mathcal{K}(\{x_2\})$ if and only if $x_1 = x_2$;
- (2) $\mathcal{T}(\{x_1\}) = \mathcal{T}(\{x_2\})$ if and only if $x_1 = x_2$;
- (3) $\mathcal{T}(\{x_1\}) \cap \mathcal{K}(\{x_1\}) = \{x_1\}$;
- (4) $\mathcal{K}(\mathcal{T}(\{x_1\})) = \mathcal{T}(\{x_1\}) \cup \mathcal{K}(\{x_1\})$.

Proof. We show (1). If $\mathcal{K}(\{x_1\}) = \mathcal{K}(\{x_2\})$, by Theorem 5.19, $\mathcal{K}(\{x_1\}) = \overline{p_1(x_1)x_1}$ and $\mathcal{K}(\{x_2\}) = \overline{p_1(x_2)x_2}$. Hence, either, $x_1 = x_2$ or $p(x_1) = x_2$ or $p(x_2) = x_1$. If $p(x_1) = x_2$, then $\mathcal{K}(\{x_1\}) = \mathcal{K}(\{x_2\}) = \overline{x_1x_2}$. This implies, by [22, Lemma 7.7.3], that $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_2} = \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} = \overline{x_1x_2}$ and $\overline{x_1x_2} \subset \mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\})$. Hence, by Theorem 5.4, X is not a pointwise smooth dendroid, a contradiction. Similarly, we obtain a contradiction if $p(x_2) = x_1$. Therefore, $x_1 = x_2$. The reverse implication is clear.

To prove (2), note that if $\mathcal{T}(\{x_1\}) = \mathcal{T}(\{x_2\})$ and $x_1 \neq x_2$, then $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_2} = \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} = \overline{x_1x_2}$ and $\overline{x_1x_2} \subset \mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\})$. Thus, by Theorem 5.4, X is not a pointwise smooth dendroid, a contradiction. Therefore, $x_1 = x_2$. The reverse implication is clear.

To see (3), let $x_3 \in \mathcal{T}(\{x_1\}) \cap \mathcal{K}(\{x_1\}) \setminus \{x_1\}$. Then, by [22, Lemma 7.7.3], we have that $x_1 \in \mathcal{T}(\{x_3\})$ and $x_3 \in \mathcal{T}(\{x_1\})$. Hence, $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_3} = \mathcal{T}(\{x_3\}) \cap \overline{x_1x_3} = \overline{x_1x_3}$

and $\overline{x_1x_3} \subset \mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_3\})$. Thus, by Theorem 5.4, X is not a pointwise smooth dendroid, a contradiction. Therefore, $\mathcal{T}(\{x_1\}) \cap \mathcal{K}(\{x_1\}) = \{x_1\}$.

To show (4), observe that since $x_1 \in \mathcal{T}(\{x_1\})$, we have that $\mathcal{K}(\{x_1\}) \subset \mathcal{K}(\mathcal{T}(\{x_1\}))$ and, by definition, $\mathcal{T}(\{x_1\}) \subset \mathcal{K}(\mathcal{T}(\{x_1\}))$. Hence, $\mathcal{T}(\{x_1\}) \cup \mathcal{K}(\{x_1\}) \subset \mathcal{K}(\mathcal{T}(\{x_1\}))$. Let $x_3 \in \mathcal{K}(\mathcal{T}(\{x_1\}))$. By Theorem 2.5, we obtain that $\mathcal{T}(\{x_3\}) \cap \mathcal{T}(\{x_1\}) \neq \emptyset$. By Theorem 5.4, we have that either $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_3} = \{x_1\}$ or $\mathcal{T}(\{x_3\}) \cap \overline{x_1x_3} = \{x_3\}$. Suppose $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_3} = \{x_1\}$, the proof of the other case is analogous. We claim that, in this case, $\overline{x_1x_3} \subset \mathcal{T}(\{x_3\})$. If this is not true, then $(\mathcal{T}(\{x_1\}) \cup \mathcal{T}(\{x_3\})) \cup \overline{x_1x_3}$ would contain a nonunicoherent subcontinuum of X (Let x_4 be the first point of the arc $\overline{x_1x_3}$ in $\mathcal{T}(\{x_3\})$, and let $x_5 \in \mathcal{T}(\{x_3\}) \cap \mathcal{T}(\{x_1\})$. Then $\overline{x_1x_4} \cup \overline{x_4x_5} \cup \overline{x_5x_1}$ is a simple closed curve contained in X), a contradiction. Hence, $x_1 \in \mathcal{T}(\{x_3\})$ and, by [22, Lemma 7.7.3], $x_3 \in \mathcal{K}(\{x_1\})$. Therefore, $\mathcal{K}(\mathcal{T}(\{x_1\})) = \mathcal{T}(\{x_1\}) \cup \mathcal{K}(\{x_1\})$. \square

Theorem 5.23. [9, Theorem 3.7] *If X is a pointwise smooth dendroid, then X does not contain an R^3 -continuum.*

The following result is [10, Theorem 16].

Theorem 5.24. *If X is a dendroid, then the following are equivalent:*

- (1) X is pointwise smooth dendroid.
- (2) X is semi-aposyndetic and X does not contain an R^3 -continuum.
- (3) X is semi-aposyndetic and X does not contain two points x_1 and x_2 such that $x_1 \in X \setminus \mathcal{T}(\{x_2\})$, $x_2 \in X \setminus \mathcal{T}(\{x_1\})$ and $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \neq \emptyset$.

Proof. We prove that (1) implies (2). Suppose X is a pointwise smooth dendroid. Let x_1 and x_2 be two points of X . By Theorem 5.4, either $x_1 \in X \setminus \mathcal{T}(\{x_2\})$ or $x_2 \in X \setminus \mathcal{T}(\{x_1\})$. Hence, by [22, Theorem 2.1.32], X is semi-aposyndetic. By Theorem 5.23, X does not contain an R^3 -continuum. The fact that (2) implies (3) follows from Corollary 5.9 and [22, Lemma 7.3.10]. To show (3) implies (1), let x_1 and x_2 be two points of X such that $\mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} \neq \{x_2\}$ and $\mathcal{T}(\{x_1\}) \cap \overline{x_1x_2} \neq \{x_1\}$. Suppose that $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \neq \emptyset$. Then, since X is hereditarily unicoherent, $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} \neq \emptyset$. If $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} = \{w\}$, for some element w of X , we have that $w \in X \setminus \{x_1, x_2\}$. Hence, $x_1 \in X \setminus \mathcal{T}(\{x_2\})$ and $x_2 \in X \setminus \mathcal{T}(\{x_1\})$ (if $x_1 \in \mathcal{T}(\{x_2\})$, then $\overline{x_1x_2} \subset \mathcal{T}(\{x_2\})$ and $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} = \mathcal{T}(\{x_1\}) \cap \overline{x_1x_2} \neq \{w\}$, a contradiction; similarly if $x_2 \in X \setminus \mathcal{T}(\{x_1\})$). Thus, we obtain a contradiction to our assumption. If $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) \cap \overline{x_1x_2} = \overline{wz}$, where $w \in \overline{x_1z}$, then, since $z \in \mathcal{T}(\{x_1\})$, we have that $z \in \mathcal{T}(\{w\})$ (if $z \in X \setminus \mathcal{T}(\{w\})$, then there exists a subcontinuum Z of X such that $z \in \text{Int}(Z) \subset Z \subset X \setminus \{w\}$, since $z \in \mathcal{T}(\{x_1\})$, we obtain that $x_1 \in Z$; thus, $w \in Z$, a contradiction). Also, since $w \in \mathcal{T}(\{x_2\})$, we have that $w \in \mathcal{T}(\{z\})$. Hence, by [22, Lemma 2.1.32], X is not semi-aposyndetic. Thus, $\mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\}) = \emptyset$. Therefore, by Theorem 5.4, X is pointwise smooth. \square

6. Strict Point \mathcal{T} -asymmetry on Dendroids

Strict point \mathcal{T} -asymmetric continua were defined by David P. Bellamy [19, p. 392]. We present new results about strict point \mathcal{T} -asymmetry.

A continuum X is *strictly point \mathcal{T} -asymmetric* if for any two distinct points p and q of X with $p \in \mathcal{T}(\{q\})$, we have that $q \in X \setminus \mathcal{T}(\{p\})$.

David P. Bellamy asked:

Question 6.1. [22, Question 8.1.9] If X is a strictly point \mathcal{T} -asymmetric dendroid, then is X smooth?

Leobardo Fernández gave a negative answer to this question [13]. The dendroid presented in Remark 5.2 (Figure 4) is an example of a nonsmooth dendroid that is strictly point \mathcal{T} -asymmetric. He proved that the reverse implication is true [13, Corollary 3.3]. He also showed that for fans strict point \mathcal{T} -asymmetry is equivalent to smoothness [13, Theorem 3.5]. We extend these results to pointwise smooth dendroids (Corollary 6.3).

We present a characterize of strictly point \mathcal{T} -asymmetric continua as semi-aposyndetic continua.

Theorem 6.2. *Let X be a continuum. Then X is point \mathcal{T} -asymmetric if and only if X is semi-aposyndetic.*

Proof. Suppose X is strictly point \mathcal{T} -asymmetric and let p and q be two points of X . We have that either $p \in X \setminus \mathcal{T}(\{q\})$ or $p \in \mathcal{T}(\{q\})$. In the second case, since X is strictly point \mathcal{T} -asymmetric, we obtain that $q \in X \setminus \mathcal{T}(\{p\})$. Therefore, X is semi-aposyndetic [22, Theorem 2.1.32].

Assume that X is semi-aposyndetic and let p and q be two points of X . Suppose $p \in \mathcal{T}(\{q\})$. Since X is semi-aposyndetic and $p \in \mathcal{T}(\{q\})$, we have that $q \in X \setminus \mathcal{T}(\{p\})$ [22, Theorem 2.1.32]. Therefore, X is strictly point \mathcal{T} -asymmetric. \square

Next, we show that pointwise smooth dendroids are strictly point \mathcal{T} -asymmetric.

Corollary 6.3. *If X is a pointwise smooth dendroid, then X is strictly point \mathcal{T} -asymmetric.*

Proof. Suppose X is not strictly point \mathcal{T} -asymmetric. Then there exist two points x_1 and x_2 of X such that $x_1 \in \mathcal{T}(\{x_2\})$ and $x_2 \in \mathcal{T}(\{x_1\})$. Since $\{x_1, x_2\} \subset \mathcal{T}(\{x_2\})$, we have that $\overline{x_1 x_2} \subset \mathcal{T}(\{x_2\})$. Similarly, we have that $\overline{x_1 x_2} \subset \mathcal{T}(\{x_1\})$. Hence, $\overline{x_1 x_2} \cap \mathcal{T}(\{x_2\}) = \overline{x_1 x_2}$ and $\overline{x_1 x_2} \cap \mathcal{T}(\{x_1\}) = \overline{x_1 x_2}$. Thus, $\overline{x_1 x_2} \subset \mathcal{T}(\{x_1\}) \cap \mathcal{T}(\{x_2\})$. Therefore, by Theorem 5.4, X is not pointwise smooth. \square

As a consequence of Theorem 6.2 and Corollary 6.3, we have:

Corollary 6.4. *If X is a pointwise smooth dendroid, then X is semi-aposyndetic.*

Remark 6.5. Note that Corollary 6.4 is also a consequence of Theorem 5.24.

Corollary 6.6. *Let X be a fan. Then the following are equivalent:*

- (1) X is strictly point \mathcal{T} -asymmetric.
- (2) X is a smooth fan.
- (3) X is a pointwise smooth fan.
- (4) X is semi-aposyndetic.

Proof. By [13, Theorem 3.5], (1) and (2) are equivalent. By Theorem 5.3, (2) and (3) are equivalent. By Theorem 6.2, (1) and (4) are equivalent. \square

We end the paper with another characterization of pointwise smooth dendroids.

Theorem 6.7. *A dendroid X is pointwise smooth if and only if X is strictly point \mathcal{T} -asymmetric and X does not contain an R^3 -continuum.*

Proof. Suppose X is a pointwise smooth dendroid. Then, by Theorem 5.24, X is semi-aposyndetic and X does not contain an R^3 -continuum. Since X is semi-aposyndetic, by Theorem 6.2, X is strictly point \mathcal{T} -asymmetric.

Assume that X is a strictly point \mathcal{T} -asymmetric dendroid such that X does not contain an R^3 -continuum. Since X is strictly point \mathcal{T} -asymmetric, by Theorem 6.2, X is semi-aposyndetic. Hence, by Theorem 5.24, X is pointwise smooth. \square

References

- [1] Borsuk K., “A theorem on fixed points”, *Bull. Acad. Polon. Sci.*, 2 (1954), 12-20.
- [2] Charatonik J.J., “On fans”, *Dissertationes Math.*, (Rozprawy Mat.), 54 (1967), 1-37.
- [3] Charatonik J.J. and Charatonik W.J., “Dendrites”, *Aportaciones Matemáticas*, Sociedad Matemática Mexicana, Serie Comunicaciones, 22 (1998), 227-253.
- [4] Charatonik J.J. and Eberhart C., “On smooth dendroids”, *Fund. Math.*, 67 (1970), 297-322.
- [5] Charatonik W.J., “Pointwise smooth dendroids have contractible hyperspaces”, *Bull. Polish Acad. Sci. Math.*, 33 (1985), 409-412.
- [6] Charatonik W.J., “ R^i -continua and hyperspaces”, *Topology Appl.*, 23 (1986), No. 3, 207-216.
- [7] Czuba S.T., “A concept of pointwise smooth dendroids”, *Upeishi Math. Nauk.*, 34 (1979), 215-217 (*Russian Math. Surveys*, 34 (1979), 169-171).
- [8] Czuba S.T., “ R^i -continua and contractibility”, Proceedings of the International Conference on Geometric Topology PWN Warszawa, Poland, 75-79, August 24-September 2, 1980.
- [9] Czuba S.T., “On pointwise smooth dendroids”, *Fund. Math.*, 114 (1981), 197-207.
- [10] Czuba S.T., “Some other characterizations of pointwise smooth dendroids”, *Comment. Math. Prace Mat.*, 24 (1984), 195-200.

- [11] Czuba S.T., “On dendroids for which smoothness, pointwise smoothness and hereditarily contractibility are equivalent”, *Comment. Math. Prace Mat.*, 25 (1985), 27-30.
- [12] Eberhart C., “A note on smooth fans”, *Colloquium Math.*, 20 (1969), 89-90.
- [13] Fernández L., “On strictly point \mathcal{T} -asymmetric continua”, *Topology Proc.*, 35 (2010), 91-96.
- [14] Fugate F.B., Gorth G.R and Lum L., “Arc-smooth continua”, *Trans. Amer. Math. Soc.*, 265 (1981), 545-561.
- [15] Grace E.E. and Vought E.J., “Weakly monotone images of smooth dendroids are smooth”, *Houston J. Math.*, 14 (1988), 191-200.
- [16] Hocking J.G. and Young G.S., *Topology*, Dover Publications Inc, New York, 1988.
- [17] Kuratowski K., *Topology*, vol. 1, Academic Press, New York, 1966.
- [18] Kuratowski K., *Topology*, vol. 2, Academic Press, New York, 1968.
- [19] Lewis W., “Continuum theory problems”, *Topology Proc.*, 8 (1983), 361-394.
- [20] Macías S., “La Estructura de los Dendroides Suaves”, *Aportaciones Matemáticas de la Sociedad Matemática Mexicana, Serie Comunicaciones*, 10 (1993).
- [21] Macías S., *Topics on Continua*, Second Edition, Springer-Cham, 2018.
- [22] Macías S., *Set Function \mathcal{T} : An Account on F.B. Jones’ Contributions to Topology, Series Developments in Mathematics*, Vol. 67, Springer, 2021.
- [23] Mohler L., “A characterization of smoothness of dendroids”, *Fund. Math.*, 67 (1970), 369-376.
- [24] Nadler S.B., *Hyperspaces of Sets: A Text with Research Questions*, Monographs and Textbooks in Pure and Applied Mathematics, Vol. 49, Marcel Dekker, New York, Basel, 1978. Reprinted in: *Aportaciones Matemáticas de la Sociedad Matemática Mexicana, Serie Textos*, 33, 2006.
- [25] Nadler S.B. *Continuum Theory: An Introduction*, Monographs and Textbooks in Pure and Applied Math., Vol. 158, Marcel Dekker, New York, Basel, Hong Kong, 1992.