

# The first record of endangered *Lycaena helle* ([Denis & Schiffermüller], 1775) for Turkey (Lepidoptera: Lycaenidae)

S. S. Çalıřkan & A. Hasbenli

## Abstract

*Lycaena helle* ([Denis & Schiffermüller], 1775), is listed as “endangered” on the European Red List of butterflies. We report the first record of this species from Turkey. The EUNIS habitat information and photographs of the area where adult individuals was collected together with habitus dorsal and ventral photographs of the species were provided. The distribution status and habitat of the species in the Palaearctic region were evaluated.

KEY WORDS: Lepidoptera, Lycaenidae, *Lycaena helle*, new record, Eunis, Turkey.

## Primer registro de *Lycaena helle* ([Denis & Schiffermüller], 1775) en peligro de extinción para Turquía (Lepidoptera: Lycaenidae)

## Resumen

*Lycaena helle* ([Denis & Schiffermüller], 1775), está en la lista Roja Europea de mariposas en peligro de extinción. Citamos el primer registro de esta especie en Turquía. Se proporciona información y fotografías del hábitat EUNIS del área donde se colectó un individuo adulto, junto con una fotografía de la especie de su habitus dorsal y ventral. Fueron evaluadas la distribución, estatus y hábitat de la especie en la región Paleártica.

PALABRAS CLAVE: Lepidoptera, Lycaenidae, *Lycaena helle*, nuevo registro, Eunis, Turquía.

## Introduction

*Lycaena helle* ([Denis & Schiffermüller], 1775), is a species with the relict Boreo-montane Palaearctic distribution (POPOVIĆ *et al.*, 2014; BOZANO, 2004; HABEL *et al.*, 2014). It is found in most of Fennoscandia throughout Central and Northern Europe (POPOVIĆ *et al.*, 2014). It extends from Siberia to the Ussuri region in Russia Belarus, Estonia, Georgia, Kazakhstan, Lithuania, Ukraine to Mongolia and China in the Far East (TUZOV *et al.*, 2000; NEKRUTENKO & TSHIKOLOVETS, 2005; BOZANO & WEIDENHOFFER, 2001; KORB & BOLSHAKOV, 2016). The presence of this species in Serbia and Bulgaria in the Balkan Peninsula is an important zoogeographic record (KOLEV & SHTINKOV, 2015).

This species uses peat and sphagnum swamps, flowering moist meadows, forest edges and openings, creeks and slopes as its habitat (TSHIKOLOVETS, 2011). It is a postglacial relict in Central Europe and lives mostly in the highlands (HABEL *et al.*, 2011; MARTIN *et al.*, 2014). However, some populations are found in moist low meadows (SKORKA *et al.*, 2007). The population of the species in Europe has expanded to places where suitable habitats are found (POPOVIĆ *et al.*, 2014).

There was a significant decrease in the populations of *Lycaena helle* in many countries in the last decade (VAN SWAAY & WARREN, 1999; KUDRNA *et al.*, 2011). Populations in Western and

Central Europe decreased considerably (by 50 to 80%) in the last century. It is currently considered extinct in Hungary, the Czech Republic, Italy, Latvia, and Slovakia (VAN SWAAY *et al.*, 2010a).

*L. helle* is listed in the European Red Data Book and in the Annexes of the Habitats Directive (VAN HELSDİNGEN *et al.*, 1996; VAN SWAAY & WARREN 1999; VAN SWAAY *et al.*, 2010a).

The aim of this study is to provide information about and discuss the distribution and habitat of this butterfly species in Turkey, which was recorded for the first time in Turkey.

## Materials and Methods

A specimen of *Lycaena helle* species was collected in Posof district of Ardahan province in 17 July 2020 (Map 1). The specimen was collected during the TANAP (Trans-Anatolian Pipeline), project monitoring studies. The line transect method was employed in monitoring studies (Fig. 1). The specimen prepared in accordance with the museum methods is preserved in the Zoology Museum of Gazi University (ZMGU, Ankara, Turkey). Photographs of the dorsal and ventral wing of the specimen were taken with a Canon camera EOS 50D (Fig. 2).




**Map 1.**– The location of the specimen of *L. helle* ([D. & Schiff.]) collected in Turkey (red pin).

## Result and discussion

Within the scope of the Trans-Anatolian Pipeline (TANAP), which is being carried out in Turkey, fauna monitoring studies were carried out along the line. Critical butterfly species were identified, and the EUNIS types of habitats where they spread were determined according to DAVIES *et al.* (2004). The habitat where the species was caught is the “Calcareous alpine and subalpine grassland”, located between 2100-2400 m (Fig. 3). The EUNIS habitat code of this habitat is “E4.4”. This habitat is suitable for *L. helle*, and it is usual for the species to survive in this habitat. The larvae feed from *Polygonum amphibium* L., *P. bistorta* L., *P. viviparum* (L.) Ronse Decr., *Rumex acetosa* L., *R. acetosella* L., *R. aquaticus* (Trautv.) Hiitonen (TSHIKOLOVETS, 2011). Records of the larval food plant of the Lepidoptera (*Polygonum bistorta* L.) were reported in the same habitat near the location where the

species was collected (TANAP, 2014). This plant species is widely found in the high mountain steppe in the Eastern Anatolia region in Turkey. *Polygonum bistorta* ssp. *carneum* (K. Koch) is a mountain element of Blacksea region and spreads out through Middle and Eastern Blacksea sections, Erzurum-Kars section and Hakkari section (GÜNER, 2012). The fact that the larval food plant is widespread indicates that the necessary conditions exist for *L. helle* to settle and reproduce in the area.

*Lycaena helle*, which was identified in the monitoring studies carried out in the province of Ardahan in 2020, is the new record for Turkey. This is an important zoogeographic discovery regarding



**Figures 1-3.**– 1. Aerial photograph of the habitat where the samples were collected. 2. Habitus of *Lycaena helle* (D. & Schiff.). 3. Acid alpine and subalpine grassland (E4.4) habitat.

the spread of this species. The closest record of the species to Turkey is known as Georgia (Abkhazeti) (DIDMANIDZE, 2004). The available data in Europe show that the very limited habitat base of the species has been deteriorating at an alarming rate over the past two decades. New records of *L. helle* shift the known area of the species in Europe further to the south, which make them very important (POPOVIĆ *et al.*, 2014). The identification of the species in Turkey supports that the current distribution area is shifting towards the south.

The reason why only one individual belonging to *L. helle* was caught in this study is that it is a monitoring study carried out along the pipeline, which is a limited area. It is necessary to urgently carry out comprehensive studies in the field to determine the distribution area of the species and the population density. Otherwise, the only known population of the violet copper in Turkey may be on the verge of extinction, as in Europe, without being included in the scope of conservation studies due to insufficient data.

### Acknowledgment

We would like to thank Dr. Üzeyir Çağlar and doctoral student Şirin Bahar Can for their contribution to the field studies during the TANAP project monitoring studies.

## BIBLIOGRAPHY

- DAVIES, C. E., MOSS, D. & HILL, M. O., 2004.– EUNIS Habitat Classification Revised.– *European Environment Agency, European Topic Centre on Nature Protection and Biodiversity*: 1-307.
- BALINT, Z., 1993.– The threatened lycaenids of the Carpathian Basin, east-central Europe: 105-110.– *In* T. R. Ed.– *New Conservation Biology of Lycaenidae (Butterflies)*: 173 pp. The IUCN Species Survival Commission, Strasburg.
- BOZANO, G. C., 2004.– Lycaenidae part III. Subfamily Theclinae. Tribe Eumaeini (partim). *Satyrium, Superflua, Armenia, Neolycaena, Rhymmaria*.– *Guide to the butterflies of the Palaearctic Region*: 94 pp. Omnes Artes, Milano.
- BOZANO, G. C. & WEIDENHOFFER, Z., 2001.– Lycaenidae part I. Subfamily Lycaeninae.– *Guide to the butterflies of the Palaearctic region*: 62 pp. Omnes Art, Milano.
- BUSZKO, J. & MASŁOWSKI, J., 2008.– *Motyle dzienne Polski*: 274 pp. Koliber, Nowy Sacz.
- DIDMANIDZE, E., 2004.– Annotated list of diurnal butterflies (Lepidoptera: Rhopalocera) of Georgia and adjacent territory from Southern Caucasus.– *Proceedings of the Institute of Zoology*, **22**: 197-226.
- GÜNER, A., 2012.– *Türkiye Bitkileri Listesi, Damarlı Bitkiler*: 1290 pp. Nemaş, İstanbul.
- HABEL, J. C., MEYER, M. & SCHMITT, T., 2014.– *Jewels in the Mist: a synopsis on the endangered butterflies species the Violet Copper, Lycaena helle*: 235 pp. Pensoft, Sofia-Moscow.
- HABEL, J. C., RODDER, D., SCHMITT, T. & NÉVES, G., 2011.– Global warming will affect the genetic diversity and uniqueness of *Lycaena helle* populations.– *Global Change Biology*, **17**: 194-205.
- HELSDINGEN, P. J. van, WILLEMSE, L. & SPEIGHT, M. C. D., 1996.– Background information on invertebrates of the Habitats Directive and the Bern Convention. Part 1 Crustacea, Coleoptera and Lepidoptera.– *Nature and Environment*, **79**: 231 pp. Council of Europe Publishing, Strasbourg.
- KOLEV, Z. & SHTINKOV, N., 2015.– Notes on the distribution and conservation status of the Violet Copper *Lycaena helle* (Lepidoptera: Lycaenidae) in Bulgaria.– *Phegea*, **43**(1): 15.
- KORB, S. & BOLSHAKOV, L. V., 2016.– Systematic catalogue of butterflies of the former Soviet Union (Armenia, Azerbaijan, Belarus, Estonia, Georgia, Kyrgyzstan, Kazakhstan, Latvia, Lithuania, Moldova, Russia, Tajikistan, Turkmenistan, Ukraine, Uzbekistan) with special account to their type specimens (Lepidoptera: Hesperioidea, Papilionoidea).– *Zootaxa*, **4160**(1): 1-324.
- KUDRNA, O., HARPKE, A., LUX, K., PENNERSTORFER, J., SCHWEIGER, O., SETTELE, J. & WIEMERS, M., 2011.– *Distributions Atlas of Butterflies in Europe*: 576 pp. Gesellschaft für Schmetterlingsschutz eV, Germany.
- MARTIN, Y., HABEL, J. C., VAN DYCK, H. & TITEUX, N., 2014.– Losing genetic uniqueness under global change: the Violet Copper (*Lycaena helle*) in Europe.– *In* J. C. HABEL, M. MEYER & T. SCHMITT (eds). *Jewels in the mist. A synopsis on the endangered Violet Copper butterfly Lycaena helle*: 165-184. Pensoft, Sofia-Moscow.
- NEKRUTENKO, Y. & TSHIKOLOVETS, V., 2005.– *The Butterflies of Ukraine*: 231 pp. Rayevsky Scientific Publishers, Kyiv.
- POPOVIĆ, M., ĐURIĆ, M., FRANETA, F., VAN DEIJK, J. R. & VERMEER, R., 2014.– First records of *Lycaena helle* ([Denis & Schiffermüller], 1775) for the Balkan Peninsula (Lepidoptera: Lycaenidae).– *SHILAP Revista de lepidopterología*, **42**(166): 287-294.
- SKÓRKA, P., SETTELE, J. & WOYCIECHOWSKI, M., 2007.– Effects of management cessation on grassland butterflies in southern Poland.– *Agriculture, Ecosystems & Environment*, **121**: 319-324.
- VAN SWAAY, C. & WARREN, M. S., 1999.– *Red Data Book of European Butterflies (Rhopalocera)*, *Nature and Environment*, **99**: 260 pp. Council of Europe Publishing, Strasbourg.
- VAN SWAAY, C., CUTTELOD, A., COLLINS, S., MAES, D., LÓPEZ-MUNGUIRA, M., ŠAŠIĆ, M., SETTELE, J., VEROVNIK, R., VERSTRAEL, T., WARREN, M., WIEMERS, M. & WYNHOF, I., 2010a.– *European Red List of Butterflies*: X + 47 pp. Publications Office of the European Union, Luxembourg.
- VAN SWAAY, C., WYNHOF, I., VEROVNIK, R., WIEMERS, M., LÓPEZ-MUNGUIRA, M., MAES, D., SASIC, M., VERSTRAEL, T., WARREN, M. & SETTELE, J., 2010b.– *Lycaena helle*.– *In* IUCN 2012. *IUCN Red List of Threatened Species*. Version 2012.2. available from <http://www.iucnredlist.org>.
- TANAP, 2014.– *Trans Anadolu Doğal Gaz Boru Hattı (Tanap) Projesi Çevresel Etki Değerlendirmesi Raporu*: 3184 pp. Ankara.
- TSHIKOLOVETS, V., 2011.– *Butterflies of Europe & the Mediterranean area*: 544 pp. Tshikolovets Publications, Pardubice.

TUZOV, V. K., BOGDANOV, P. V., CHURKIN, S. V., DANTCHENKO, A. V., DEVYATKIN, A. L., MURZIN, D. J., SAMODUROV, G. D. & ZHDANKO, A. B., 2000.– *Guide to the Butterflies of Russia and Adjacent territories (Lepidoptera, Rhopalocera): Libytheidae, Danaidae, Nymphalidae, Rionidae, Lycaenidae*, 2: 580 pp. Pensoft, Moscow.

\*S. S. Ç.  
Gazi University  
Science Faculty  
Biology Department  
TR-06500 Ankara  
TURQUÍA / TURKEY  
E mail: selma@gazi.edu.tr  
E-mail: selmaseven@gmail.com  
<https://orcid.org/0000-0002-5839-3172>

A. H.  
Gazi University  
Science Faculty, Biology Department  
TR-06500 Ankara  
TURQUÍA / TURKEY  
E mail: hasbenli@gazi.edu.tr  
<https://orcid.org/0000-0002-2919-7594>

\*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 16-III-2021)  
(Revisado y aceptado / *Revised and accepted* 8-IV-2021)  
(Publicado / *Published* 30-III-2022)