

## Length-weight relationship for two species of river fish in the state of Mato Grosso of Sul, Brazil

Relação peso-comprimento para duas espécies de peixes de rios do estado do Mato Grosso do Sul, Brasil

Relación talla-peso para dos especies de peces de río en el estado de Mato Grosso do Sul, Brasil

Soinski TA, Costa MS, Brambilla EM, Smith WS. Length-weight relationship for two species of fish from Upper Paraguay Basin in Mato Grosso do Sul, Brazil. Rev Colombiana Cienc Anim. Recia. 2020; 12(1):e748. <https://doi.org/10.24188/recia.v12.n1.2020.748>

Universidad de Sucre, Colombia

Los autores permiten a RECIA reimprimir el material publicado en él. En caso de que un autor quiera traducir o usar una publicación parcial o completa de nuestro Diario, el autor debe obtener un permiso por escrito del editor de la revista.

Copyright © 2020. El (los) autor (es), Revista Colombiana de Ciencia Animal - RECIA. 2020. Este es un artículo de acceso abierto distribuido bajo los términos de Creative Commons Attribution 4.0 (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), El uso, distribución o reproducción está permitido, siempre que se acrediten al autor original y al propietario del copyright y que se cite la publicación original en esta revista, de acuerdo con la práctica académica aceptada. No se permite el uso, distribución o reproducción que no cumpla con estos términos.



# Length-weight relationship for two species of river fish in the state of Mato Grosso of Sul, Brazil

Relação peso-comprimento para duas espécies de peixes de rios do estado do Mato Grosso do Sul, Brasil

Relación talla-peso para dos especies de peces de río en el estado de Mato Grosso do Sul, Brasil

**Thais A. Soinski** M.Sc.

Universidade Paulista. Laboratório de Ecologia Estrutural e Funcional de Ecossistemas, Av. Independência, 210, Jardim Éden, CEP 18087-101, Sorocaba. Brasil.

Universidade de São Paulo. Escola de Engenharia de São Carlos/ EESC, Centro de Recursos Hídricos e Estudos Ambientais/CRHEA, Programa de Pós-Graduação em Ciências da Engenharia Ambiental, Rodovia Domingos Innocentini, km 13, Itirapina (SP), Brasil.

[thaissoinski@outlook.com](mailto:thaissoinski@outlook.com)

<https://orcid.org/0000-0003-4147-0212>

DOI: <https://doi.org/10.24188/recia.v12.n1.2020.748>

**Matheus S. Costa**

Universidade Paulista. Laboratório de Ecologia Estrutural e Funcional de Ecossistemas, Av. Independência, 210, Jardim Éden, CEP 18087-101, Sorocaba. Brasil.

[matheussouza\\_18075@hotmail.com](mailto:matheussouza_18075@hotmail.com)

<https://orcid.org/0000-0002-6182-0331>

**Eduardo M. Brambilla** Ph.D.

Universidade Estadual Paulista "Júlio de Mesquita Filho", Departamento de Zoologia, Botucatu, São Paulo, Brasil.

[eduardo.brambilla@gmail.com](mailto:eduardo.brambilla@gmail.com)

<https://orcid.org/0000-0003-2901-2526>

**Welber Senteio Smith** Ph.D.

Universidade Paulista. Laboratório de Ecologia Estrutural e Funcional de Ecossistemas, Av. Independência, 210, Jardim Éden, CEP 18087-101, Sorocaba. Brasil.

Universidade de São Paulo. Escola de Engenharia de São Carlos/ EESC, Centro de Recursos Hídricos e Estudos Ambientais/CRHEA, Programa de Pós-Graduação em Ciências da Engenharia Ambiental, Rodovia Domingos Innocentini, km 13, Itirapina (SP), Brasil.

Programa de Pós-Graduação em Aquicultura e Pesca, Avenida Francisco Matarazzo, 455, 05033-970, Parque da Água Branca, Barra Funda, São Paulo SP, Brasil.

[welber\\_smith@uol.com.br](mailto:welber_smith@uol.com.br)

<https://orcid.org/0000-0001-9803-7394>

Recepción: 15 febrero 2020

Aprobación: 4 mayo 2020

Publicación: 8 mayo 2020

## ABSTRACT

The study of the weight-length relationship of a fish species can be used to address several aspects that involve the distinction of small taxonomic units, providing basic information for the study of species biology, in addition, this relationship can also be used as quantitative indicator of the degree of health or well-being of the species in the environment. The objective of this study was to present unpublished data on the weight-length relationship (LWRs) for two species of fish. The fish were sampled in the months of January and July 2017 and January 2018 in rivers belonging to the Upper Paraná and Upper Paraguay basin along the MS-163, collected through sieves and trawls. 49 individuals were captured, distributed in two orders and two families, being 31 from *Hisonotus francirochai* and 18 from *Roeboides*

**Como citar (Vancouver).**

Soinski TA, Costa MS, Brambilla EM, Smith WS. Length-weight relationship for two species of fish from Upper Paraguay Basin in Mato Grosso do Sul, Brazil. Rev Colombiana Cienc Anim. Recia. 2020; 12(1):e748. <https://doi.org/10.24188/recia.v12.n1.2020.748>

*descalvadensis*. The results showed that the two species are within the expected for both parameters (a and b), even with a relatively low sample number. In addition, the data this study contribute with biological information for the species studied in this work.

**Keywords:** Freshwater fish, *Hisonotus francirochai*, rivers, *Roeboides descalvadensis*.

## RESUMO

O estudo da relação peso-comprimento de uma espécie de peixe pode ser utilizado para abordar diversos aspectos que envolvem a distinção de pequenas unidades taxômicas, contribuindo com informações básicas para o estudo da biologia das espécies, além disso, esta relação pode ser também usada como indicador quantitativo do grau de saúde ou bem-estar da espécie no ambiente. O objetivo deste trabalho foi apresentar dados não publicados sobre a relação peso-comprimento (LWRs) para duas espécies de peixes. Os peixes foram amostrados nos meses de Janeiro e Julho de 2017 e Janeiro de 2018 em rios pertencentes à bacia do Alto Paraná e Alto Paraguai ao longo da MS-163, coletados através de peneiras e rede de arrasto. Foram capturados 49 indivíduos, distribuídos em duas ordens e duas famílias, sendo 31 de *Hisonotus francirochai* e 18 de *Roeboides descalvadensis*. Os resultados mostraram que as duas espécies estão dentro do que é esperado para ambos os parâmetros (a e b), mesmo apresentando número amostral relativamente baixo. Além disso, os dados deste estudo contribuem com informações biológicas para as espécies estudadas neste trabalho.

**Palavras chaves:** Peixes de água doce, *Hisonotus francirochai*, rios, *Roeboides descalvadensis*.

## RESUMEN

El estudio de la relación peso-longitud de una especie de pez se puede utilizar para abordar varios aspectos que implican la distinción de pequeñas unidades taxonómicas, proporcionando información básica para el estudio de la biología de las especies, además, esta relación también se puede utilizar como indicador cuantitativo del grado de salud o bienestar de las especies en el medio ambiente. El objetivo de este estudio fue presentar datos no publicados sobre la relación peso-longitud (LWR) para dos especies de peces. Los peces fueron muestreados en los meses de enero y julio de 2017 y enero de 2018 en ríos pertenecientes a la cuenca del Alto Paraná y Alto Paraguay a lo largo de la MS-163, recolectados a través de tamices y redes de arrastre. Se capturaron 49 individuos, distribuidos en dos órdenes y dos familias, 31 de *Hisonotus francirochai* y 18 de *Roeboides descalvadensis*. Los resultados mostraron que las dos especies están dentro de lo esperado para ambos parámetros (ayb), incluso con un número de muestra relativamente bajo. Además, los datos de este estudio contribuyen con información biológica para las especies estudiadas en este trabajo.

**Palabras clave:** Peces de agua dulce, *Hisonotus francirochai*, ríos, *Roeboides descalvadensis*.

## INTRODUCTION

The correlation between weight and length includes characteristics selected for the population, providing very important information about the specific population (1,2). Among the most frequent uses of the weight and length ratio, the possibility of indirect weight determination by length or vice versa, the indication of the condition of the fish, in relation to fat storage or even the indirect analysis of growth (3). The knowledge of quantitative aspects, such as a weight / length ratio, is of great importance for a better understanding of the role of environmental characteristics in the ecology of species, as well as tools for studies of fisheries biology, threats to the management and conservation of stocks, in addition to population development (4).

Given the above, the objective of this study was to present unpublished data on the weight-length relationship (LWRs) of two species *Hisonotus francirochai* and *Roeboides descalvadensis* sampled in rivers along the MS-163 highway in Mato Grosso do Sul.

## MATERIAL AND METHODS

The state of Mato Grosso do Sul is drained by two sub-units of the Prata basin - the Upper Paraná basins to the east and the Paraguay basin to the west (5), belonging to the Upper Paraguay basin, one of the the main basins of the Central-West

Region of Brazil and to the Upper Paraná basin. The length-weight ratio is an important tool for describing fish growth and development, regardless of age (6). With this tool we can estimate the condition factor, used as an inference parameter of the health of the sample and its biomass in natural populations to analyze the relationships with the environmental conditions and the behavioral aspects of the species (7,8).

The samples were collected in January and July 2017 and January 2018 in rivers belonging to the state of Mato Grosso of Sul along the MS-163, using a 50 by 50 cm sieve and a 5 mm mesh, and a drag of three meters of mesh of 1 cm. After capture, the fish were euthanized in the field under the effect of the benzocaine anesthetic, after that they were weighed with the aid of on a digital balance (0.00001 g precision) and measured with the aid of a tape measure and analogical caliper (0.1 mm precision) and then placed in 10% formaldehyde solution for fixation. In the laboratory the individuals were transferred to 70% alcohol for preservation and identified with the aid of identification keys and specialized literature (5). In this way, species were identified, and all collected specimens vouchers were deposited in the fish collection at the Laboratório de Ecologia Funcional e Estrutural de Ecossistemas, Universidade Paulista, Sorocaba campus, São Paulo, Brazil (catalog numbers LEEF 0070-0071). All of the collections were carried out with authorization of the IBAMA license (Authorization N° 658/2015).

The length-weight ratio for both species was obtained based on Keys (9). Different types of length measurements can be used for length-weight relationship (LWR) studies. The most commons are total length, standard length and fork length. The type of length utilized vary according to study. In our case, we have chosen to use total length because is a measure that include the whole caudal fin and all the individuals sampled in this study presented this structure intact. In cases where the caudal fin presents damages, usually by predation or sampling methodologies, it is better use standard length. As example, the following studies have used total length to calculate LWRs: (10,11,12).

## RESULTS

A total of 49 specimens were collected, distributed in two orders (Siluriformes and Characiformes) and two families (Loricariidae and Characidae respectively), being 31 of *Hisonotus francirochai* and 18 of *Roeboides descalvadensis*. The parameters of LWRs are summarized in Table 1.

For *Hisonotus francirochai* there is no published formula for TLxSL, to transform TL (total length) into SL (standard length), so it was not possible to confirm that the value found in TL is greater than the SL record. For *Roeboides descalvadensis*, there is a formula published and disclosed by FishBase:  $TL = 0 + 1.182 \times SL$ , where the transformation was performed and the TL found is equivalent to 7.87 cm of SL, unfortunately smaller than the largest record (8.9 cm).

**Table 1.** Total length (cm) and weight (g) ratio for the two species sampled in rivers along the MS-163 highway in Mato Grosso do Sul, where  $W = aL^b$  and  $n$  = number of specimens used in the analysis,  $R^2$  = correlation coefficient. The length-weight ratios for the two species were significant in  $p < 0.0$ .

Family	Taxa	n	WT range (g)	TL range (cm)	a (95% CI)	b (95% CI)	R <sup>2</sup>
Loricariidae	<i>Hisonotus francirochai</i>	31	0.22-0.79	2.80-4.30	0.0096 (0.0062-0.0130)	2.97 (2.65-3.30)	0.923
Chacacidae	<i>Roeboides descalvadensis</i>	18	1.24-6.87	4.70-9.30	0.0116 (0.0059-0.0174)	2.88 (2.51-3.25)	0.944

## DISCUSSION

The values of parameter b of LWRs ranged from 2.88 to 2.97 (Table 1), which is adequate as demonstrated by Froese (13). Comparing the results with Fishes Bayesian LWR predictions (14), the two species are within what is expected for both parameters (a and b), even presenting slightly low values, which may be related to the smaller variation in size. The  $r^2$  values were considered acceptable for the species, even though they had a relatively low sample number (Table 1) (15).

Biometric variables and the type of growth of fish species can be influenced by several factors, such as food availability, population density, sex, physiology or abiotic factors that can affect the estimated values of use relationships (16,17,18,19). Therefore, more data, a larger sample size, is needed to confirm these growth values.

Although easily obtained, LWRs data are scarce for several species in the Neotropical region, thus, the results of this study contribute with biological information for the species studied in this work.

## REFERENCES

1. Giarrizzo T, Oliveira RRS, Andrade MC, Gonçalves AP, Barbosa TAP, Martins AR, et al. Relações peso-comprimento e comprimento-comprimento para 135 espécies de peixes do rio Xingu (bacia amazônica, Brasil). *J Appl Ichthyol.* 2015; 31(2):514-424. <https://doi.org/10.1111/jai.12677>
2. Baitha R, Sinha A, Koushlesh SK, Chanu TN, Kumari K, Gogoi P, et al. Relação comprimento-peso de dez espécies de peixes de água doce indígenas do rio Gandak, Bihar, Índia. *J Appl Ichthyol.* 2018; 31(1):233-236. DOI: <https://doi.org/10.1111/jai.13555>
3. Agostinho AA, Gomes LC. Reservatório de segredo: bases ecológicas para o manejo, Maringá: EDUEM; 1997.
4. Lizama MA, Ambrósio AM. Crescimento, recrutamento e mortalidade do pequi *Moenkhausia intermedia* (Osteichthyes, Characidae) na planície de inundação do alto rio Paraná, Brasil. *Acta Sci Biol Sci.* 2003; 25(2):329-333. <https://doi.org/10.4025/actasciobiolsci.v25i2.2020>
5. Froehlich O, Cavallaro M, Sabino J, Suárez YR, Alencar VMJ. Checklist da ictiofauna do Estado de Mato Grosso do Sul, Brasil. *Iheringia Sér Zool.* 2017; 107(Supl):e2017151. <http://dx.doi.org/10.1590/1678-4766e2017151>
6. Lemos JRG, Tavares-Dias M, Marcon JL, Lemos PEM, Affonso EG, Zaiden SF. Relação peso-comprimento e fator de condição em espécies de peixes ornamentais do rio Negro, Estado do Amazonas, Brasil. *CIVA*; 2006.
7. Gomiero LM, Braga FMS. 2005. The condition factor of fishes from two river basins in São Paulo state, Southeast of Brazil. *Acta Sci Biol Sci.* 2005; 27(1):73-78. <https://doi.org/10.4025/actasciobiolsci.v27i1.1368>
8. Camara EM, Caramaschi EP, Petry AC. Fator de condição: Bases conceituais, aplicações e perspectivas de uso em pesquisas ecológicas com peixes. *Oecologia Australis.* 2011; 15(2):249-274. <https://doi.org/10.4257/oeco.2011.1502.05>
9. Keys AB. The length-weight relation in fishes. *PNAS.* 1928; 14(12):922-925. <https://doi.org/10.1073/pnas.14.12.922>
10. Koutrakis ET, Tsikliras AC. Length-weight relationships of fishes from three northern Aegean estuarine systems (Greece). *J Appl Ichthyol.* 2003; 19(4):258-260. <https://doi.org/10.1046/j.1439-0426.2003.00456.x>
11. Oliva-Paterna FJ, Torralva M, Carvalho ED. Length-weight relationships for 20 species collected in the Jurumirim reservoir (Paranapanema Basin, Brazil). *J Appl Ichthyol.* 2009; 25(3):360-361. <https://doi.org/10.1111/j.1439-0426.2009.01194.x>
12. Moutopoulos DK, Stergiou KI. Length-weight and length-length relationships of fish species from the Aegean Sea (Greece). *J Appl Ichthyol.* 2002; 18(3):200-203. <https://doi.org/10.1046/j.1439-0426.2002.00281.x>
13. Froese R. Cubelaw, condition factor and weight-length relationships: History, meta-analysis and recommendations. *J Appl Ichthyol.* 2006; 22(4):241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>
14. Froese R, Thorson JT, Reyes Jr RB. A Bayesian approach for estimating length-weight relationships in fishes. *J Appl Ichthyol.* 2014; 30(1):78-85. <https://doi.org/10.1111/jai.12299>
15. Azevedo Santos VM, Coelho PN, Brambilla EM, Lima FP, Nobile AB, Britton JR. Length-weight relationships of four fish species from the upper Paraná River basin, Southeastern Brazil. *J Appl Ichthyol.* 2018; 34(1):237-239. <https://doi.org/10.1111/jai.13542>
16. Silva LMA, Oliveira MSB, Florentino AC, Tavares-Dias M.. Relação comprimento-peso de 11 espécies de peixes de um afluente do sistema do rio Amazonas no norte do Brasil. *J Appl Ichthyol.* 2015; 31(4):816-817. <https://doi.org/10.1111/jai.12791>

17. Cella-Ribeiro A, Hauser M, Nogueira LD, Doria CRC, Torrente-Vilara G. Relações peso-comprimento de peixes do rio Madeira, Amazônia brasileira, antes da construção de usinas hidrelétricas J Appl Ichthyol. 2015; 31(5):939-945. <https://doi.org/10.1111/jai.12819>
18. Dieb-Magalhães L, Florentino AC, Soares MGM. Relações comprimento-peso e comprimento na primeira maturidade para nove espécies de peixes de lagos de várzea na Amazônia Central (bacia Amazônica, Brasil). J Appl Ichthyol. 2015; 31(6):1182-1184. <https://doi.org/10.1111/jai.12919>
19. Freitas TMS, Prudente BS, Montag LFA.. Relação comprimento-peso em dez espécies de peixes do rio Nhamundá, na Bacia Amazônica, Brasil. *Acta Amazonica*. 2017; 47(1):75-78. <https://doi.org/10.1590/1809-4392201601272>