



**The vinasse and yellow trap as an alternative in the ecological control of radish  
(*Raphanus sativus L.*). A practical approach**

**La vinaza y trampa amarillo una alternativa en el control ecológico del rabanito  
(*Raphanus sativus L.*). Una aproximación práctica**

Legua-Cárdenas José Antonio\*<sup>ID</sup>, Cruz-Nieto Dante Daniel<sup>ID</sup>, Nunja-García José Vicente<sup>ID</sup>, Caro-Soto Félix Gil<sup>ID</sup>

**Article Data**

José Faustino Sánchez Carrión National University.  
Mercedes Indacochea Avenue N° 600.  
Huacho - Huaura.  
Peru.  
Tel: +511 232 2918.

\* Contact address:

José Antonio Legua-Cárdenas

José Faustino Sánchez Carrión National University.  
Mercedes Indacochea Avenue N° 600.  
Huacho - Huaura.  
Peru.  
Tel: +511 232 2918.

E-mail: [jlegua@unjsc.edu.pe](mailto:jlegua@unjsc.edu.pe)

**Abstract**

The objective was to determine the dose for whitefly control. The methodology is based on applied research, so the Completely Randomized Block Design was implemented, consisting of 3 blocks and 5 treatments which were: T<sub>1</sub> 0, T<sub>2</sub> 10, T<sub>3</sub> 20, T<sub>4</sub> 30 and T<sub>5</sub> with 40 mL of vinasse/1.6 m<sup>2</sup>, these were applied from the first week to the fourth week with doses of 10 mL of vinasse/32 mL of water/week/1.6 m<sup>2</sup>. The physical characteristics of the crop, the number of flies trapped in the trap and the percentage of damage to the radish crop were also evaluated and statistically evaluated using the analysis of variance and Duncan's statistical method. It was determined that the T<sub>5</sub> treatment with 40 mL/1.6m<sup>2</sup> excelled in the most optimal results, which were the following values: Crops affected:31.67 %, whiteflies per trap:7, crop size: 26.58 cm, plant weight:37.26 g, agricultural yield:7.008 t/ha and bulb diameter:3.15 cm. It is concluded that the application of vinasse is an agro ecological alternative for whitefly control, with the best results being observed with treatment T<sub>5</sub>.

**Keywords:**

Vinasse,  
yellow trap,  
white fly,  
radish,  
performance.

***J. Selva Andina Biosph.***  
**2022; 10(1):46-56.**

**Article ID: 113/JSAB/2021**

**Article history.**

Received November, 2021.  
Returned January, 2022.  
Accepted March, 2022.  
Available online, May 2022.

**Edited by: Selva Andina  
Research Society**

**Palabras clave:**

Vinaza,  
trampa amarilla,  
mosca blanca,  
rabanito,  
rendimiento.

**2022. *Journal of the Selva Andina Biosphere*®. Bolivia. All rights reserved.**

**Resumen**

El objetivo fue determinar la dosis para el control de mosca blanca. La metodología se basa en investigación aplicada, por lo que se implementó el Diseño de Bloques Completamente al Azar que constaron de 3 bloques y 5 tratamientos los cuales fueron: T<sub>1</sub> con 0, T<sub>2</sub> con 10, T<sub>3</sub> con 20, T<sub>4</sub> con 30 y T<sub>5</sub> con 40 mL de vinaza/1.6 m<sup>2</sup>, estas se aplicaron desde la primera semana hasta la cuarta semana con dosis de 10 mL de vinaza/32 mL de agua/ semana/1.6 m<sup>2</sup>. Asimismo, se evaluaron las características físicas del cultivo, cantidad de moscas atrapadas en trampa y porcentaje de daño al cultivo rabanito, con esta información se evaluó estadísticamente, utilizando el análisis de varianza y método estadístico de Duncan. Se determinó que el tratamiento T<sub>5</sub> con 40 mL/1.6m<sup>2</sup> sobresalió en los resultados más óptimos que fueron los valores siguientes: Cultivos afectados:31.67 %, moscas blancas por trampa:7, tamaño de cultivo: 26.58 cm, peso de planta:37.26 g, rendimiento agrícola:7.008 t/ha y diámetro de bulbo:3.15 cm. Se concluye que la aplicación de vinaza es una alternativa agroecológica para el control de la mosca blanca, observándose los mejores resultados con el tratamiento T<sub>5</sub>.

**2022. *Journal of the Selva Andina Biosphere*®. Bolivia. Todos los derechos reservados.**



## Introduction

For many years, sugarcane organic wastes such as bagasse, vinasse and other wastes have not been optimally treated or disposed of in the environment. As a result, this causes harm to public health and negative environmental impact. Research has shown that industrial activities generate emissions of particles such as soot and smoke due to the use of fuel oil and bagasse from the burning of sugar cane<sup>1</sup>. This situation affects health, as the Ministry of Health has shown that inadequate management of these residues has negative effects on health, ranging from simple laryngitis to more complex diseases such as cancer, neuropsychiatric disorders and vascular diseases worldwide<sup>2</sup>.

Vinasse, an effluent derived from sugarcane residues (SR), obtained from large volumes of alcohol distillation, whose final disposal is not adequately treated, on average, 10 to 15 L of vinasse/L of alcohol produced are generated<sup>3</sup>. These fresh by-products of the sugar industry (cachaza, bagasse and vinasse) incorporated into the soil generate a negative impact on plants<sup>4</sup>. There is another assessment that explains that cachaza and vinasse, residues from the sugar agroindustry, have a negative impact when discharged into bodies of water and a positive impact when applied to the soil, the application of vinasse mainly benefits the chemical and biological properties of the soil, they conclude that vinasse benefits the soil by providing a large amount of K and organic matter (OM)<sup>5</sup>. Due to this problem, it is necessary to develop an alternative solution, which could be its use as a complement in the preparation of compost, or its application in pest control, due to its chemical and physical characteristics. It should also be mentioned that these effluents are characterized by high temperatures, acid pH, high COD (chemical organic demand) and total solids<sup>6</sup>.

Hence, the incorporation of SR such as cachaza, bagasse, vinasse, has favored agriculture, whether in the preparation of compost, due to its concentration of nutrients, such as nitrogen and microelements, which supplement their availability in the soil, reducing environmental pollution and fertilizer costs. The applications of pure vinasse as an organic amendment, produce marked residual effects, positively influencing chemical properties, increasing assimilable OM, phosphorus and potassium contents, which made it possible to dispense with mineral fertilization and maintain high agricultural yields<sup>7</sup>. Likewise, the disposal of composted and/or biodigested vinasse in soils is a way of reducing the costs of distribution and application of these and at the same time adding a certain value, in carrying out various treatments prior to their application<sup>8</sup>.

It has been mentioned that there is an important need to carry out research on the biocidal effect of agro-industrial and livestock by-products as a basis or complement for the registration of a series of biopesticides, which would be very useful in agriculture<sup>9</sup>. It should be noted that vinasse, due to its properties, can be used as a sustainable ecological alternative for pest control in vegetables, either because of its expandable odor, acid pH, as a repellent for some insects, which are attracted by the yellow color and get trapped in traps installed for this purpose, thus reducing the cost of production, obtaining ecological fruit and at the same time reducing environmental pollution. Currently, the biocidal effect of agro-industrial and livestock by products will make it possible to comply with international environmental and health commitments. The use of vinasse in aerobic fermentation processes is feasible because it is rich in carbon and some salts such as potassium and calcium, making it an important source of microbial growth<sup>10</sup>. Its

use for the ecological control of pests, as a substrate for the propagation of microorganisms such as *Trichoderma* spp. can have effects on some diseases or nematodes, reducing damage to the crop and increasing its yield<sup>11</sup>. The study of its physicochemical properties, its importance if biological treatments are to be used<sup>12</sup>. Therefore, the dose of vinasé can reduce the damage of the white fly (*Trialeurodes vaporariorum*), a pest of economic importance in vegetables, compared to synthetic insecticides that have generated resistance with adverse effects for the environment. The high sugars and organic matter of the SR (vinasé), could be used as substrates to favor the

multiplication of beneficial microorganisms such as *Trichoderma* spp.<sup>13</sup>.

Therefore, this research was carried out on the use of vinasé as an ecological alternative for pest control and to reduce the damage caused by the white fly in vegetables in the area.

## Materials and methods

*Location of the experiment.* The research was carried out in the district of Barranca, province of Barranca, located in the Lima region, temperature 23 to 25° C, relative humidity 85 to 88 %, sandy loam soil type<sup>14</sup>.

**Table 1 Treatment of vinasé doses**

Treatments	Doses	TA	Application timing (mL/L of water/ha/week)				TH
			Week 1	Week 2	Week 3	Week 4	
T <sub>1</sub>	0	0	0	0	0	0	0.00
T <sub>2</sub>	10	10	62.5/200	0	0	0	62.5
T <sub>3</sub>	10	20	62.5/200	62.5/200	0	0	125.0
T <sub>4</sub>	10	30	62.5/200	62.5/200	62.5/200	0	187.5
T <sub>5</sub>	10	40	62.5/200	62.5/200	62.5/200	62.5/200	250.0

Dose: mL/32 mL water/week, TA Total: mL/1.6 m<sup>2</sup>, TH Total application: L./ha  
The water used to dilute the vinasé was groundwater.

A Completely Randomized Block Design (CRBD) was used, consisting of 3 blocks and 5 treatments which were: T<sub>1</sub> with 0, T<sub>2</sub> with 10, T<sub>3</sub> with 20, T<sub>4</sub> with 30 and T<sub>5</sub> with 40 mL of vinasé/1.6 m<sup>2</sup>, taking into account a control treatment and the vinasé was dosed at a rate of 10 mL/32 mL of water/week (Table 1), applied with doses of 8 mL/24 mL of water/1.6 m<sup>2</sup>/week<sup>15</sup>, to reduce or eliminate the action of the earthworm, the dose used was proportional to that used<sup>16</sup>. Radish seed was sown at distances of 0.10 m between plants and 0.50 m between rows, and yellow traps were placed in all the demonstration plots, smeared with glue for traps. Data were recorded from sowing to harvesting of the radish crop in each plot and the number of whiteflies stuck in the yellow traps.

The application of the vinasé doses of 10 mL of vinasé/32 mL of water/1.6 m<sup>2</sup> from the first week to

the fourth week, according to Table 1. Experiment procedures. Field preparation was carried out in a conventional manner, i.e., the way it is done by farmers in the area. Soil samples were taken from the experimental area in a staggered manner and only 1 kg was taken to the National Institute for Agrarian Innovation (INIA) for evaluation<sup>17</sup>. The physical characteristics of the radish crop were then evaluated, such as: plant size (TS), plant weight (PW), commercial yield (CY) and equatorial bulb diameter (EBD), and the data were processed using analysis of variance and Duncan's test at 5 % error. Finally, yellow traps were placed in each plot to attract and trap the insects; the trap measures 40 cm long by 20 cm wide. Five days after planting and at harvest time, the insects stuck in each trap were counted. Once the data were obtained, they were processed by statistical

analysis. Analysis of variance. The experimental information was processed with the analysis of variance, which allowed determining if there was an effect of the vinasse dose on the physical characteristics and on the ecological control of this pest.

*Duncan's test.* This statistical operation was performed when it was determined that there was statistical significance in the physical characteristics of the crop.

**Table 2 Basic soil fertility analysis for radish crop<sup>18</sup>**

E.C. mS/cm 1:2.5	pH (1:2.5)	O.M. (%)	N (%)	P (ppm)	K (ppm)	CaCO <sub>3</sub> (%)	Cation exchange (mEq/100 g soil)				CEC-E
							Ca	Mg	Na	K	
1.19	6.88	1.37	0.07	12	212	1.76	16.99	0.68	0.29	0.54	18.50

EC: electrical conductivity, OM: organic matter, mEq: milliequivalent, CEC-E: cation exchange capacity, ppm: parts per million.

The concentration of microelements, low values of copper and boron, normal iron, zinc in excess, so it was necessary to apply organic matter to supply these elements (Table 3).

**Table 3 Trace element concentrations in the area of the experiment<sup>19</sup>**

Microelements			
Fe (ppm)	Cu (ppm)	Zn (ppm)	Bo (ppm)
103.90	7.62	11.04	1.56
Normal	Low	Excess	Low

*Characteristics of the vinasse.* Percentage of total solids 8.2 %, total ashes 3.99 % and others. In nutrients, a concentration of 2.35 mg of nitrogen was obtained. Also in the physical characteristics a pH of 4.92 was obtained, electrical conductivity a value of 365.7 mS/cm. (Table 4).

Percentage of plants affected. Affected by whitefly per week, is shown in Table 5, T<sub>5</sub> with 31.67 %.

*Statistical analysis of the number of whiteflies per yellow trap.* Table 6, there was no significance. Treatment T<sub>5</sub> with 7 whiteflies attached to the trap stood out in relation to the other treatments.

## Results

*Soil analysis:* neutral pH (6.8 - 7.3), low MO concentration (2-4 %), nitrogen (0.1-0.2 %), phosphorus (12-36 ppm), potassium (12-250 ppm), calcium carbonate medium concentration and no danger of salts (Table 2).

**Table 4 Physical and chemical characteristics of vinasse<sup>15</sup>**

Parameters	Value
Total solids (%)	8.2
Total ash (%)	3.99
Crude fat (%)	1.33
Total nitrogen (mg L <sup>-1</sup> )	2.35
Crude fibre (%)	1.97
pH	4.92
Electrical conductivity (mS cm <sup>-1</sup> )	36.7
Brix degrees (°Bx)	10.0

*Plant size.* The TP is detailed in Table 7, it indicates that there were no statistical differences, its coefficient of variation was 12.21 %, T<sub>5</sub> with 25.68 cm stood out in relation to the other treatments.

*Plant weight.* The PP shown in Table 7 shows that there was no significance in the treatments and it is observed that T<sub>5</sub> with 37.26 g obtained higher weight in relation to the others.

*Commercial yield.* Table 7 shows that there were no statistical differences, however, T<sub>5</sub> with 7.008 t/ha obtained the highest yield. *Equatorial diameter.* As shown in Table 7, there was no effect of dose in the treatments, T<sub>5</sub> with 3.15 cm differed from the others.

**Table 5 Percentage of Plants Affected by Whiteflies**

Treatments	TA	Percentage of damaged plants				Average (%)
		Week 1	Week 2	Week 3	Week 4	
T <sub>1</sub>	0	33.33	40.00	53.33	66.67	48.33
T <sub>2</sub>	10	30.00	36.67	46.67	60.00	43.33
T <sub>3</sub>	20	26.67	33.33	43.33	53.33	39.17
T <sub>4</sub>	30	20.00	30.00	40.00	56.67	36.67
T <sub>5</sub>	40	16.67	23.33	36.67	50.00	31.67

TA Total, of vinasé application (mL/1.6 m<sup>2</sup>)**Table 6 Statistical analysis of number of whiteflies per yellow trap**

Treatments	TA	Q
T <sub>5</sub>	40	7.00 a
T <sub>4</sub>	30	6.33 a
T <sub>3</sub>	20	5.00 a
T <sub>2</sub>	10	4.00 a
T <sub>1</sub>	0	3.00 a
Significance		**
Coefficient of variation %		43.31

TA Total, of vinasé application (mL/1.6 m<sup>2</sup>)Q Quantity (N whiteflies/trap Yellow fly/1.6 m<sup>2</sup>)**Table 7 Physical characteristics of the radish crop**

Treatments	Total	PS	PW	CY	BED
T <sub>5</sub>	40	26.58 a	37.26 a	7.008 a	3.15 a
T <sub>4</sub>	30	25.79 a	34.10 ab	6.137 a	3.02 ab
T <sub>3</sub>	20	25.36 a	32.18 ab	5.403 a	2.64 ab
T <sub>2</sub>	10	24.69 a	28.01 ab	4.667 a	2.40 ab
T <sub>1</sub>	00	22.67 a	26.33 b	4.240 a	2.34 b
Significance		**	**	**	**
Coefficient of variation		12.21	14.90	38.41	14.52

Total, of vinasé application (mL/1.6 m<sup>2</sup>), CY commercial yield (t/ha),  
TP Plant size (cm), PP Plant weight (g), DE Bulb equatorial diameter (cm)

## Discussion

According to the results, the percentage of affected plants shown in Table 5, it can be seen that the T<sub>5</sub> treatment with the lowest level of affection that registered a value of 31.67 %, compared to the rest of the treatments. Therefore, at this dose of vinasé, a control effect on the propagation of this insect was verified, showing a lower level of damage (deterioration of the plant, reflected mainly in the leaves and

in the agricultural yield, due to the action of the whitefly). This is due to the fact that the dose of higher concentration of vinasé and consequently also higher volatilization of its components, causes an intense odor that makes a hostile effect to the white fly to the radish crop, causing less damage to the plant and increasing the agricultural yield, in this regard Senatore et al.<sup>20</sup>, indicate, with the purpose of reducing the resistance of this insect, by the excess-

sive use of synthetic insecticides, environmental pollution and the cost of production, the application of vinasse and its different effects on the microbiota of the soil is important. It is important to mention that vinasse is the main by-product resulting from the production of bioethanol, it is a dark brown effluent with a strong odor and acid pH. This is due to its high contamination power, which is favored by substances such as: phytotoxins, antibacterials and recalcitrant compounds such as phenols, polyphenols and heavy metals, which have negative effects on microorganisms and plants<sup>21</sup>. The vinasse is composed of organic materials and mineral nutrients, which are part of plant compounds and constituents such as amino acids, proteins, lipids, various acids, enzymes, sugars and hormones<sup>22</sup>. With respect to the number of whiteflies per yellow trap, which is detailed in Table 6, it can be seen that there was no significance, i.e. the applications of vinasse did not have a statistical influence on the control of this insect, however, the highest dose, which is treatment T<sub>5</sub> with 7 whiteflies trapped, stood out in relation to the other treatments. It is an acid liquid (pH 3.7 to 5), with a characteristic odor, whose coloration varies from amber yellow to dark brown and which presents high temperature when produced<sup>23</sup>. The vinasse dose of 250 m<sup>3</sup> ha<sup>-1</sup> increased the K content in the cachaza compost and increased the P content<sup>24</sup>. However, a problem that persists is that the possible agronomic application of vinasse (both in the field and in experimental conditions) is its intense and persistent odor due to the presence of melanoidins<sup>25</sup>. In the evaluations of the physical characteristics of the radish crop such as PS, Table 7, the applications of vinasse did not influence, however, T<sub>5</sub> with 26.58 cm stood out in comparison

to the others. At this dose of vinasse, greater availability of nutrients such as nitrogen, potassium and other micronutrients was obtained, and its physical characteristics mentioned influenced in repelling insects, which affected the development of PH, presenting good architecture. The application of vinasse in soil and its possible agricultural use in coriander (*Coriandrum sativum*), in the length of stems treated with 20 % v/v vinasse are greater than those obtained in T<sub>0</sub>. The treatments with 5.152 and 7.879 L/ha had the greatest height<sup>26</sup>. The application of fortified vinasse (T<sub>3</sub>) showed the best vegetative growth 30 days after sowing, surpassing by 25 cm in height on average all the treatments in corn crops<sup>27</sup>. It was investigated that Spirulina and vinasse have properties that favor the growth and development of crops, thus improving their nutritional quality<sup>28</sup>. The PW, Table 7, did not show statistical significance, it was also observed that the T<sub>5</sub> treatment with 37.26 g stood out in comparison to the other treatments. Therefore, it can be interpreted that there was an effect of the dose of vinasse on the PW, the dose of vinasse influenced the higher PW, at this dose, concentrations of nitrogen, phosphorus, potassium and microelements were added to the soil, and the greater availability of these elements promoted the absorption of other nutrients that influenced the development of the plant and therefore its weight. It should be noted that the contents of assimilable phosphorus and potassium in the soil benefited from the application of 60 m<sup>3</sup> ha<sup>-1</sup> of vinasse; the pH was not influenced by the application of this product. Likewise, the results obtained allow us to identify the potential of vinasse as an alternative for fertilization<sup>29</sup>. In relation to the application of vinasse for fertilizer purposes, some authors maintain that there was a positive effect observed during the

application of vinasse, increasing total organic carbon and inorganic nitrogen, which was used by corn plants<sup>30</sup>. The CY, Table 7, indicates that there was no significance, in other words, there was no effect of vinasse application, but it should be noted that T<sub>5</sub> with 7.008 t/ha obtained the highest yield compared to the other treatments. This result indicates that at this dose of vinasse, nutrients such as nitrogen, phosphorus, potassium and other nutrients were incorporated into the soil, which increased the availability and greater absorption of these elements that influenced the development of the plant and therefore the yield. In relation to the favorable effects of the application of vinasse in agriculture, there is information that the composting of vinasse effluent in organic residues of sugarcane bagasse, in order to increase the nitrogen in the final composition of this fertilizer, guano from guinea pigs was incorporated into this mixture, also taking into account the basic analysis of soil fertility, for the radish crop (*R. Sativus, L.*)<sup>31</sup>. Another study treated SR, with the objective of using it as a fertilizer to minimize the use of synthetic fertilizers. To obtain this fertilizer, compost was made with 16.5 kg of stubble, 16.5 kg of guano from guinea pigs, 11 kg of bagasse and 11 liters of vinasse, the statistical analysis applied was analysis of variance and Duncan, it was determined that T<sub>4</sub> excelled in yield with 15.39 tn/ha, PW 44.66 g, PL 25.16 cm, DE 3.60 cm, polar diameter with 4.80 cm, RL 10.35 cm<sup>32</sup>. In ED Table 7, it was determined that there was no dose effect of vinasse applications in the treatments. However, T<sub>5</sub> with 3.15 cm stood out compared to the other treatments. This result analyzes that this dose of vinasse influenced the bulb thickness, hence with this dose of vinasse nutrients such

as nitrogen, phosphorus, potassium and other micro-nutrients were incorporated into the soil, which increased the availability and greater efficiency of absorption of these elements that influenced the development of the plant and therefore the thickness of the fruit. It is worth mentioning that many studies have concluded that the use of vinasse is an organic alternative to reduce inorganic (chemical) fertilization, mainly due to its high nutritional content at low cost and soil chemical conditions<sup>33</sup>. The vinasse used rationally fully satisfies the needs of N, K and S, reducing fertilizer costs<sup>34</sup>. It is concluded that the current application of vinasse contributes to improve productivity by increasing crop biomass.

## Source of financing

The present study was partially financed by the Universidad Nacional José Faustino Sánchez Carrión, with access to information and laboratory analysis.

## Conflicts of interest

There is no conflict of interest in the development of the research.

## Acknowledgments

The authors would like to thank the Universidad Nacional José Faustino Sánchez Carrión for logistical support.

## Ethical considerations

For the development of this research, the established norms regarding ethics and morality were followed.

## Research limitations

Financial difficulties and restrictions were encountered in finding similar research to ascertain the state of knowledge of our research.

## Authors' contribution

*José Antonio Legua Cárdenas*, statistical processing and data analysis and interpretation. *Dante Daniel Cruz Nieto*, collaborated in the field data collection and conduct of all the technical aspects of the research. *Jose Vicente Nunja García*, participated in the critical interpretation of the information from the physical-chemical analysis and *Félix Gil Caro Soto*, in the revision of style and writing of the scientific article.

## Cited Literature

- Bustamente Marín CK, Hernández Suárez BMR (dir). Análisis de la situación actual de las emisiones del Ingenio Central Progreso, Veracruz [tesis licenciatura]. [Veracruz]: Universidad Veracruzana; 2014 [citado 26 de octubre de 2021]. Recuperado a partir de: <https://cdigital.uv.mx/handle/123456789/46736>
- Tovar Artunduaga M, Losada Salazar GM, García TF. Impacto en la salud por el inadecuado manejo de los residuos peligrosos. Ingenierías USBMed 2015;6(2):46-50. DOI: <https://doi.org/10.21500/20275846.1731>
- Torres Gaviria LF, Ocampo Vélez JC, Socarrás Cárdenas A. Reducción del nivel de potasio en vinaza de destilería utilizando resinas de intercambio iónico. Rev Investig Agrar Ambient 2019;10(1):107-18. DOI: <https://doi.org/10.22490/21456453.2177>
- Bohórquez A, Puentes YJ, Menjivar JC. Evaluación de la calidad del compost producido a partir de subproductos agroindustriales de caña de azúcar. Cienc Tecnol Agropecu 2014;15(1):73-81. DOI: [https://doi.org/10.21930/rcta.vol15\\_num1.art:398](https://doi.org/10.21930/rcta.vol15_num1.art:398)
- Quiroz Guerrero I, Pérez Vázquez A. Vinaza y compost de cachaza: efecto en la calidad del suelo cultivado con caña de azúcar. Rev Mex Cienc Agríc 2013;4(Suppl 5):1069-75. DOI: <https://doi.org/10.29312/remexca.v0i5.1313>
- Sandoval Rojas ME, Hernández Muñoz AF (dir), Houbron E (dir). Tratamiento de vinazas provenientes de etanol en un reactor de lecho fluidizado inverso [tesis doctoral]. [Madrid]: Universidad Politécnica de Madrid; 2015 [citado 16 de octubre de 2021]. Recuperado a partir de: <https://oa.upm.es/40681/>
- Armengol JE, Lorenzo R, Fernández N. Utilización de la vinaza como enmienda orgánica y su influencia en las propiedades químicas de vertisoles y en los rendimientos de la caña de azúcar. Cultrop 2003;24(3):67-71.
- Santos M, Martín F, Diánez F, Carretero F, García-Alcázar M, de Cara M, et al. Efecto de la aplicación de vinaza de vino como biofertilizante y en el control de enfermedades en el cultivo de pepino [Internet]. Almería: Universidad de Almería; 2007 [citado 12 de octubre de 2021]. 9 p. Recuperado a partir de: [http://webantigua.agroecologia.net/recursos/publicaciones/publicaciones-online/2009/eventos-seae/cds/congresos/actas-bullas/seae\\_bullas/verd/sesiones/16%20S4CSANIDAD%20\(III\)/S4C8.pdf](http://webantigua.agroecologia.net/recursos/publicaciones/publicaciones-online/2009/eventos-seae/cds/congresos/actas-bullas/seae_bullas/verd/sesiones/16%20S4CSANIDAD%20(III)/S4C8.pdf)
- Valeiro A, Portocarrero R, Ullivarri E, Vallejo J. Los residuos de la Industria Sucro-alcoholera Argentina [Internet]. Buenos Aires: Instituto Nacional de Tecnología Agropecuaria; 2017 [citado 2 de

- octubre de 2021]. 16 p. Recuperado a partir de: [https://inta.gob.ar/sites/default/files/inta\\_residuos\\_sucro\\_alcoholera\\_argentina.pdf](https://inta.gob.ar/sites/default/files/inta_residuos_sucro_alcoholera_argentina.pdf)
10. Campos CR, Mesquita VA, Silva CF, Schwan RF. Efficiency of physicochemical and biological treatments of vinasse and their influence on indigenous microbiota for disposal into the environment. *Waste Manag* 2014;34(11):2036-46. DOI: <https://doi.org/10.1016/j.wasman.2014.06.006>
11. Carbajo Romero MS, Ojeda Fermoselle AC, Meneguzzi N, Canteros BI, Rodríguez G. Comportamiento de hongos en medios con vinaza de caña de azúcar. En: Monetta P, Paroldi HE, Miguel RE, editores. II Simposio de Residuos Agropecuarios y Agroindustriales del NOA y Cuyo. Instituto Nacional de Tecnología Agropecuaria [Internet]. San Juan: Instituto Nacional de Tecnología Agropecuaria; 2019 [citado 3 de septiembre de 2021]. p. 47-50. Recuperado a partir de: <https://inta.gob.ar/documentos/los-residuos-de-la-industria-sucro-alcoholera-argentina>
12. Aristizábal Alzate CE. Caracterización físico-química de una vinaza resultante de la producción de alcohol de una industria licorera, a partir del aprovechamiento de la caña de azúcar. Ing. USBMed 2015;6(2):36-41. DOI: <https://doi.org/10.21500/20275846.1729>
13. Aguilar-Astudillo E, Rodríguez-Hernández C, Bravo-Mojica H, Soto-Hernández RM, Bautista-Martínez N, Guevara-Hernández F. Repelencia de adultos de mosca blanca *Trialeurodes vaporariorum* (Heteroptera: Aleyrodidae) con clavo y pimienta. *Rev Colomb Entomol* 2020;46(2): e7520. DOI: <https://doi.org/10.25100/socolen.v46i2.7520>
14. Senatore D. Vinaza como fertilizante de caña azucarera: efectos sobre la comunidad bacteriana del suelo [tesis maestría]. [Montevideo]: Universidad la República; 2013 [citado 16 de septiembre de 2021]. Recuperado a partir de: <https://www.colibri.udelar.edu.uy/jspui/handle/20.500.12008/6446>
15. Gálvez Torres E, Legua Cárdenas J, Cruz Nieto D, Huamán Carranza M. Efecto de dosis de vinaza en el cultivo de rabanito (*Raphanus sativus L.*), para el control ecológico del gusano de tierra. *AS* 2020;13(2):274-85. DOI: <https://doi.org/10.32911/as.2020.v13.n2.740>
16. Viteri Vizuete ED. Evaluación de la vinaza de caña como abono orgánico y su posible efecto tóxico en el cultivo de rábano (*Raphanus sativus*) [tesis licenciatura]. [Quito]: Universidad Central del Ecuador; 2015 [citado 26 de octubre de 2021]. Recuperado a partir de: <http://www.dspace.uce.edu.ec/handle/25000/6428>
17. Bazán Tapia R. Manual de procedimientos de los análisis de suelos y agua con fines de riego [Internet]. Lima: Instituto Nacional de Innovación Agraria; 2017 [citado 12 de octubre de 2021]. 92 p. Recuperado a partir de: [https://repositorio.inia.gob.pe/bitstream/20.500.12955/504/1/Bazan\\_Manual\\_de\\_procedimientos\\_de\\_los.pdf](https://repositorio.inia.gob.pe/bitstream/20.500.12955/504/1/Bazan_Manual_de_procedimientos_de_los.pdf)
18. Instituto Nacional de Innovación Agraria (INIA) – Huaral. Análisis básico de fertilidad de suelos. Hoja análisis de suelo del distrito de Barranca. Laboratorio de Suelos de Instituto Nacional de Innovación Agraria - Huaral. Perú. 2018
19. Universidad Nacional Agraria LA Molina (UNALM). El Laboratorio de Análisis Agua, Suelo y Medio Ambiente. Resultado de análisis de vinaza. Universidad Agraria La Molina. 2018, Lima-Perú.
20. Senatore D, Queirolo A, Wajswol S, Bajsa N. Monitoreo de la aplicación de vinaza como fertili-

- zante en caña de azúcar con indicadores microbianos de suelo. INNOTECH 2017;(13);92-7. DOI: <https://doi.org/10.26461/13.09>
21. Prasad MN. Recovery of resources from biowaste for pollution prevention. In: Prasad MNV, Shih K, editors. Environmental Materials and Waste. Ámsterdam: Elsevier B.V; 2016. p. 1-19. DOI: <https://doi.org/10.1016/B978-0-12-803837-6.00001-9>
22. Lugo P. Evaluación del uso de la vinaza (residuo industrial) para la obtención de ácido cítrico empleando *Aspergillus niger* [tesis maestría]. [Valencia]: Universidad de Carabobo; 2017 [citado 6 de octubre de 2021]. Recuperado a partir de: [http://mriuc.bc.uc.edu.ve/bitstream/handle/12345\\_6789/4386/plugo.pdf?sequence=1](http://mriuc.bc.uc.edu.ve/bitstream/handle/12345_6789/4386/plugo.pdf?sequence=1)
23. García A, Rojas CA. Posibilidades de uso de la vinaza en la agricultura de acuerdo con su modo de acción en los suelos. Técnica [Internet]. 2005;9 (17): 3-13. Recuperado a partir de: <https://xdoc.mx/documents/posibilidades-de-uso-de-la-vinaza-en-la-agricultura-de-acuerdo-con-5e1cd14799680>
24. Zérega ML, Echner E, Hernández T, Arrieche A, Franco N, López A. Efecto de la vinaza y la fertilización química sobre el suelo y tres variedades de caña de azúcar en Venezuela. En: VI Congreso Asociación de Técnicos Azucareros de Latinoamérica y el Caribe. Guayaquil, Ecuador; 2006. p. 249-61.
25. Reynolds TM. Chemistry of nonenzymic browning. I. The reaction between aldoses and amines. Adv Food Res 1963;12:1-52. DOI: [https://doi.org/10.1016/s0065-2628\(08\)60005-1](https://doi.org/10.1016/s0065-2628(08)60005-1)
26. Arcila Hernández A. Evaluación de la aplicación de vinaza en suelo y su posible uso agrícola en plantas de cilantro (*Coriandrum sativum*) [tesis licenciatura]. [Maracay]: Universidad de Carabobo; 2017. [citado 16 de octubre de 2021]. Recuperado a partir de: <http://www.riuc.bc.uc.edu.ve/handle/123456789/4583>
27. Fernández Pelicó NY. Respuesta del crecimiento y rendimiento (Kg/ha) del cultivo, *Zea mays* L. Poaceae “Maíz” con tres tipos de vinaza y su efecto en las propiedades químicas del suelo [tesis licenciatura]. [Mazatenango]: Universidad San Carlos Guatemala; 2016 [citado 16 de octubre de 2021]. Recuperado a partir de: <https://repositorio-siidca.csuca.org/Record/RepoUSAC5310>
28. Núñez-Vázquez MC, Delgado-Acosta C, López-Padrón I, Martínez-González L, Reyes-Guerrero Y, Pérez-Domínguez G, et al. Nuevo bioestimulante y su influencia en la producción del frijol común. Cultrop 2020;41(4).e08.
29. Pineda Ruiz E, Chico Lamas Y, Vidal Díaz ML, Becerra de Armas E, Acosta Hernández F, Fernández Denis I, et al. Uso alternativo de la vinaza en la fertilización de la caña de azúcar, efectos sobre el cultivo y el suelo. Cent Agríc 2015;42(1):31-6.
30. Tuesta Popolizio DA. Efecto de la aplicación de vinazas de la industria del tequila en el cultivo del maíz y en la asociación planta-hongos micorrízicos arbusculares (HMA) [tesis licenciatura]. [Guadalajara]: Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C.; 2017 [citado 26 de octubre de 2021]. Recuperado a partir de: <https://ciatej.repositorio.institucional.mx/jspui/bitstream/1023/382/1/Diego%20Tuesta%20Popolizio.pdf>
31. Inga Sotelo MA. Estudio de compostaje del efluente vinaza de la actividad industrial azucarera para obtener mayor rendimiento ecológico en el cultivo de rabanito (*Raphanus Sativus* L) [tesis doctoral]. [Huacho]: Universidad Nacional José Faustino Sánchez Carrión; 2020 [citado 16 de octubre de 2021]. Recuperado a partir de: <http://repositorio.unjfsc.edu.pe/handle/UNJFSC/4212>

- 32.Gálvez Torres E, Legua Cárdenas J, Cruz Nieto D, Caro Soto F, Inga Sotelo M. Evaluación de Abono Orgánico de Vinaza y Bagazo de la Caña de Azúcar para la producción ecológica de rabanito (*Raphanus sativus* L.). AS 2019;12(2):236-49.  
DOI: <https://doi.org/10.32911/as.2019.v12.n2.645>
- 34.Korndörfer GH, Nolla A, Gama AJM. Manejo, aplicación y valor fertilizante de la vinaza para caña de azúcar y otros cultivos. Tecnicaña [Internet]. 2010 [citado 5 de octubre de 2021]; (24): 23-28. Recuperado a partir de: [https://issuu.com/relistatecnicana/docs/tec\\_no24\\_2010](https://issuu.com/relistatecnicana/docs/tec_no24_2010)

**Editor's Note:**

*Journal of the Selva Andina Biosphere (JSAB) remains neutral with respect to jurisdictional claims published on maps and institutional affiliations.*

- 33.Jaramillo Chamba RA, Novoa V (dir). Efecto de la vinaza, en el rendimiento de una mezcla forrajera establecida en un Andisol [tesis licenciatura]. [Quito]: Escuela Politécnica Nacional; 2010 [citado 10 de octubre de 2021]. Recuperado a partir de: <https://bibdigital.epn.edu.ec/bitstream/15000/1688/1/CD-2651.pdf>