

Physical and chemical properties of soils in Gera moist Afromontane forest, southwest Ethiopia

Propiedades físicas y químicas de los suelos en el bosque húmedo afromontano de Gera, suroeste de Etiopía

Egeta Dereje^{*ID}, Yadesa Giche^{ID}, Eshete Abeje^{ID}, Belay Hailu Belay^{ID}

Article Data

Ethiopian Forestry Development Jimma Center.
Research department: Ecosystem management re-
searcher.
Street: Addis Ababa City, Ethiopia.
Avenue: Jimma Town, Ethiopia.

***Contact address:**

Dereje Egeta.

Ethiopian Forestry Development Jimma Center.
Research department: Ecosystem management re-
searcher.
Street: Addis Ababa City, Ethiopia.
Avenue: Jimma Town, Ethiopia.
E-mail: egetaidereje@gmail.com

Keywords:

Bulk density,
cation exchangeable capacity,
correlation,
degraded forest,
organic carbon,
soil composite,
soil fertility,
soil texture.

J. Selva Andina Biosph.
2023; 11(2):118-130.

Article ID: 136/JSAB/2023

Article history

Received June 2023.
Returned in August 2023.
Accepted September 2023.
Available online November 2023.

Edited by:
Selva Andina
Research Society

Palabras clave:

Densidad masiva,
capacidad intercambiable de cationes,
correlación,
bosque degradado,
carbón orgánico,
compuesto de suelo,
fertilidad del suelo,
textura de la tierra.

Abstract

Soil is the centre of plant growth and, in turn, plants are important elements in soil creation. This points to a strong positive correlation between forest growth dynamics and soil qualities. However, the decline in the biological, chemical and physical characteristics of forest soils alters this link. Therefore, the objective of the study was to determine the physical and chemical properties of soils in the Afromontane rainforest of Gera as inputs for vegetation and land management. A systematic random sampling design was used to collect a soil sample. Twenty main plots with a size of 20 m by 20 m pits measuring (0.5) m by (0.5) m, one in the centre and four in the corners, were dug for the collection of soil samples. Soil samples were collected to a depth of 0 - 20 cm using an auger because the soil has a similar profile and is mixed to form a soil composite. From the composite sample, twenty samples were randomly selected as a working sample. Descriptive statistics and inferential statistics were carried out to analyze the data using SAS software. The sampled soil contains an average of 56.93 % sand, 26.22 % silt and 16.36 % clay. The average bulk density of the forest is 1.49 g cm⁻³. The forest contains 3.91 and 6.74 % organic carbon and matter, respectively. The analysis revealed that sandy loam accounts for 70 % of the texture classes in the Gera forest. Total nitrogen was highly correlated with organic carbon and matter, and magnesium and calcium were highly correlated with cation exchange capacity, while sodium was highly correlated with electrical conductivity. Further study on other soil physico-chemical properties will be required for rapid rehabilitation of the Gera forest for site species that match and sustain the forest.

2023. *Journal of the Selva Andina Biosphere*[®]. Bolivia. All rights reserved.

Resumen

El suelo es el centro de crecimiento de las plantas y, a su vez, las plantas son elementos importantes para la creación del suelo. Esto señala una fuerte correlación positiva entre la dinámica del crecimiento forestal y las cualidades del suelo. Sin embargo, la disminución de las características biológicas, químicas y físicas de los suelos forestales altera este vínculo. Por lo tanto, el objetivo del estudio fue determinar las propiedades físicas y químicas de los suelos del bosque húmedo afromontano de Gera como insumos para la vegetación y el manejo del territorio. Se utilizó un diseño sistemático de muestreo aleatorio para recolectar una muestra de suelo. Se cavaron veinte parcelas principales con un tamaño de 20 m por 20 m pozos que miden (0.5) m por (0.5) m, una en el centro y cuatro en las esquinas, para la recolección de muestras de suelo. Las muestras de suelo se recolectaron hasta una profundidad de 0 - 20 cm usando un sinfín debido a que el suelo tiene un perfil similar y se mezcla para formar un compuesto de suelo. De la muestra compuesta, se seleccionaron al azar veinte muestras como muestra de trabajo. Se llevaron a cabo estadísticas descriptivas y estadísticas inferenciales para analizar los datos utilizando el software SAS. El suelo muestreado contiene una media de 56.93 % de arena, 26.22 % de limo y 16.36 % de arcilla. La densidad aparente promedio del bosque es de 1.49 g cm⁻³. El bosque contiene 3.91 y 6.74 % de carbono orgánico y materia, respectivamente. El análisis reveló que el franco arenoso representa el 70 % de las clases de textura en el bosque de Gera. El nitrógeno total estaba muy correlacionado con el carbono orgánico y la materia, y el magnesio y el calcio estaban altamente correlacionados con la capacidad de intercambio catiónico, mientras que el sodio estaba altamente correlacionado con la conductividad eléctrica. Se requerirá más estudio sobre otras propiedades fisicoquímicas del suelo para la rehabilitación rápida del bosque de Gera para especies del sitio que coincidan y sostengan el bosque.

2023. *Journal of the Selva Andina Biosphere*[®]. Bolivia. Todos los derechos reservados.



Introduction

Soil is one component of natural resources that is made up of mineral matter, water, air, and organic matter (OM)¹. Soil is the growth center for plants due to the give support, nutrients, and water required for plant development, and in turn, plants are important elements for soil creation through geological processes and the chemistry of the above-ground atmosphere².

This indicates a strong positive correlation between forest growth dynamics and soil qualities³. However, the decline in the biological, chemical and physical characteristics of forest soils disrupts this link and reduces their ability to operate fully, whether temporarily or permanently⁴.

In the forest ecosystem, deforestation, forest degradation, population growth and agricultural expansion are major causes of soil degradation that decrease soil fertility and alter ecology⁵. Therefore, rehabilitation of degraded forests should remain one aspect of the interventions aimed at combating the effects of soil fertility loss, global warming, and climate change. Rehabilitation of degraded forests demands an understanding of both the physical and chemical properties of the soil. Thus, the understanding of soil physicochemical properties provides insight into the environmental requirements of the plant species needed for successful ecological restoration and biodiversity protection⁶ specifically, the natural distribution of forest tree species, growth, and forest biomass is governed by soil physical properties such as soil texture, bulk density (BD), and porosity. Besides soil, physical properties also soil chemical properties such as soil: pH, OM, total nitrogen (TN), available phosphorus (AVP), and exchangeable bases, also affect plant growth and distribution⁷.

Most studies on soil properties have focused on the sustainability of agricultural production, but many

have not focused on the soil properties of forest production⁸. However, forest soils are the fundamental agents for maintaining tree growth and diversity⁹. The physical, chemical and biological properties of forest soil are vital for improving productivity and maintaining tree growth and diversity. In addition, its physicochemical properties are essential for sustainable management of natural forests¹⁰.

In southwestern Ethiopia, waterlogging is one of the main problems in the region, which in turn can cause leaching of nutrients under plant roots, making the soil deficient in some nutrients, and affecting the acceleration of the restoration of degraded forests proposed by the Ethiopian government as a green legacy¹¹. This large-scale afforestation of degraded lands and forests is practiced in southwestern Ethiopia, particularly in the degraded Gera forest, for climate and erosion control, carbon sequestration, food security enhancement and biodiversity conservation¹². However, the success stories were below expectations and the problem remains immense. This could be related to the lack of species and site matching¹³. In general, no research has been conducted on soil properties in the degraded humid Afromontane rainforest (AF) of Gera. Therefore, knowledge of soil physical and chemical properties is crucial and a prerequisite for restoration of degraded forests and sustainability of natural vegetation.

The objective of the study was to determine the physical and chemical properties of soils in the wet Gera PA as inputs for vegetation and soil management.


Materials and methods

Study sites selection. Following a full reconnaissance survey to select the moist AF of Southwest Ethiopia, Gera moist AF was purposefully selected for this

study based on its level of degradation based on FAO¹⁴ forest degradation assessment guidelines.

Description of the study area. The research was done in Gera District, Jimma Zone, Oromia Regional State, in Southwest Ethiopia. The district is located at an elevation of 1300 to 2400 meters above sea level, within latitudes of 7° 27' to 7° 55' north, and a longitude of 38° 01' to 36° 24' east. The mean annual temperature of the district ranged between 15 and 22° C, and the vast area of the district's annual rainfall varies between 1300 and 1700 mm¹⁵. The agro ecological zones of the district are 3.7, 50.19, and 46.11 % lowland, mid-highland, and highland, respectively. The forest is located about 440 km southwest of Addis Ababa, the capital city of Ethiopia¹⁶.


Figure 1 map of study area



Soil sampling. Initially, a reconnaissance survey was conducted through a field visit and physical observation across the Gera forest to gain a general impression of the forest. Using this as a baseline distance between the transect lines was determined to be 430 m and the distance between consecutive quadrants was 500 m¹⁷. A systematic random sampling design was used to collect soil samples. Sample plots of 20 x 20 m were laid out along four transect lines following¹⁸. In each main plot, pits measuring 0.5 x 0.5 m were dug, one at the center and four at the corners, a total of one hundreds soil samples were collected from twenty plots before composite¹⁹. Soil samples were collected up to a depth of 0-20 cm since the soil

has a similar profile it was mixed to form a soil composite²⁰. This composite sample was divided into different parts, and twenty samples were randomly selected as working samples and transported to Jimma University, College of Agriculture and Veterinary Medicine soil laboratory located in Jimma town, Ethiopia, for soil physical and chemical analysis.

Figure 2 Sampling design for data collected



Furthermore, some selected properties of the soil, such as soil pH, exchangeable cations, electrical conductivity (EC), moisture contents (MC), AVP, TN, OM, exchangeable bases (K, Na, Mg, and Ca), cation exchange capacity (CEC), BD, and soil texture, were analyzed following the standard soil analysis methodology employed by the National Soil Laboratory of Ethiopia^{21,22}. The OMC of the soil was determined by the wet combustion procedure of Walkley-Black²³. The TN content of the soil was determined by the wet-oxidation procedure of the Kjeldahl method²⁴. The AVP content of the soil was determined by the P-Olsen method²⁵. The exchangeable cation content and the CEC of the soils were determined by the 1 M ammonium acetate (pH 7) method according to the percolation tube procedure²⁶. The effective CEC was calculated as the sum of the exchangeable cations extracted by ammonium acetate buffered at pH 7 plus the extractable Al extracted by 1 M KCl. BD was determined by the core method²⁷. The pH and exchangeable cation were calculated using a 1:2.5 soil-to-water suspension ratio. Soil texture

was determined using the hydrometer method²⁸.

Data analysis. Descriptive statistics (i.e., percentage, mean, maximum, and minimum) and inferential statistics (like correlation analysis) were carried out to analyze the data using SAS software. Furthermore, a Pearson correlation analysis was carried out between the soil physical and chemical properties.

Results

Physical properties of the soil. Forest soil showed that sand had the highest percentage value, ranging from 24 to 64.8 %. The silt content ranged from 20 to 55 %, while the clay content ranged from 9.6 to 28 % (Table 1).

Table 1 Values of determined physical properties of the studied soils

No of samples	Sand (%)	Silt (%)	Clay (%)	Textural Class	BD (g cm ⁻³)	Porosity (%)	MC (%)
1	64	21.6	14.4	Sandy loam	.97	63.3	71.20
2	60	24	16	Sandy loam	1.34	49.43	53.03
3	59	27	14	Sandy loam	1.23	53.58	51.03
4	42	30	28	Loam	1.44	45.66	52.12
5	62.6	22.4	15	Sandy loam	1.57	40	45.77
6	58.8	31.6	9.6	Sandy loam	1.615	39.05	46.05
7	60	22	18	Sandy loam	1.66	37.35	50.15
8	58	23	19	Sandy loam	1.36	48.67	64.55
9	63	24	23	Sandy loam	1.32	50.18	51.15
10	54	20	16	Sandy clay loam	1.63	38.84	51.40
11	62.4	20.2	27	Sandy loam	1.57	40.75	44.01
12	24	55	11	Silt loam	1.68	36.6	49.39
13	62	27	11	Sandy loam	1.62	38.86	47.79
14	52	26	22	Sandy clay loam	1.77	33.20	43.26
15	62	27.4	10.6	Sandy loam	1.38	47.92	44.85
16	64.8	23.6	11.6	Sandy loam	1.42	46.41	45.50
17	59	23.6	17.4	Sandy loam	1.60	39.62	39.17
18	53	26	21.5	Sandy loam	1.27	52.07	68.52
19	62	26	12	Sandy loam	1.86	29.80	34.69
20	56	24	20	Sandy loam	1.49	56.22	43.68
Mean	56.93	26.22	16.36	Sandy clay loam	1.49	44.38	49.87

MC moisture contents, BD bulk density.

Soil texture or the proportion of sand, silt, and clay-sized particles that make up the mineral fraction of the soil contains different ratios. Texturally, the studied soils were sandy loams with a sandy content ranging from 53 to 64.8 %, a silt content ranging from 31.6 to 21.6 %, and a clay content ranging from 14.4 to 27 %. Loam is contents 42 % sand, 30 % silt, and 28 % clay. The other textural class was sandy clay loam, which contained sand from 52 to 56 %, silt from 20 to 26 %, and clay from 16 to 22 %. Finally, silt loam contained 24 % sand, 55 % silt, and 11 % clay. The values of BD in the study area ranged between .97 g cm⁻³ and 1.86 g cm⁻³. The values of soil porosity in the study area oscillated between 29.8 and 63.3 %. Finally, the MC of the studied soil ranged

between 34.69 and 71.2 % (Table 1).

Chemical Properties of the Soil. The chemical properties of the studied soil were given in Table 2. The pH of the studied soils ranged from 4.00 to 5.75. The soil can be said to be slightly weak acidic based on the logarithmic pH scale. The exchangeable cation of the studied soils was found between 59.05 and 137.1. The organic carbon (OC) of the studied soil ranged between 4.68 and 5.03 %. TN content in the soils ranged from (.4) to (.43) % while the values of AVP in the soil were between 10.54 and 22.2 ppm. OM of the studied soil ranged between 8.06 and 8.67 %. CEC of the studied soil ranged between 23.32 and 27.38 meq 100 g⁻¹.

Table 2 Values of determined chemical properties of the studied soils

No of samples	pH	EC (ms cm ⁻¹)	OC %	TN %	AVP (ppm)	OM %	CEC (meq 100 g ⁻¹)
1	4.97	77.51	4.68	.40	10.54	8.06	23.32
2	4.80	93.73	3.12	.27	14.11	5.37	18.74
3	5.07	92.83	3.90	.33	14.59	6.72	20.44
4	4.53	59.05	3.23	.27	22.20	5.58	25.16
5	5.20	74.40	3.63	.31	10.76	6.25	20.00
6	5.36	77.54	4.056	.35	11.17	6.99	18.78
7	5.36	62.70	3.93	.34	11.18	6.79	22.58
8	4.00	71.54	5.03	.43	13.61	8.67	26.84
9	4.85	130.13	4.10	.35	12.17	7.05	20.60
10	5.29	86.36	4.17	.35	19.35	7.19	27.38
11	4.65	99.81	3.74	.32	21.57	6.45	21.40
12	5.27	65.83	4.29	.37	11.23	7.39	20.28
13	4.25	61.44	3.59	.32	12.90	6.18	17.86
14	4.89	60.39	4.21	.36	19.33	7.26	25.52
15	5.48	88.71	4.87	.42	11.67	8.40	22.10
16	4.77	74.08	3.02	.26	13.71	5.21	16.22
17	5.75	137.1	4.02	.34	13.46	6.92	22.54
18	5.52	93.09	3.20	.27	11.42	5.51	21.52
19	5.3	60.30	3.51	.30	11.08	6.05	18.10
20	5.63	107.89	3.96	.34	11.84	6.83	23.66

EC electrical conductivity, OC organic carbon, TN total nitrogen, AVP available phosphorus, OM organic matter, CEC cation exchange capacity.

Table 3 Values of determined exchangeable cations of the studied soils

No of samples	K (meq 100 g ⁻¹)	Mg (meq 100 g ⁻¹)	Ca (meq 100 g ⁻¹)	Na (meq 100 g ⁻¹)
1	8.43	2.41	18.18	.02
2	11.29	1.94	14.61	.02
3	11.67	2.11	15.93	.02
4	18.26	2.60	19.61	.01
5	8.60	2.07	15.59	.02
6	8.94	1.94	14.64	.02
7	8.94	2.34	17.60	.01
8	10.89	2.78	20.92	.01
9	9.74	2.13	16.06	.03
10	15.57	2.83	21.34	.02
11	17.26	2.21	16.68	.02
12	8.98	2.10	15.81	.01
13	10.32	1.85	13.92	.01
14	15.46	2.64	19.89	.01
15	9.33	2.29	17.23	.02
16	10.97	1.68	12.64	.01
17	10.77	2.33	17.57	.03
18	9.13	2.23	16.77	.02
19	8.86	1.87	14.11	.01
20	9.47	2.45	18.44	.02

K potassium, Mg magnesium, Ca calcium, and Na sodium.

Exchangeable Cations (K, Mg, Ca and Na). The values of exchangeable cations in the studied soils were K (8.43-18.26 meq 100 g⁻¹), Mg (1.68-2.83 meq 100

g⁻¹), Ca (18.18-21.34 meq 100 g⁻¹), and Na (.01-.03 meq 100 g⁻¹) (Table 3).

Correlation between soil physicochemical proper-

ties. The result of the Pearson's correlation analysis between soil properties of the study sites, Table 5.

Table 4 Descriptive statistical values of chemical properties and exchangeable cations of the studied soils

Parameters	Minimum	Maximum	Mean	Standard Deviation
pH	4	5.75	5.05	.46
OC (%)	3.02	5.03	3.91	.56
OMC (%)	5.21	8.67	6.74	.96
TN (%)	.26	.43	.33	.05
AVP (%)	10.54	22.2	13.90	3.69
Na (meq 100 g ⁻¹)	.011	.03	.02	.004
K (meq 100 g ⁻¹)	8.43	18.26	11.14	3.02
Ca (meq 100 g ⁻¹)	12.64	21.34	16.88	2.37
Mg (meq 100 g ⁻¹)	1.68	2.83	2.24	.32
CEC (meq100 g ⁻¹)	16.22	27.38	21.65	3.05
EC (ms cm ⁻¹)	59.05	137.1	83.72	22.38

EC electrical conductivity, OC organic carbon, TN total nitrogen, AVP available phosphorus, OMC organic matter content, CEC cation exchange capacity.

Table 5 Pearson's correlation coefficients between soil physicochemical properties

Parameters	pH	EC	OC	Sand	Clay	Silt	TN	AVP	OMC	CEC	K	Mg	Ca	Na	MC	BD	Porosity
pH	1																
EC	.36	1															
OC	.14	.06	1														
Sand	-.21	.19	-.18	1													
Clay	-.02	.01	-.06	-.50*	1												
Silt	.09	-.26	-.05	-.42	-.41	1											
TN	.14	.06	1.00*	-.17	-.08	-.03	1										
AVP	-.48*	.08	-.17	-.29	.47*	-.06	-.17	1									
OMC	.14	.06	1.00*	-.18	-.06	-.05	1.00*	-.17	1								
CEC	.11	.03	.56*	-.49	.75*	-.32	.55*	.29	.56*	1							
K	-.48*	.09	-.16	-.29	.46*	-.05	-.17	1.00*	-.16	.28	1						
Mg	.10	.03	.55*	-.49	.76*	-.32	.54*	.30	.55*	1.00*	.29	1					
Ca	.11	.03	.56*	-.49	.75*	-.32	.55*	.29	.56*	1.00*	.28	1.00*	1				
Na	.35	1.00*	.09	.19	.03	-.29	1.00	.08	.09	.07	.08	.07	.07	1			
MC	-.32	.00	.05	-.12	.23	-.11	.05	.03	.05	.23	.03	.23	.23	.03	1		
BD	.18	-.49*	.05	-.29	-.01	.09	.04	-.06	.05	-.04	-.06	-.05	-.04	-.51*	-.60*	1	
Porosity	-.05	.57*	-.03	.23	.08	-.12	-.01	.02	-.03	.11	.02	.12	.11	.59*	.48*	-.94*	1

Correlation is significant at $P \leq .05$, Values with * are values for significant pair.

EC electrical conductivity, OC organic carbon, TN total nitrogen, AVP available phosphorus, OMC organic matter content, CEC cation exchange capacity, MC moisture contents, BD bulk density.

Exchangeable cation has a low correlation with pH (.36). Clay was moderately negative and significantly correlated with sand (.50). Silt has a low negative correlation with an exchangeable cation of (.26) and sand has a correlation of (.42) and clay has a correlation of (.41). TN is highly and significantly correlated (1.00) with OC. AVP is inversely related to pH (.48) and significantly related to clay (.47). OM is highly correlated with OC and TN (1.00). Cata-

strophic exchange capacity is highly and significantly correlated with clay (.75) moderately with OC (.56) TN (.55) and OM (.56) and negatively with sand (.49) and silt (.32). Potassium (K) was negatively correlated with pH (.48) clay (.46) and very highly correlated with AVP (1.00), as well as negatively correlated with sand (.29) and positively correlated with CEC (.28). Magnesium (Mg) was highly correlated with CEC (1.00) and clay (.76) moderately cor-

related with OC (.55) TN (.54) and OM (.55) negatively correlated with sand (.49) and silt (.32) and positively correlated with AVP (.30) and potassium (.29). Calcium is highly correlated with magnesium (1.00) and CEC (1.00), is moderately correlated with OC (.56) TN (.55) and OMC, MO (.56) is negatively correlated with sand (.49) and silt (.32) and is positively correlated with AVP (.29) and potassium (.28). Sodium has a high correlation with higher exchangeable cation and TN (1.00), a positive correlation with lower pH (.35) and a negative correlation with lower silt (.29). MC is negatively correlated with (.32) and positively correlated with clay (.23) CEC (.23) magnesium, and calcium. Table 5 shows that BD was negatively correlated with EC (.49), moderately correlated with sodium (.51) and MC (.60) and negatively correlated with sand (.29). Porosity is moderately positively correlated with EC (.57) sodium (.59) and MC (.48) but highly negatively correlated with BD (.94).

Discussion

Soil is the most important natural resource because it provides the basis for all other activities required for life. So its protection is a prerequisite for nature conservation. Soil physicochemical properties are one technique for determining the state of the soil and recovering it. Our research plays a great role in minimizing the cost required for the restoration of degraded forests in southwest Ethiopia due to the physical and chemical properties of the soil is known. As a result, it keeps species and sites from being mismatched. Additionally, it is crucial for various stakeholders involved in the development of forestry in southwest Ethiopia.

When we consider particle size, the soils studied have the highest mean percentage of sand and the lowest of clay, with silt in between (Table 1). Our results agree with the findings of²⁹ and³⁰, who re-

ported that sand has the highest particle size distribution in the areas studied, and agree with those of²¹, who indicated that sand has the highest particle size and clay the lowest in the native forest areas studied. This similarity could be due to the effects of the depth at which the sample was collected. In addition, our results agree with those of³¹, who reviewed different literature on different forest types in Kerala, noted that sand has the largest particle size, silt is medium and clay is the smallest. Our findings contradict those of³², which reported that sand particle size is the lowest and clay the highest in natural upland forests of northwestern Ethiopia. This difference is due to the fact that particle size distribution varies over an extended period as a consequence of erosion, deposition, eluviation, weathering and parent materials. In particular, temperature and precipitation are the two main active soil-forming factors contributing to soil weathering and development. Our observation also contradicts the findings of^{33,34} which pointed out that sand has the lowest particle size at the studied site. This difference could be due to the effects of soil depth.

Soil porosity has a direct relationship to the grain size distribution of the sand (Table 1). This general truth means that sandy soil has no water-holding capacity, and good soil water-holding capacity is an important factor in soil fertility. Clay has the lowest porosity, which means that it has the highest water holding capacity. Our results agree with those of³⁴, who reported that the total soil porosity of the studied site was recorded 63 %, which is in our range. This similarity is due to the dominance of clayey soil of the studied site. Our results contradict the findings of³⁵, which indicated that the porosity of natural forests was 74.14 %. This difference could be due to the impact of soil compaction that will damage soil pores and the level of disturbance.

The BD of the studied forest was recorded at an average of 1.49 g cm^{-3} (Table 1). Our observation

agrees with the findings of³³, who noted that the BD of Harena forest and Yayu forest were recorded as 1.42 and 1.23 g cm⁻³ respectively and contradicts his findings of Bonga forest and Berhane-Kontir forest, which were recorded as 1.01 and 1.07 g cm⁻³ respectively. This difference is due to the frequency of rainfall and seasonal climatic variations at the two sites. Our results contradict those of³⁶, which found that DB of three ecclesiastical uplands, mixed midland and lowland-midland forests were recorded as (.07), (.05) and (.21) g cm⁻³, respectively. This difference is due to differences in species composition, disturbance, decomposition rates, etc. Soil BD is influenced by factors like OMC, management practices, texture, soil fauna, etc. Our result also contradicts the findings of³⁴, which pointed out that the BD of natural forests of different land uses is recorded as (.98) g cm⁻³. This difference is due to soil depth and level of disturbance. Our results contradict the findings of²¹, which reported that the BD of native forests in the studied areas is (.78) g cm⁻³. Soil BC could be affected by the availability of parent materials.

The soil MC of our studied forest was recorded at the mean of 49.87 % (Table 1). Our results are in agreement with those of³⁵, who observed that the MC of the studied areas was recorded in the ranges of 71.83 and 43.38 %. This similarity could be due to the fact that soils with a very high proportion of sand have a very low water holding capacity due to the large pore spaces between the particles, which allow water to move freely into the deeper layers, leaving the upper layers practically dry. Our result contradicts the findings of⁵, who reported that the soil MC of the studied site was recorded at 26.4 and 15.0, respectively. This difference could be due to rainfall and other climatic conditions in the two areas. Our observation also contradicts the findings of³³, which observed lower soil MC in Harena forest, Bonga forest, Berhane-Kontir forest and Yayu forest. This difference could

be due to variations in the proportions of soil texture, vegetation cover, climatic gradients and the time of data collection.

The TN of our studied forest ranges from (.26) to (.43) or an average of (.33) % (Table 2). In northeastern Ethiopia⁵, the NT content of Gatira George church forest and Gemeshat natural forest were reported to be (.42) and (.36) respectively. This similarity could be due to the soil having good aeration, drainage and permeability. NT is higher due to the low level of soil disturbance by erosion. Our findings contradict those of³⁴, which pointed out that the TN of the studied site was recorded as (.59), which is far from our results. This difference could be due to the reduction of aboveground biomass due to different impacts of human and livestock activities and levels of forest degradation. Our results contradict the findings of³⁷, which refers that the studied NT is recorded at (.09) to (.12) % which is lower than our findings. This difference is due to the fact that TN is affected by storage time, temperature and MC of the areas.

The result of the analysis implies that the average pH of the studied soil sample was pH 5.05 (Table 4). According to the soil pH classification³⁸, the sample has a neutral soil reaction class (within the range of 6.7-7.3) and an acid soil reaction class if it is below 6.7. Therefore, it needs to be upgraded to neutral. Our results indicate that the soil sampled was acidic. The acidic soil may be due to acids released by the decomposition of organic residues obtained from forest vegetation. Our result was similar to the findings of³⁹, which noted that the pH of the Jibat forest soil was 5.66. This similarity is due to the characteristics of the vegetation, the nature of the parent material and rainfall. Our observation is in agreement with the findings of⁴⁰, who noted that the pH of the studied site is slightly acidic. Our observation is in agreement with the findings of³¹, who reviewed different forest

types in Kerala and recorded the pH value of the evergreen forest as 5.1, or acidic. This is due to high rainfall in humid tropical areas, which causes leaching of base ions from soils. Soil pH values can be affected by the quality of the litter falling from the plants.

The mean exchangeable cations of the soil sampled as a whole was 83.72 ms cm^{-1} ($.83 \text{ ds m}^{-1}$). The EC measurements are used as an indication of the total amounts of soluble salts in the soils. According to⁴¹, the soil is classified as non-saline ($<2 \text{ ds m}^{-1}$). Therefore, this sampled soil was a non-saline soil.

The mean OMC of the sampled soil was 6.74 (Table 4). This means that in terms of OMC the sampled soil was high, which means that the soil is fertile as OMC is a good indicator of soil fertility. This could be due to the fact that OMC was added to the soils through litter fall and fine root decomposition. The OMC value of the soil is low (below the range of 1-2 %)⁴², below the range of (.8) - 2.6 %⁴³, and below the range of (.86) - 2.59 %³⁸. Our findings contradict those of²⁰, which found that the OMC of the studied soil ranged from 1.68 to 2.60 %. This difference could be due to microbial activities and other biotic and abiotic factors. Our results also contradict the findings of⁴⁴ who reported that the OMC content of the studied site ranged from 2.14 to 4.07 %. This difference could be due to the nature of the plant species, rooting system, plant height, canopy, growth level, soil formation processes and season of the year. Our observation contradicts the findings of⁵, who pointed out that the OMC of the studied forests accounted for half of our findings. This difference could be due to the high removal of fallen litter means humus or topsoil in the forests studied by⁵ and the diversity of vegetation cover in our case. The difference could be due to being influenced by time storage, temperature and MC. The OC of the studied forest ranged from 3.02 to 5.03 (Table 2). This observation contradicts the findings

of⁴⁴ which indicated that the OC of the studied areas ranged from 1.24 to 2.36 %. This difference is due to the level of forest disturbance, such as grazing, which affects the aboveground biomass of the forest. The difference in soil OC accumulation is attributed to vegetation type, soil type, human disturbance through vegetation removal, overgrazing by livestock, parent material and other climatic factors. OC is affected by the depth of soil where data are collected. The OC in our study forest was twice that of the Gatira George church forest and Gemeshat natural forest in northeastern Ethiopia, according to the study of⁵. This difference could be due to soil layers and locations due to climatic, edaphic, biological and land management practices, vegetation cover, aspect and topography of the studied areas.

The AVP of the studied forest was observed in the ranges of 10.54 to 22.3 ppm respectively (Table 2). These results agree to some extent with the findings of²⁰, who reported that the total AVP of the studied site was recorded at 10.92 and 18.22 % respectively. The high AVP values can be attributed to the high OMC content in the soil.

CEC, is a measure of the total amount of negative charges within the soil that adsorb plant nutrient cations such as calcium, magnesium, potassium, and Na. The cation exchange capacity of the studied forest was dominated by potassium and calcium (Table 3). These findings contradict the result obtained by³⁵, which observed that calcium was the lowest cation exchange capacity for the natural forests studied. This difference could be due to the rate of loss of natural forests. Forest loss affects soil PMC. Such a reduction in the supply of PMC will, of course, cause a reduction in soil nutrient contents such as N, P, Ca, Mg and K. The cation exchange capacity of our study was observed in the range of 16.22 to 27.38 meq 100 g^{-1} . Our result is in line with the findings of³⁵, who pointed out that the exchange capacity cations of the

studied natural forest is 26.10 meq 100 g⁻¹. This similarity is due to the vegetation composition of the studied forest.

Generally, the analysis revealed that sandy loam accounts for 70 % of the textural classes in the Gera evergreen forest. Soil porosity is inversely proportional to BD. TN was very highly correlated with OC and OMC, magnesium and calcium were highly correlated with CEC, whereas sodium was highly correlated with exchangeable cation. Our research provides the basis for the restoration of degraded forests and adjacent land use in south-west Ethiopia by giving an overview of the soil properties of the site to ensure the right species for the site (species-site matching). Also give highlighted information to policymakers for sustainable land management and to take measures for soil management practices to improve the soil quality of the area. Further study will be required on other soil physicochemical properties for fast rehabilitation of the Gera forest for site species matching and sustaining the existing forest. Additionally, awareness creation and job opportunities should be made available to the local community that lives around and within the forest area, which helps prevent forest degradation and deforestation that leads to soil fertility loss. For future rehabilitation of the degraded forest of Gera, one should consider the soil properties of the forest to plan plant species agreements with the soil.

Source of financing

This research received no external funding

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We want to say thank you to Jimma University College of Agriculture and Veterinary Medicine for research equipment or field material support and soil laboratory.

Ethical considerations

I hereby declare that this paper is not being submitted to other journal publishers for potential publication. Furthermore, I affirmed that there has been no scientific wrongdoing in this outcome.

Research limitations

Micronutrients (trace elements such as iron, zinc, copper and manganese were not included in the present research due to budget scarcity, soil laboratories, and field equipment for soil sample collection like augers and core samplers under the Jimma Forestry Development Center.

Authors' contribution

Dereje Egeta, analyzed and interpreted data and full write-up. *Giche Yadesa*, data collection, data encoding, and editing. *Abeje Eshete*, research design and editing. *Hailu Belay*, data collection and editing.

Cited Literature

1. Rajput A, Yadav SS. Physical and chemical properties of soil: indicator of forest health and ecosystem of math wad range. *Int J Life Sci Res* 2019;7 (3):93-9.
2. Neupane A, Raya SB, Bhattarai S, Neupane S. Organic matter and properties of soil in forest and agriculture areas. *Int J Geogr Geol Environ* 2019;1 (2):28-33.

- <https://doi.org/10.22271/27067483.2019.v1.i2a.107>
3. Soong JL, Janssens IA, Grau O, Margalef O, Stahl C, Van Langenhove L, et al. Soil properties explain tree growth and mortality, but not biomass, across phosphorus-depleted tropical forests. *Sci Rep* 2020;10(1):2302. DOI: <https://doi.org/10.1038/s41598-020-58913-8>
 4. Agbeshie AA, Abugre S, Atta-Darkwa T, Awuah R. A review of the effects of forest fire on soil properties. *J For Res* 2022;33:1419-41. DOI: <https://doi.org/10.1007/s11676-022-01475-4>
 5. Ayalew A, Tesfaye A, Mulat Y. Effect of species diversity and forest structure on soil chemical properties of Gatira George's Church Forest and Gemeshat Natural Forest, in North East Ethiopia. *Agrotechnology* 2022;11(11):1000290. DOI: <https://doi.org/10.35248/2168-9881.22.11.290>
 6. Birhanu L, Bekele T, Tesfaw B, Demissew S. Soil seed bank composition and aboveground vegetation in dry Afromontane forest patches of North-western Ethiopia. *Trees For People* 2022;9:100292. DOI: <https://doi.org/10.1016/j.tfp.2022.100292>
 7. Jamil N, Sajjad N, Ashraf H, Masood Z, Bazai ZA, Khan R, et al. Physical and chemical properties of soil quality indicating forests productivity: a review. *American-Eurasian J Toxicol Sci* 2016;8(2):60-8. DOI: <https://doi.org/10.5829/idosi.aejts.2016.8.2.10312>
 8. Fentie SF, Jembere K, Fekadu E, Wasie D. Land use and land cover dynamics and properties of soils under different land uses in the Tejibara Watershed, Ethiopia. *Sci World J* 2020;2020:1479460. DOI: <https://doi.org/10.1155/2020/1479460>
 9. Zhao Y, Zhao M, Qi L, Zhao C, Zhang W, Zhang Y, et al. Coupled relationship between soil physico chemical properties and plant diversity in the process of vegetation restoration. *Forests* 2022;13:648. DOI: <https://doi.org/10.3390/f13050648>
 10. Eshetu Z. Forest soils of Ethiopian highlands: their characteristics in relation to site history. Studies based on stable isotopes [doctoral thesis]. [Umeå]: Swedish University of Agricultural Sciences; 2000 [cited 16 May 2023]. Retrieved from: <https://pub.epsilon.slu.se/id/document/20356428>
 11. Sori G, Iticha B, Takele C. Spatial prediction of soil acidity and nutrients for site-specific soil management in Bedele district, Southwestern Ethiopia. *Agric Food Secur* 2021;10:59. DOI: <https://doi.org/10.1186/s40066-021-00334-5>
 12. Lemenih M, Kassa H. Re-Greening Ethiopia: history, challenges and lessons. *Forests* 2014;5(8):1896-909. DOI: <https://doi.org/10.3390/f5081896>
 13. Mamo N, Sterba H. Site index functions for *Cupressus lusitanica* at Munessa Shashemene, Ethiopia. *For Ecol Manag* 2006;237(1-3):429-35. DOI: <https://doi.org/10.1016/j.foreco.2006.09.076>
 14. Organización de las Naciones Unidas para la Agricultura y la Alimentación. Base referencial mundial del recurso suelo. Un marco conceptual para la clasificación, correlación y comunicación internacional [Internet]. Roma: Organización de las Naciones Unidas para la Agricultura y la Alimentación; 2008 [citado 16 de mayo de 2023]. Informe sobre recursos mundiales de suelos No.: 103. Recuperado a partir de: <https://www.fao.org/3/a0510s.pdf>
 15. Diro S, Erko B, Fikirie K. Production and adoption constraints of improved coffee varieties in Jimma Zone, Southwest Ethiopia. *Agric Food Sci Res* 2019;6(1):41-9. DOI: <https://doi.org/10.20448/journal.512.2019.61.41.49>
 16. Arage A, Ketema M, Gemechu A. Factors affecting honey market outlet choices in Gera district, Jimma zone, Oromia region of Ethiopia. *Int J Innov Res Multidiscip Field* 2020;6(2):121-9. DOI: <https://doi.org/10.2015/IJRMF.2455.0620/202002023>

17. Esubalew E. Woody species diversity, regeneration and population structure along altitude gradient in Alemsaga Forest, South Gondar, North Western Ethiopia. *Biodivers Endanger Species* 2021;9(2):1-9.
18. Sewale B, Mammo S. Analysis of floristic composition and plant community types in Kenech Natural Forest, Kaffa Zone, Ethiopia. *Trees For People* 2022;7:100170. DOI: <https://doi.org/10.1016/j.tfp.2021.100170>
19. Castro M, Machado-Perez F, Raven R. Alpine wet meadows: soil properties' effect on plant diversity. *CEC Res* 2018;2(2):1-11. DOI: <https://doi.org/10.21973/N33957>
20. Opeyemi AO, Adewunmi BI, Oluwaseyi AI. Physical and chemical properties of soils in Gambari Forest Reserve Near Ibadan, South Western Nigeria. *J Biores Manag* 2020;7(2):57-67. DOI: <https://doi.org/10.35691/JBM.0202.0132>
21. Yimer F, Ledin S, Abdelkadir A. Concentrations of exchangeable bases and cation exchange capacity in soils of cropland, grazing and forest in the Bale Mountains, Ethiopia. *For Ecol Manag* 2008;(6):1298-302. DOI: <https://doi.org/10.1016/j.foreco.2008.06.047>
22. Mulugeta T, Melese A, Wondwosen T. Effects of land use types on selected soil physical and chemical properties: The case of Kuyu District, Ethiopia. *Eurasian J Soil Sci* 2019;8(2):94-109. DOI: <https://doi.org/10.18393/ejss.510744>
23. Walkley A, Black AI. An examination of the degtjareff method for determining soil organic matter, and the proposed modification of the chromic acid titration method. *Soil Sci* 1934;37:29-38. DOI: <https://doi.org/10.1097/00010694-193401000-00003>
24. Bremner JM, Mulvaney CS. Nitrogen total. In: Page AL, editor. *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*. Nueva Jersey: John Wiley & Sons, Inc.; 1983. p. 595-624. DOI: <https://doi.org/10.2134/agronmonogr9.2.2ed.c3>
25. Olsen SR, Sommers LE. Phosphorus. In: Page AL, editor. *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*. Nueva Jersey: John Wiley & Sons, Inc.; 1983. p. 403-30. DOI: <https://doi.org/10.2134/agronmonogr9.2.2ed.c2>
26. Van Reeuwijk LP. Procedures for soil analysis [Internet]. Netherlands: Food and Agriculture Organization of the United Nations; 2002 [cited 22 April 2023]. 119 p. Retrieved from: https://www.isric.org/sites/default/files/ISRIC_TechPap09.pdf
27. Blake GR, Hartge KH. Bulk density. In: Klute A, editor. *Methods of Soil Analysis: Part 1 Physical and Mineralogical Methods*. Nueva Jersey: John Wiley & Sons, Inc.; 1986. p. 363-75. DOI: <https://doi.org/10.2136/sssabookser5.1.2ed.c1>
28. Gee GW, Bauder JW. Particle size analysis. In: Klute A, editor. *Methods of Soil Analysis: Part 1 Physical and Mineralogical Methods*. Nueva Jersey: John Wiley & Sons, Inc.; 1986. p. 383-411. DOI: <https://doi.org/10.2136/sssabookser5.1.2ed.c1>
29. Chinevu NC, Unanaonwi EO, Amonum IJ. Physical and chemical characteristics of forest soil in southern Guinea Savanna of Nigeria. *Agric For Fish* 2013;2(6):229-34. DOI: <https://doi.org/10.11648/j.aff.20130206.15>
30. Seyoum B. Assessment of soil fertility status of Vertisols under selected three land uses in Girar Jarso District of North Shoa Zone, Oromia National Regional State, Ethiopia. *Environ Syst Res* 2016;5:18. DOI: <https://doi.org/10.1186/s40068-016-0069-y>
31. Vishnu PS, Sandeep S, Sujatha MP. Physico-chemical properties of forest soils in Kerala-a review. *J Environ Sci Toxicol Food Technol* 2017;11(1):23-6. DOI: <https://doi.org/10.9790/2402-1101032326>

32. Molla E, Getnet K, Mekonnen M. Land use change and its effect on selected soil properties in the northwest highlands of Ethiopia. *Heliyon* 2022;8(8):e10157. DOI: <https://doi.org/10.1016/j.heliyon.2022.e10157>
33. Kufa T. Characterizing major soil physical properties of coffee forest ecosystems in Ethiopia. *Int J Soil Sci Agron* 2013;1(1):1-9.
34. Abera W, Assen M. Dynamics of selected soil quality indicators in response to land use/cover and elevation variations in Wanka watershed, Northwestern Ethiopian Highlands. *Ekológia* 2019;38(2):126-39. DOI: <https://doi.org/10.2478/eko-2019-0010>
35. Wasis B. Soil properties in natural forest destruction and conversion to agricultural land, in Gunung Leuser National Park, North Sumatera Province. *J Man Hut Trop* 2012;18(3):206-12. DOI: <https://doi.org/10.7226/jtfm.18.3.206>
36. Kifle TE. Physico-chemical characteristics of the soils in three church forest of Central Ethiopia. *Res Ecol* 2022;2(4):1-8. DOI: <https://doi.org/10.30564/re.v2i4.2368>. Retraction in: Kifle TE. Physico-chemical characteristics of the soils in three church forest of Central Ethiopia 2022;2(4):1-8. DOI: <https://doi.org/10.30564/re.v2i4.2368>
37. Keeney DR, Nelson DW. Nitrogen-inorganic forms. In: Page AL, editor. *Methods of soil analysis: Part 2 Chemical and microbiological properties*. New York: John Wiley & Sons, Inc; 1983. p. 643-98.
38. Dubale P. Availability of phosphorus in the coffee soils of southwest Ethiopia. In: Mamo T, Haile M, editors. *Soil: the resource base for survival. Proceeding of the 2nd Conference of the Ethiopian Society of Soil Science: 23-24 September 1993*. Addis Ababa [Internet]. Ethiopia: Addis Ababa; 1996 [cited 22 April 2023]. p.119-29.
39. Tolessa T, Senbeta F. The extent of soil organic carbon and total nitrogen in forest fragments of the central highlands of Ethiopia. *J Ecol Environ* 2018;42:20. DOI: <https://doi.org/10.1186/s41610-018-0081-4>
40. Balagopalan M, Jose AI. Soil chemical characteristics in a natural forest and adjacent exotic plantations in Kerala, India. *J Trop For Sci* 1995;8(2):161-6.
41. Allison LE, Bernstein L, Bower CA, Brown JW, Fireman M, Hatcher JT, et al. *Diagnosis and Improvement of Saline Alkali Soils*. Richards LA, editor. *Agriculture Handbook No. 60* [Internet]. Washington: United States Department of Agriculture; 1954 [cited 16 May 2023]. 166 p. Retrieved from: https://www.ars.usda.gov/ARSUserFiles/20360500/hb60_pdf/hb60complete.pdf
42. Murphy HF. A report on fertility status and other data on some soils of Ethiopia. *Experimental Station Bulletin No. 44, Collage of Agriculture HSIU*. 1968. p. 551.
43. Debele B. The Physical criteria and their rating proposed for land evaluation in the highland region of Ethiopia. Land use planning and regulatory department. Addis Ababa, Ethiopia: Ministry of Agriculture; 1980.
44. Olorunfemi IE, Fasinmirin JT, Akinola FF. Soil physico-chemical properties and fertility status of long-term land use and cover changes: A case study in forest vegetative zone of Nigeria. *Eurasian J Soil Sci* 2018;7(2):133-50. DOI: <https://doi.org/10.18393/ejss.366168>

Editor's Note:
Journal of the Selva Andina Biosphere (JSAB). All statements expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, editors and reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.