

## Short communication. The effect of oestrus at slaughter on carcass and meat quality of gilts intended for dry-cured ham production

S. Calvo<sup>1</sup>, G. Ripoll<sup>1</sup>, J. A. Rodriguez-Sánchez<sup>1</sup> and M. A. Latorre<sup>2\*</sup>

<sup>1</sup> Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA).

Avda. Montaña, 930. 50059 Zaragoza. Spain

<sup>2</sup> Facultad de Veterinaria. Universidad de Zaragoza. Avda. Miguel Servet, 177. 50013 Zaragoza. Spain

### Abstract

On the day previous to slaughter, a total of 467 Duroc × (Landrace × Large White) gilts with approximately 130 kg body weight and 242 ± 3 days of age intended for Teruel ham production were observed to detect oestrus. The behavioural manifestations described in the literature were used for this. A total of 24 gilts were positively detected and individually identified (O gilts). From the same gilts observed, a total of 24 that clearly did not manifest oestrus were also individually identified (NO gilts). Therefore, a total of 48 gilts were used to study the effect of the presence of oestrus at slaughter on carcass and meat quality. No carcass characteristics were affected by oestrus at slaughter. However, all the carcasses (100%) from NO gilts were suitable for cured ham while 12.5% of carcasses from O gilts were not suitable because of the lack of the minimum backfat thickness required (16 mm) ( $p = 0.09$ ). The influence of oestrus at slaughter on meat characteristics was scarce. Under our experimental conditions, it can be concluded that avoiding the slaughter of gilts in oestrus might be recommendable when a minimum covering fat in carcass is needed to improve the quality of the end product. However, new trials would be needed to confirm this.

**Additional key words:** loin quality; pig females; stress.

### Resumen

**Comunicación corta. Efecto de la presencia de estro en el momento del sacrificio sobre la calidad de la canal y la carne de cerdas destinadas a la producción de jamón curado**

El día previo al sacrificio, se observaron un total de 467 cerdas Duroc × (Landrace × Large White), con aproximadamente 130 kg de peso vivo y 242 ± 3 días de edad destinadas a la producción de Jamón de Teruel, para detectar la presencia de estro mediante las manifestaciones descritas en la bibliografía. Un total de 24 cerdas fueron detectadas positivamente e identificadas individualmente (cerdas O). Entre las cerdas observadas, un total de 24 que claramente no manifestaban celo fueron también identificadas individualmente (cerdas NO). Por tanto, se usaron 48 cerdas para estudiar el efecto de la presencia de estro durante el sacrificio sobre la calidad de la canal y la carne de cerdas destinadas a Jamón de Teruel. Ninguna característica de la canal se vio afectada por la presencia de estro. Sin embargo, todas las canales (100%) de las cerdas NO fueron aptas para la producción de jamón curado, mientras que un 12,5% de las canales de cerdas O no lo fueron por no cumplir el mínimo espesor de grasa de cobertura requerido (16 mm) ( $p = 0,09$ ). El estro en el sacrificio apenas afectó a las características de la carne. Podemos concluir que podría ser recomendable evitar el sacrificio de cerdas en estro cuando se exige un mínimo de grasa de cobertura en la canal para mejorar la calidad del producto final. Sin embargo, se necesitarían más ensayos para confirmarlo.

**Palabras clave adicionales:** calidad del lomo; estrés; hembras porcinas.

Spain is the world leader in dry-cured ham and shoulder production with a total of 46 million pieces in 2008 (MARM, 2010). Currently, the only Protected

Designation of Origin (PDO) of dry-cured ham from crossbred commercial pigs (not autochthonous) accepted by the Spanish government is «Teruel ham». The

\* Corresponding author: malatorr@unizar.es  
Received: 01-03-10; Accepted: 07-09-10.

Abbreviations used: a\* (redness), b\* (yellowness), BW (body weight), C\* (chroma), DFD (dark, firm and dry), H\* (hue angle), L\* (lightness), m. GM (gluteus medius muscle), NO (not oestrus gilts), O (oestrus gilts), p (statistical significance level), PDO (Protected Designation of Origin), PSE (pale, soft and exudative), SE (standard error of the mean).

production of Teruel ham has increased drastically in recent decades from 2,000 pieces in 1985 to 700,000 in 2009 (CRPDO Jamón de Teruel, 2010). The Regulation of PDO Teruel ham establishes a minimum of 84 kg for carcass weight, of 16 mm for fat thickness over the *gluteus medius* muscle (m. GM) and of 11.3 kg for fresh ham to improve the uniformity and quality of the end product (BOA, 1993). In a recent study, Latorre *et al.* (2008) recommended an average slaughter weight of 130 kg to optimize the growth performance and carcass quality of these pigs. However, at this slaughter weight, all the carcasses from barrows were suitable for Teruel ham but a third of the carcasses from gilts were rejected because of the limited amount of covering fat depth (Latorre *et al.*, 2009a). There are several factors that can affect the production and quality of Teruel ham from gilts (weight/age at slaughter, health status, feeding or stress). Among causes of stress, the presence of oestrus at slaughter might be important. Traditionally, the slaughter of gilts in oestrus was avoided because it was believed that it negatively affected meat quality. Currently, the Regulation of PDO Teruel ham prohibits the slaughter of gilts in oestrus (BOA, 1993). Also, Iberian gilts which are slaughtered at 160 kg of body weight (BW) are usually castrated to avoid possible negative effects of oestrus on growth performance or meat quality (Serrano *et al.*, 2008). However, to our knowledge, there is no research study that justifies the effect of oestrus at slaughter on meat characteristics of gilts.

A total of 467 Duroc × (Landrace × Large White) gilts of approximately 130 kg BW and 242 ± 3 days of age intended for Teruel ham production were observed at farm during the previous day of the slaughter to detect oestrus. The behavioural manifestations described in the literature (Whittemore, 1998) were used: jaw champing, frothing at the mouth, erect ears and immobilization. A total of 24 gilts resulted positive (in oestrus; O gilts) and were individually identified. The same number (24) of gilts that resulted clearly negative (no oestrus; NO gilts) were also individually identified. Therefore, a total of 48 gilts were used to study the effect of the presence of oestrus at slaughter on carcass and meat quality. The feeding program was common to all of the animals and was based on wheat, barley, maize and soybean meal. A growing diet from 20 to 70 kg BW (2,210 kcal NE kg<sup>-1</sup>, 16.9% CP and 1.00% lys) and a finishing diet from 70 kg BW to slaughter (2,340 kcal NE kg<sup>-1</sup>, 14.4% CP and 0.86% lys) were provided.

When animals achieved approximately 130 kg BW, they were starved for 7 h and, on the same day, trans-

ported 130 km to a commercial abattoir plant (Jamones y Embutidos Alto Mijares, S.L., Teruel, Spain). At the abattoir, pigs were rested for 10 h with access to water but not to feed. Hot carcass weight was individually recorded because it is the first criterion used to identify those carcasses suitable for the production of Teruel ham. Therefore, carcasses that weighed less than 84 kg are not acceptable. At 45 min *postmortem*, pH and temperature in the *semimembranosus* muscle using a pH-meter equipped with a glass electrode (model 52-00, Crison Instruments S.A., Barcelona, Spain), carcass length from the posterior edge of the symphysis pubis to the anterior edge of the first rib, ham length from the anterior edge of the symphysis pubis to the hock joint, and ham circumference at its widest point were measured on the left side of each carcass. In addition, fat depth over the m. GM, and backfat thickness between the 3<sup>rd</sup> and 4<sup>th</sup> last ribs on the midline of the carcass (skin included) were measured, with a ruler with a precision of 0.5 mm, because it is the second criterion used to identify those carcasses suitable for Teruel ham production; carcasses with fat depth over the m. GM less than 16 mm are not acceptable. After 2 h at 2°C, carcasses were processed according to the simplified EC reference method (Branscheid *et al.*, 1990). Loins, hams, and shoulders were then trimmed of external fat and weighed to calculate trimmed loin, ham, and shoulder yield. Ham weight is the third criterion used to identify those carcasses suitable for Teruel ham production; carcasses with fresh ham weight less than 11.3 kg are not acceptable. At 24 h *postmortem*, pH was measured as described before and hams were weighed again to calculate ham shrink losses. The incidence of pale, soft and exudative meat (PSE, as % carcasses with pH at 45 min < 5.8) and of dark, firm and dry meat (DFD, as % carcasses with pH at 24 h > 6.2) were calculated.

A section of 300 ± 20 g of *longissimus thoracis* muscle was excised at the level of the last rib from each left loin of carcass. Meat samples were stored in individual plastic bags and vacuum-packaged at -20°C until subsequent analyses. Thawing and drip losses were determined as described by Latorre *et al.* (2003). Meat colour was evaluated with a chromameter (model CM 2002, Minolta Camera, Osaka, Japan) using objective measurements (CIE, 1976). Lightness (L\*), redness (a\*) and yellowness (b\*) were recorded and chroma (C\*) and hue angle (H\*) were calculated as C\* = √(a\*<sup>2</sup> + b\*<sup>2</sup>) and as H\* = tan<sup>-1</sup>(b\*/a\*) × 57.29, respectively. Chroma is related to the quantity of pigments and high values represent a more vivid color denoting lack of grayness

while Hue is the attribute of a color perception denoted by blue, green, yellow, red, purple, etc. related to the state of pigments (Wyszecki and Stiles, 1982). Cooking losses were determined by the method described by Honikel (1998) using a water bath (Precisterm, J.P. Selecta S.A., Barcelona, Spain) and a handheld temperature probe (model HI 9063, Hanna Instruments, Woonsocket, RI). Afterwards, tenderness was determined by a Warner-Bratzler device attached to an Instron Universal testing machine attached to a computer (Instron model 5543, Instron Ltd, Buckinghamshire, UK). Data were analyzed using the GLM procedure of SAS v. 6.12 (SAS, 1990). The treatment (presence or not of oestrus at slaughter) was included at the model as main effect. Each treatment was replicated 24 times. The incidence of PSE and DFD meat were analyzed by a  $\chi^2$  test.  $p$  values lower than 0.05 were classified as a significant difference and  $p$  values between 0.05 and 0.10 were classified as a trend.

A total of 857 pigs were observed at farm, 54.4% being barrows and 45.6% gilts. From all the females, a total of 24 manifested oestrus which is 6.14% of the total gilts. The proportion of O gilts was lower than expected. The reason could be that only females with clear oestrus symptoms were identified, not taking into account those that were at the beginning or at the end of the oestrus period, or delays in the puberty and silent oestrus (without clear symptoms) (Tummaruk *et al.*, 2007).

Carcass parameters (weight, covering fat depth and yield of primal lean cuts) were similar to those reported by Latorre *et al.* (2009a) for similar gilts slaughtered at 130 kg BW and also intended for dry-cured ham (Table 1). The presence of oestrus at slaughter did not modify any carcass characteristics of gilts. However, treatment affected the percentage of carcasses suitable for Teruel ham (Table 2). No effect of the presence of oestrus at slaughter was detected on the proportion of suitable carcasses with the minimum carcass and ham weights required for PDO Teruel ham. In fact, 100% of the carcasses fulfilled those requirements. However, NO gilts tended to show a higher proportion of final carcasses suitable for Teruel ham (carcasses that fulfill the three named requirements) than O gilts (91.6 vs. 84.4%;  $p = 0.09$ ) because of lack of backfat thickness at m. GM (100 vs. 87.5%;  $p = 0.09$  for NO and O gilts, respectively). This might be due to a reduction in feed intake during oestrus days as a consequence of stress. Some authors have mentioned that oestrus could lead to a lower feed consumption (Zeng *et al.*, 2002; Serrano *et al.*, 2008) but it has been not scientifically demonstrated. Le Bellego *et al.* (2002) reported a reduction in daily feed intake and in backfat thickness in growing pigs under a stress situation; housed at a high (29°C) vs. a thermoneutral (22°C) temperature. However, these effects could be due to the hot temperature and/or to the stress caused by hot temperature. In fact, Morrison *et al.* (2003) detected a reduced backfat depth under

**Table 1.** The effect of oestrus at slaughter on carcass characteristics of gilts intended for dry-cured Teruel ham

|   | Oestrus | No oestrus | SE <sup>a</sup><br>(n=24) | p <sup>b</sup> |
|---|---------|------------|---------------------------|----------------|
| Carcass weight, kg  | 103.9   | 103.2      | 2.05                      | NS             |
| Backfat depth at 3 <sup>rd</sup> -4 <sup>th</sup> last ribs, mm | 26.4    | 26.5       | 1.01                      | NS             |
| Fat depth at m. <i>gluteus medius</i> , mm                      | 23.7    | 23.3       | 0.97                      | NS             |
| Carcass length, cm  | 85.7    | 86.1       | 0.89                      | NS             |
| Ham length, cm  | 38.4    | 38.2       | 0.28                      | NS             |
| Ham perimeter, cm   | 76.8    | 77.0       | 0.58                      | NS             |
| Weight of trimmed lean cuts, kg                                 |         |            |                           |                |
| Shoulder  | 14.8    | 14.7       | 0.311                     | NS             |
| Loin  | 6.6     | 6.6        | 0.147                     | NS             |
| Ham   | 26.9    | 26.8       | 0.511                     | NS             |
| Total <sup>c</sup>  | 48.3    | 48.0       | 0.916                     | NS             |
| Yield of trimmed lean cuts, % carcass                           |         |            |                           |                |
| Shoulder  | 14.2    | 14.2       | 0.188                     | NS             |
| Loin  | 6.4     | 6.4        | 0.098                     | NS             |
| Ham   | 25.9    | 26.0       | 0.219                     | NS             |
| Total <sup>c</sup>  | 46.5    | 46.6       | 0.410                     | NS             |
| Ham shrink losses, %  | 1.01    | 1.05       | 0.046                     | NS             |

<sup>a</sup> SE: standard error of the mean. <sup>b</sup> p: significance. NS: not significant ( $p > 0.05$ ). <sup>c</sup> Shoulder + Loin + Ham.

**Table 2.** The effect of oestrus at slaughter in gilts on suitable (%) carcasses intended for dry-cured Teruel ham

| Requirements for Teruel ham <sup>a</sup>      | Oestrus | No oestrus | SE <sup>b</sup><br>(n=24) | p <sup>c</sup> |
|---|---------|------------|---------------------------|----------------|
| Carcass weight ≥ 84 kg                        | 100     | 100        | —                         | NS             |
| Fat depth at m. <i>gluteus medius</i> ≥ 16 mm | 87.5    | 100        | 0.05                      | +              |
| Fresh ham weight ≥ 11.3 kg                    | 100     | 100        | —                         | NS             |
| All <sup>d</sup>                              | 87.5    | 100        | 0.05                      | +              |

<sup>a</sup> According to BOA (1993). <sup>b</sup> SE: standard error of the mean. <sup>c</sup> p: significance. NS: not significant ( $p > 0.05$ ). +:  $p < 0.10$ .

<sup>d</sup> Carcasses fulfill all the above requirements.

other stress situation (a reduced pen space) but the authors concluded that it was not associated with changes in feeding behaviour but with social stress. Irrespective of the oestrus presence at slaughter, 87.5% of the gilts studied were suitable for Teruel ham. Similar results were reported by Latorre *et al.* (2008) who found that 84.2% of gilts slaughtered at 130 kg BW were suitable.

The influence of the presence of oestrus at slaughter on meat characteristics of gilts was scarce (Table 3). Meat parameters (color, water holding capacity and tenderness) were similar to those showed by Latorre *et al.* (2009b) for similar gilts slaughtered at 130 kg BW and also intended for dry-cured ham. Possible differences in meat between O and NO gilts were expected for pH, temperature or DFD/PSE meat incidence because oestrus is stressful to the gilt. A marked increase in social activities was demonstrated in gilts in

oestrus which includes snout contacts between pigs, ano-genital sniffing, flank nosing and mounting (Pedersen, 2007). Meat from O gilts tended to show higher lightness ( $p = 0.08$ ) and had a yellower ( $p < 0.01$ ) and more intense ( $p < 0.01$ ) colour than meat from NO gilts. This could be related to stress. It has been widely reported that a stress situation just before slaughter results in low pH values at 45 min *postmortem* and higher lightness and moisture losses (Warris, 2000). However, no named variable was affected by treatment. In fact, although O gilts showed a numerically higher incidence of PSE and DFD meat than NO gilts (20.8 vs 18.2% for PSE and 16.6 vs 9.0% for DFD), the differences were not significant ( $p > 0.05$ ). Also, water holding capacity was not affected by the presence of oestrus. Perhaps, a higher number of replicates per treatment might have permitted the detection of significant

**Table 3.** The effect of oestrus at slaughter on meat characteristics of gilts intended for dry-cured Teruel ham

|  | Oestrus | No oestrus | SE <sup>a</sup><br>(n=24) | p <sup>b</sup> |
|--|---------|------------|---------------------------|----------------|
| <i>Measurements postmortem at m. Semimembranosus</i> |         |            |                           |                |
| pH 45 min  | 5.98    | 6.04       | 0.061                     | NS             |
| pH 24 h  | 5.90    | 5.97       | 0.047                     | NS             |
| Temperature 45 min, °C                               | 37.6    | 37.3       | 0.17                      | NS             |
| <i>Measurements at m. Longissimus thoracis</i>       |         |            |                           |                |
| Color  |         |            |                           |                |
| L*   | 48.2    | 46.6       | 0.622                     | +              |
| a*   | 1.04    | 1.05       | 0.176                     | NS             |
| b*   | 5.33    | 2.27       | 0.277                     | **             |
| C*   | 5.51    | 4.47       | 0.265                     | **             |
| H*   | 78.4    | 66.7       | 5.532                     | NS             |
| Water holding capacity, %                            |         |            |                           |                |
| Drip losses  | 0.83    | 0.89       | 0.088                     | NS             |
| Thawing losses                                       | 7.36    | 6.90       | 0.449                     | NS             |
| Cooking losses                                       | 20.05   | 19.30      | 0.697                     | NS             |
| Total losses   | 28.9    | 27.0       | 1.091                     | NS             |
| Shear force, kg                                      | 3.00    | 2.80       | 0.129                     | NS             |

<sup>a</sup> SE: standard error of the mean. <sup>b</sup> p: significance. NS: not significant ( $p > 0.05$ ). +:  $p < 0.10$ . \*\*:  $p < 0.01$ .

differences in some of these traits. In addition, a sensorial analysis would be interesting because possible differences in flavour might be found. However, the «sexual odour in females» is only a hypothesis and it has not been demonstrated.

Under our experimental conditions, it may be concluded that a low proportion of gilts were in oestrus when females were slaughtered at 130 kg BW, although the result might have been underestimated. The presence of oestrus in gilts at slaughter had scarce effect on carcass and meat characteristics. It did not affect the proportion of PSE/DFD meat but increased yellowness and intensity in meat colour. It also tended to reduce the proportion of suitable carcasses intended for PDO Teruel ham production because of a lack of backfat depth. However, new trials (with a higher proportion of replicates and also a sensorial analysis) would be needed before recommending the industry to avoid the slaughter of gilts in oestrus if a minimum covering fat is needed to improve the quality of the end product.

## Acknowledgements

This research was supported by the Ministry of Science and Technology (CDTI Project 20060926). The authors thank L. Ariño and staff of the slaughterhouse for help with sample collection.

## References

- BOA, 1993. Orden del Departamento de Agricultura, Ganadería y Montes por la que se aprueba el Reglamento de la Denominación de Origen Jamón de Teruel y su Consejo Regulador. Boletín Oficial de Aragón 29/07/1993. [In Spanish].
- BRANSCHEID W., DOBROWOLSKI A., SACK E., 1990. Simplification of the EC-reference method for the full dissection of pig carcasses. Fleischwirtsch 70, 565-567.
- CIE, 1976. Colorimetry: Official Recommendations of the International Commission on Illumination. Publication CIE No. 15 (E-1.3.1). Commission International de l'Eclairage, Paris.
- CRPDO JAMÓN DE TERUEL, 2010. Data base. Consejo Regulador of the Protected Designation of Origin of Teruel ham. Available in <http://www.jamondeteruel.com> [25 January, 2010]. [In Spanish].
- HONIKEL K.O., 1998. Reference methods for the assessment of physical characteristics of meat. Meat Sci 49,447-457.
- LATORRE M.A., MEDEL P., FUENTETAJA A., LÁZARO R., MATEOS G.G., 2003. Effect of gender, terminal sire and age at slaughter on performance, carcass characteristics and meat quality of heavy pigs. Anim Sci 77, 33-45.
- LATORRE M.A., GARCÍA-BELENGUER E., ARIÑO L., 2008. The effects of gender and slaughter weight on growth performance and carcass traits of pigs intended for dry-cured hams from Teruel (Spain). J Anim Sci 86, 1933-1942.
- LATORRE M.A., RIPOLL G., GARCÍA-BELENGUER E., ARIÑO L., 2009a. The increase of slaughter weight in gilts as strategy to optimize the production of Spanish high quality dry-cured ham. J Anim Sci, 87, 1464-1671.
- LATORRE M.A., RIPOLL G., GARCÍA-BELENGUER E., ARIÑO L., 2009b. The effect of gender and slaughter weight on loin and fat characteristics of pigs intended for Teruel dry-cured ham production. Span J Agric Res 7, 407-416.
- LE-BELLEGO L., VAN MILGEN J., NOBLET J., 2002. Effect of high temperature and low-protein diets on the performance of growing-finishing pigs. J Anim Sci 80, 691-701.
- MARM, 2010. Anuario de Estadística Agraria. Data base. Ministerio de Medio Ambiente y Medio Rural y Marino. Available in <http://www.marm.es>. [17 January, 2010]. [In Spanish].
- MORRISON R.S., HEMSWORTH P.H., CRONIN G.M., CAMPBELL R.G., 2003. The effect of restricting pen space and feeder availability on the behaviour and growth performance of entire male growing pigs in a deep-litter, large group housing system. Appl Anim Behav Sci 83, 163-176.
- PEDERSEN L.J., 2007. Sexual behavior in female pigs. Hormones Behav 52, 64-69.
- SAS, 1990. SAS Institute user's guide: statistics. Statistical Analysis Systems. Version 6, 4<sup>th</sup> ed. Cary, NC, USA.
- SERRANO M., VALENCIA D.G., NIETO M., LÁZARO R., MATEOS G.G., 2008. Influence of sex and terminal sire line on performance and carcass and meat quality of Iberian pigs reared under intensive production systems. Meat Sci 78, 420-428.
- TUMMARUK P., TANTASUPARUK W., TECHAKUMPHU M., KUNAVONGKRIT A., 2007. Age, body weight and thickness at first observed in crossbred Landrace × Yorkshire gilts, seasonal variations and their influence on their subsequent performance. Anim Reprod Sci 99, 167-181.
- WARRIS P.D., 2000. Meat science: an introductory text. CABI Pub Inc, NY, USA. 310 pp.
- WHITTEMORE C., 1998. Ciencia y práctica de la producción porcina. Ed Acritiba. 800 pp. [In Spanish].
- WYSZCECKI G., STILES W.S., 1982. Color science. Concepts and methods: quantitative data and formula, 2<sup>nd</sup> ed. John Wiley, NY.
- ZENG X.Y., TURKSTRA J.A., TSIGOS A., MELOEN R.H., LIU X.Y., CEN, F.Q., SCHAAPEL W.M.M., OONK H.B., GUO D.Z., VAN DE WIEL D.F.M., 2002. Effects of active immunization against GnRH on serum LH, inhibin A, sexual development and growth rate in Chinese female pigs. Theriogenology 58, 1315-1326.